BBA 86127

Electron transfers in chemistry and biology

R.A. Marcus ^a and Norman Sutin ^b

^a Arthur Amos Noyes Laboratory of Chemical Physics, California Institute of Technology, Pasadena, CA 91125, and
^b Department of Chemistry, Brookhaven National Laboratory, Upton, NY 11973 (U.S.A.)

(Received October 1st, 1984)

II. Electron-transfer reactions in solution. Theory 2 A. Classical theory. 2 B. Quantum-mechanical aspects. 2 C. Electronic and steric aspects. 2 D. Nuclear and electron tunneling. 2 E. Relaxation-limited rates. 2 III. Comparisons of predictions with experimental measurements on non-biological systems. 2 A. Cross-reactions. 2 B. Self-exchange reactions. 2 C. Relation between rates and equilibrium constants. 2 D. Mixed-valence systems. 2 E. Electrode reactions. 2 IV. Extension to biological electron transfers. 2 A. Similarities and differences with small molecule electron transfers. 2 B. Reorganization parameters. 2 C. Electron tunneling barriers. Distance dependence of electron-transfer rate in nonbiological systems. 2 D. Work terms for bimolecular reactions involving metalloproteins. 2 V. Applications to biological electron transfers. 2 A. Reactions of cytochrome c with metal complexes. 2 B. The cytochrome c with metal complexes. 2 B. The cytochrome c self-exchange reaction. 2 <			
I.	Introduction	266	
II.	Electron-transfer reactions in solution. Theory	266	
	A. Classical theory.	266	
		272	
	C. Electronic and steric aspects.	275	
		275	
	E. Relaxation-limited rates.	276	
Ш	. Comparisons of predictions with experimental measurements on non-biological systems.	276	
		276	
	B. Self-exchange reactions.	278	
	C. Relation between rates and equilibrium constants.	280	
		281	
	E. Electrode reactions	283	
IV.		283	
	A. Similarities and differences with small molecule electron transfers.	283	
	B. Reorganization parameters.	284	
	C. Electron tunneling barriers. Distance dependence of electron-transfer rate in nonbiological systems.	286	
	D. Work terms for bimolecular reactions involving metalloproteins.	288	
v.	Applications to biological electron transfers.	291	
		291	
	B. The cytochrome c self-exchange reaction.	292	
	C. Intramolecular electron transfer.	296	
	D. Photosynthetic reaction centers.	298	
	E. Reactions of cytochrome c with cytochrome b_5 and cytochrome c oxidase	301	
	F. Reactions of blue copper proteins.	303	
	G. Cytochrome c peroxidase	306	
VI.	Conclusions.	308	
Ap	pendix I. Expressions for activation- and diffusion-controlled reaction rates. A. No steric effects.	308 308	
Ab	breviations: Cyt c, cytochrome c; cp, cyclopentadienyl radi-		

Wentzel-Kramers-Brillouin; CcP, cytochrome c peroxidase;

EXAFS, extended X-ray-absorption fine structure.

cal; hfac, hexafluoroacetylacetonate ion; bpy, 2,2'-bipyridine; phen, 1,10-phenanthroline, (FC), Franck-Condon factor; WKB,

B. Steric effects included	310
C. Dependence of H_{AB} on nuclear configuration	311
D. Miscellaneous.	313
Appendix II. Franck-Condon factors and absolute rates.	314
Acknowledgements	316
References	316

I. Introduction

considerable experimental study during the past three decades. Experimental results have also been obtained on related phenomena, such as reactions between ions or molecules and electrodes, chargetransfer spectra, photoelectric emission spectra of ionic solutions, chemiluminescent electron transfers, electron transfer through frozen media, and electron transfer through thin hydrocarbon-like films on electrodes. For 'outer-sphere' electron-transfer reactions in solution there have also been extensive develop-

Electron-transfer reactions between ions and

molecules in solution have been the subject of

ments in the theory. Quantitative predictions of the theory, without adjustable parameters, include [1-3] (1) rate constants of 'cross-reactions' from those of self-exchange reactions, (2) the relation between the rate constant of a reaction and the driving force for the reaction (standard free energy of reaction ΔG^0), (3) nonspecific solvent effects on the rate constant, (4) the relation between rates of self-exchange reactions in solution and rate of electron transfer between one of the reactants and an electrode, (5) the calculation of rate constants of electron-transfer reactions from molecular size, bond length changes, vibration frequencies, and solvent properties, and (6) the 'inverted' effect.

By and large, as shown later in this review, these predictions have been successful for a large body of experimental data. In some cases, cited later, there have been major deviations, which have sometimes been used as diagnostic mechanistic tools. In some cases, too, there have been trends in relatively minor discrepancies and these, too, are discussed.

Some of these theoretical predictions (notably those arising from the dependence of the activa-

tion-free energy barrier on 'intrinsic' (λ) and thermodynamic (ΔG^0) factors) have been extended to other quite different types of reactions, such as atom and proton transfers [4-6], group transfers [7,9], hydride transfers [10,11], proton-bound dimers [12], gas phase ion-molecule reactions [13,14] and conformational equilibria [15]. The main questions, however, for the present article, are: (1) what predictions for reactions in solution can be extended to electron transfers in biological systems, and (2) what can the theory say about the interpretation of some current experimental results? Both of these questions will be addressed.

310

We begin with a review of the concepts and equations for electron transfers in solution.

II. Electron-transfer reactions in solution. Theory

IIA. Classical theory

The theory of electron-transfer reactions in solution has been summarized and reviewed in many texts and articles, e.g., Refs. 16-27. In the present section we briefly summarize the relevant concepts and equations, so as to serve as a basis for the comparison with experiment and for their extension to biological systems.

We shall be concerned in this section primarily with what has been termed outer-sphere bimolecular electron-transfer reactions, i.e., with reactions in which the two reactants do not share a common atom or group, or, more generally, reactions in which the interaction of the relevant electronic orbitals of the two centers is weak. This circumstance greatly simplifies the theoretical development.

For reaction to occur, the reactants must approach each other to enhance the coupling of their electronic orbitals, and at the same time there must be fluctuations in other coordinates. As in any reaction the system as a whole proceeds from configurations (coordinates) in the vicinity of the equilibrium ones for the reactants to those for the products. In the present case, these coordinates involve vibrational coordinates of the reactants, when their equilibrium values differ before and after electron transfer, as well as the orientational coordinates of the surrounding solvent molecules: averaged equilibrium orientations of those solvent molecules differ for reactants and products because each reactant undergoes a change of charge.

The potential energy of the reactants and surrounding medium is a function of all of these nuclear coordinates (thousands of them), and defines a many-dimensional potential-energy surface. A one-dimensional profile of such a many-dimensional surface is given schematically in Fig. 1, and is labelled R. The profile for the products and surrounding medium is labelled P. If there are N coordinates the intersection point in Fig. 1 represents an (N-1)-dimensional surface in the N-dimensional coordinate space \P .

Because the electron being transferred is such a light particle, the Franck-Condon principle § can be applied to this weak overlap electron transfer (small overlap of the electronic orbitals). It then follows that transfer will occur only at or near nuclear configurations for which the total potential energy of the reactants and surrounding medium is equal to that of the products and




Fig. 1. Profile of many-dimensional potential surface of reactants plus surrounding medium (R) and that of products plus medium (P). Dotted lines indicate splitting due to electronic interaction of reactants. A and C denote nuclear coordinates for an equilibrium configuration of reactants and for products, respectively; B denotes nuclear configuration at the intersection of the two potential-energy surfaces.

surrounding medium, i.e., at the intersection in Fig. 1. Thermal fluctuations of the coordinates from values near the minimum of the R surface are needed in order for the reacting system to reach the intersection region in Fig. 1.


Fluctuations in the vibrational coordinates need be considered when a reactant has different equilibrium bond lengths (or bond angles) in its two redox states. The systems studied vary widely in this respect, as we shall see in Section III. In addition, fluctuations in the orientational coordinates of the solvent molecules are needed when the solvent is polar. In the latter case the mean orientation of each solvent molecule at any position near the reacting pair is significantly different for reactants and products, as already noted.

Once the system reaches the intersection of the two curves in Fig. 1 the probability of going from the R to the P surface depends on a number of factors, such as the extent of coupling of the electronic orbitals of the two reactants, which in turn depends on the separation distance r of the reactants or, more precisely, on the dominant range of separation distances contributing to the reaction (cf. discussion later).

It is useful at this point to consider the effective potential energy curve for the *electronic* motion, and relate it to plots such as the one in Fig. 1 for the *nuclear* motion. A series of such curves for

The actual profile of the many-dimensional surface is more complicated than Fig. 1 when there are solvation contributions – there are many local potential-energy minima, for example. In that case only the profile of the *free energy* (in effect, a Boltzmann-weighted profile) of a system consisting of the pair of reactants, at a given separation, and their environment, along the reaction coordinate could have a quadratic appearance. Such an approximation is much less drastic than assuming that the potential energy is a quadratic function of the coordinates.

[§] According to the Franck-Condon principle, during the actual (instantaneous) electron transfer the nuclei do not have time to change either their positions or their momenta. To satisfy reasonably both conditions simultaneously, the transfer has to occur at nuclear configurations stated in this paragraph, namely at or near the intersection region in Fig. 1. The entropy of the reactants (plus environment) and that of the products (plus environment) are equal on the intersection surface, as are their energies and hence, also, their free energies [2].



corresponding (nuclear) potential-energy curves on the right. The electronic energies in the two wells are indicated on the left for three nuclear configurations A, B, and C for the reaction $\operatorname{red}_1 + \operatorname{ox}_2 \to \operatorname{ox}_1 + \operatorname{red}_2$. The levels in the wells are vertical ionization energies: the filled and open circles denote, respectively, ionization of the reduced state at its equilibrium nuclear configuration and at the equilibrium configuration appropriate to its oxidized state; the half-filled circles refer to ionization of the reduced state at the nuclear configuration appropriate to the intersection region. The level in well 1 is initially occupied in (a) and the level in well 2 in (c).

Fig. 2. Electronic potential-energy curves (on left) and the

different nuclear configurations (A, B, and C in Fig. 1) is shown in Fig. 2a-c. The effective one-electron potential-energy curve with two wells is given in Fig. 2a. Also given are two electronicenergy levels of the entire system, one when an electronic orbital of reactant 1 is occupied, and the other when an orbital of reactant 2 is occupied. Initially, in a reaction between the reduced form of reactant 1 and the oxidized form of reactant 2, an electronic orbital of reactant 1 is occupied. The nuclear configuration is labelled A in Fig. 1 (and in Fig. 2a), and is a configuration that is an equilibrium one for the reactants ('red₁' and 'ox₂'). The relation of these two electronic-energy levels in Fig. 1 to the nuclear potential-energy curves is indicated by the dashed lines in Fig. 2a.

After a suitable fluctuation of positions of the nuclei in the system prior to electron transfer, a nuclear configuration such as B in Fig. 1 is formed

and the positions of the two electronic-energy levels are indicated on the left-side of Fig. 2b; that of the electronic orbital in well 1 has been raised and that of well 2 has been lowered. The relationship of these levels to Fig. 1 is again indicated by dashed lines in Fig. 2b. Now the electron can transfer from well 1 to well 2.

A further fluctuation in the positions of the nuclei in the entire system to form a set of nuclear configurations which is an equilibrium one for the products leads to Fig. 2c. The electronic energy when well 2 is occupied now is the lower one. The relation of these energy levels to Fig. 1 is again indicated by dashed lines in Fig. 2c.

We shall denote by $\kappa(r)$ the transmission coefficient or averaged transition probability for electron transfer per passage of the system through the intersection region of Fig. 1, at a fixed separation distance r of the two reactants. This $\kappa(r)$, discussed in Section IV, is usually assumed to vary at large r approximately as $\exp(-\beta r)$. When suitably averaged over r, $\kappa(r)$ will be denoted by κ . For a reaction in which there is substantial electronic coupling between the reactants ('adiabatic reactions'), κ is about unity [25].

The calculation of the rate constant of an electron-transfer reaction involves calculating the probability of reaching the many-dimensional intersection region, due to the fluctuations described above, multiplied by a suitably weighted frequency for crossing the intersection region and by a transition probability for going from the R to the P surface. The net result of this calculation is that the rate constant k of a bimolecular electron-transfer reaction is given by

$$k = \kappa A \sigma^2 \exp(-\Delta G^*/RT), \tag{1}$$

where $A\sigma^2$ has dimensions of collision frequency (its value is given in Appendix I), σ is the average center-to-center distance in the reacting pair during the electron transfer (the mean separation distance in the transition state of the reaction), ΔG^* is the free energy of activation that is related to λ (described below), to ΔG^0 , the 'standard' free en-

[¶] Evidence that in some systems the distance dependence may be somewhat less than exponential is given in Ref. [28].

the work of bringing the reactants (w^r) or products (w^p) to the mean separation distance σ [3,27]:

ergy of reaction in the prevailing medium, and to

$$\Delta G^* = w^{\mathsf{r}} + \frac{\lambda}{4} \left(1 + \Delta G^{0} / \lambda \right)^2 \tag{2}$$

Molecular theory (specifically the use of molec-

ular vibrational coordinates) was used to calculate

$$\Delta G^{0\prime} = \Delta G^0 + w^p - w^r \tag{3}$$

the contribution λ_i of changes in the bond lengths of the reactants to λ . To obtain a simple expression for the contribution λ_o of changes in solvent orientation coordinates to λ , dielectric continuum theory was used. (A statistical mechanical treatment is also given in Ref. 3.) In this way λ was found to be given by [3,27]:

$$\lambda = \lambda_i + \lambda_o \tag{4}$$

$$\lambda_i = \Sigma_j \frac{f_j^{\text{r}} f_j^{\text{p}}}{f_j^{\text{r}} + f_j^{\text{p}}} (\Delta q_j)^2$$
 (5)

$$\lambda_{o} = (\Delta e)^{2} \left[\frac{1}{2a_{1}} + \frac{1}{2a_{2}} - \frac{1}{r} \right] \left[\frac{1}{D_{op}} - \frac{1}{D_{s}} \right]$$
 (6)

where $f_i^{\rm r}$ is the jth normal mode force constant in

the reactants. f_j^p is that in the products, Δq_j is the change in equilibrium value of the *j*th normal coordinate, Δe is the charge transferred from one reactant to the other, a_1 and a_2 are the radii of the two (spherical) reactants, r is the center-to-center separation distance (r equals the σ in Eqn. 1), and D_s and D_{op} are the static and optical (square of refractive index) dielectric constants of

In the derivation of Eqns. 1-6, the vibrations

¶ The ΔG^0 introduced here is obtained from concentration

measurements in the prevailing medium and in Refs. 3 and 27 is shown to be the appropriate quantity to be used in

the solvent, respectively.

within the reactants were treated as harmonic oscillators, but no such restrictive assumption was made for the solvent outside of the coordination

shells. What was assumed was, rather, that the dielectric polarization outside the coordination shells responded linearly to any change in (real or

fictitious) charges, and hence that free energy depended quadratically on charging parameters ('dielectric unsaturation approximation') [1-3] §. Such a treatment allows the individual solvent dipoles to move very anharmonically, as indeed

they do in a liquid. Equations 1-6 can be given a physical interpretation discussed elsewhere [26]. For example, when $D_{\rm s}$ equals $D_{\rm op}$, the solvent molecules are nonpolar and $\lambda_{\rm o}$ vanishes. Again, the larger the radii of the

reactants, the smaller the charge-solvent interac-

tions and the smaller λ_0 .

The 'cross-relation' [2,3,27] follows from these equations, by noting from Eqns. 4-6 that λ for the cross-reaction (λ_{12}), Eqn. 7,

$$\operatorname{red}_{1} + \operatorname{ox}_{2} \xrightarrow{k_{12}} \operatorname{ox}_{1} + \operatorname{red}_{2} \tag{7}$$

is approximately the mean of the λ 's of the self-exchange reactions λ_{11} and λ_{22} .

$$\operatorname{red}_{1} + \operatorname{ox}_{1} \xrightarrow{k_{11}} \operatorname{ox}_{1} + \operatorname{red}_{1}$$

$$k_{22}$$
(8)

$$\operatorname{red}_{2} + \operatorname{ox}_{2} \xrightarrow{k_{22}} \operatorname{ox}_{2} + \operatorname{red}_{2} \tag{9}$$

From this result one finds when the transmission

coefficients κ are approximately unity, and when the self-exchange rate constants k_{ii} are either corrected for the work terms or when the latter approximately cancel, that k_{12} is given by Eqn. 10:

$$k_{12} = \left(k_{11}k_{22}K_{12}f_{12}\right)^{1/2} \tag{10}$$

Eqns. 1-3. It is slightly different from the standard free energy of reaction which is based on standard states. The present ΔG^0 was denoted by a primed quantity in Refs. 3 and 27 but we use a simpler notation here and use a primed quantity for the work-corrected free energy of reaction, defined in Eqn. 3. The ΔG^{0} in Eqn. 3 equals the free energy of reaction when the reactants are a distance r apart in the prevailing medium.

As indicated earlier, such an assumption about the free energy, a thermodynamic quantity which involves statistical averaging over many coordinates, is much less drastic than assuming that the potential-energy function for the surrounding solvent molecules is a quadratic function of their coordinates.

7 in the prevailing medium, i.e., K_{12} is equal to $\exp(-\Delta G^0/RT)$, and f_{12} is a known function of k_{11}, k_{22} and $K_{12} (\ln f_{12} = \frac{1}{4} (\ln K_{12})^2 / \ln(k_{11}k_{22}/Z^2)$, where $Z = A\sigma_{11}\sigma_{22}$; in practice f_{12} is often close to unity). A modification of this equation when the κ 's are less than unity has been given [21].

where K_{12} is the equilibrium constant of reaction

Section III, including comparison of the cross-relation Eqn. 10 with the experimental data, ab initio calculation of rate constants, study of nonspecific

Several applications of Eqns. 1–6 are given in

solvent effects, and relation between rates and equilibrium constants. We have already noted that a range of separation distances r contributes to the rate con-

stant. To make $\kappa(r)$ large, r should be small; on

the other hand, any short-range or Coulombic

repulsion between the reactants will favor large r.

A compromise in r results, and we obtain approximately Eqn. 1 (cf. Ref. 3 and Appendix I).

More refined rate expressions can also be derived.

Expressions have also been derived for the outer-sphere electron transfer between an ion or molecule and an electrode [27,29-31]. The expressions are similar to Eqns. (1-6), only now $A\sigma^2$ is replaced by a related quantity which has dimensions of a collision frequency of the ion with unit area of the electrode ($\approx 1 \cdot 10^4 \,\mathrm{cm} \cdot \mathrm{s}^{-1}$), r is now twice the distance of the center of the ion to the electrode surface, $1/2a_2$ is missing from Eqn. 6 (a_2) tends to infinity), the summation in Eqn. 5 is over the vibrational coordinates of one ion instead of two, and ΔG^0 is replaced by the activation overpotential. When the rate constant for self-exchange in solution k_{ex} is then compared with the rate constant for reaction between one of these ions and an electrode at zero activation overpotential, $k_{\rm el}$, one finds from Eqns. 1-6 and from the

$$\left(\frac{k_{\rm ex}}{Z_{\rm soln}}\right)^{1/2} \gtrsim \frac{k_{\rm el}}{Z_{\rm el}} \tag{11}$$

equations for the corresponding reaction at an

electrode that, approximately,

where Z_{soln} denotes $A\sigma^2$ for the solution reaction and $Z_{\rm el}$ the corresponding quantity for the electrode reaction. Equality exists when the averaged

center-to-center separation distance σ of the reactants in the homogeneous reaction is twice that from the center of the reactant to the electrode. However, an adsorbed solvent layer on the electrode may cause the latter separation distance to be somewhat larger. The effect of this increased ron $k_{\rm el}$ causes the inequality in Eqn. 11, and the effect can be calculated. The equations are applied to electrode systems in Section III.

In deriving the preceding formulae the motion of the nuclei was treated classically. In practice there may be some nuclear tunneling through the barrier in Fig. 1, e.g., depicted schematically by a horizontal motion from a to b in Fig. 3. The nuclear tunneling in Fig. 3 at room temperature typically accounts, in the so-called normal region. for only a relatively minor factor (Γ in Table II) in the rate constants, which due to variations in λ and ΔG^0 , may vary by some 15 or more orders of magnitude from system to system. Examples are given later in Table II. The cross-relation in the normal region is even less affected, because of partial cancellation of the quantum corrections in the ratio $k_{12}/(k_{11}k_{22}K_{12})^{1/2}$. Nuclear tunneling in the 'inverted region', defined below, has a larger

effect, as discussed later.

When $\Delta G^{0\prime}$ is made increasingly negative at constant λ , for example, by suitable variation of a ligand in a reactant, one sees from Eqn. 2 that the free-energy barrier to reaction ΔG^* initially decreases and that, when $-\Delta G^0$ exceeds λ , ΔG^* begins to increase. This region where $-\Delta G^{0} > \lambda$ has been termed the inverted region [3], and some consequences are described in Section III. Physi-




Fig. 3. As in Fig. 1, but describing nuclear tunneling through the barrier from the R to the P surface (from a to b).

cally, in terms of Fig. 1, this effect of making ΔG^0 more negative at constant λ can be seen to correspond to lowering the P surface vertically (or raising the R surface). The intersection of the R and P surfaces eventually occurs at the minimum of the R surface and there is then no barrier. Further lowering raises the point of intersection, as in Fig. 4, and hence now raises the barrier, i.e., increases ΔG^* .

In the case of Fig. 4, nuclear tunneling at some energy, from the R to the P surface, is represented by a horizontal motion from a to b, as in Fig. 5. Here, the slopes of the potential-energy surfaces R and P have the same sign near the intersection region, unlike the case in Fig. 3, and the semiclassical theory of nuclear tunneling shows that tunneling is now much more important. Typically, nuclear tunneling is treated in the case of Figs. 3 and 5 either by calculation of quantum mechanical Franck-Condon factors (appearing in the next section) or by semiclassical nuclear tunneling equations. The theoretical equations for these quantum




Fig. 4. Highly exothermic reaction to form the electronic ground-state products (surface P) or products where one is electronically excited (surface P*).




Fig. 5. Nuclear tunneling in a highly exothermic reaction, from a to h

calculations are referred to in Section IIB. Examples of the results are given in Section III.

A corollary of Fig. 4 involves the formation of electronically excited products, i.e., a chemiluminescent reaction occurs: although the ground electronic-state potential-energy surface P for the products may intersect R at a high point, the surface in which one of the products is electronically excited, P*, may intersect in a more favorable region, as in Fig. 4, and then one may obtain preferential formation of the less thermodynamically favored products [33,34]. This preferential ($\approx 100\%$) formation of electronically excited products has been observed experimentally [35], and one explanation is the predicted 'inverted effect'. Direct experimental confirmation of the inverted effect has recently been obtained [36-38] (cf. Ref. 39 and also a discussion of other studies [27]).

Various other classical treatments of the problem have also been given, including the treatment

motion. It is small when the region of space occupied by the reactants' relevant wavefunction for the nuclear motion is rather different from that occupied by the products' wavefunction, for example, when there is a large difference of the equilibrium bond lengths and when, at the same time, the two wavefunctions refer to relatively unexcited vibrational quantum states. When the overlap of two such wavefunctions of the same energy is small, it can be shown that an equivalent semiclassical interpretation is that the states are connected not by a classically allowed path but by a nuclear tunneling through a barrier separating the reactants' and products' regions of vibrational coordinate space. (The argument is based on a semiclassical evaluation of the overlap integral, related to that given in Ref. 32.)

[¶] In most chemical reactions nuclear tunneling is usually important only for protons. However, for the present reactions, the potential-energy barrier for changes in metal-ligand bond lengths is so narrow (e.g., as seen from the Δd_0 's in Table II) that even ligands can tunnel at room temperature (e.g., see Γ factors in Table II).

[§] A particular Franck-Condon term in the Franck-Condon sum (FC) in Eqn. 16 of Section IIB is an overlap integral of a reactants' and a products' wavefunction for the nuclear

of vibrations and solvation [40] and of vibrations alone [41,42]. The expressions in Ref. 40 for the solvation are the same as those in Ref. 1[¶]. The vibrations were treated using calculated force fields rather than experimental vibration frequencies. The vibrational results in Refs. 41 and 42 are for self-exchange reactions, rather than cross-reactions, and a simple force field was assumed (no cross-terms in bond force constants).

In the foregoing discussion we have explicitly considered outer-sphere electron transfers. The latter differ from 'inner-sphere' electron transfers. In the latter, the two reactants approach each other closely enough that they actually share in common a ligand or atom in their coordination shells. This strong interaction is then followed by either an electron, atom, or radical transfer. Such bond formation and rupture requires a modified treatment from that leading to Eqns. 1-6. The cross-relations can still apply [4,9,10,44], provided the cross-reaction and the self-exchange reactions have a similar mechanism.

IIB. Quantum-mechanical aspects

treated classically. Quantum mechanical treatments of the nuclear coordinates have been made for cases where additional approximations have been imposed (e.g., 45-56). These treatments were almost exclusively for reactions having very weak electronic coupling, namely having $\kappa \ll 1$ in Eqn.

In Eqns. 1–6 the motion of the nuclei was

The first-order rate constant for electron transfer at fixed r is now given by

$$k = \frac{2\pi}{\hbar} H_{\rm AB}^2(FC),\tag{12}$$

where H_{AB} is the electronic matrix element describing the electronic coupling of the reactants' electronic state with the products', and is equal to one-half the separation of the dotted curves at the intersection of the R and P surfaces in Fig. 1. The quantity (FC) is the 'Franck-Condon' factor: it is a sum of products of overlap integrals of the vibrational and solvational wavefunctions of the reactants with those of the products, suitably weighted by Boltzmann factors. Purely vibrational Franck-Condon factors are available in various

reviews and articles, e.g., Refs. 21, 45 and 57.

Typically, quantum-mechanical theories [45–56] for the nuclear motion have either:

solvent reorganization in Ref. 1.

- (i) treated all coordinates (solvent alone or inner-shell plus solvent) at fixed r as a collection of independent oscillators, harmonic anharmonic ('Morse') §, or
- (ii) treated quantum-mechanically the innershell coordinates as harmonic (or Morse) oscillators, but treated the motion (orientational, libra-

tional) of the solvent outside the inner-coordina-

tion shell as classical, using the free energy of

With this classification in mind we consider next the relation between Eqn. 12 and its classical counterpart for treatment of the motion of the nuclei. When the solvent molecules are treated as a collection of oscillators, namely when assumption (i) is made, Eqn. 12 may be rewritten as

$$k = \frac{2\pi}{\hbar} H_{AB}^2 \sum_{v_r, v_p} (FC)_v p(v_r)$$
 (13)

where $(FC)_v$ is the Franck-Condon (vibrational overlap) factor for any given set of vibrational quantum numbers v_r of the reactants' and v_p of the products' system (including the solvent oscillators), $p(v_r)$ is the equilibrium (Boltzmann) probability of finding the system in the vibrational state v_r , and the sum is over all v_r and v_p . When all of the vibrational frequencies v_i in (FC) are relatively small, e.g., $h\nu_i/2kT \le 1$, and when they are unchanged by reaction, Eqn. 13 reduces to

entire system has been given [58].

[¶] Recently, another treatment of the solvent for organic systems has been assumed in which the solvent near a singly charged organic ion is assigned a harmonic oscillator force constant (for the ion-solvent interaction) which is considerably changed when the ion loses its charge [43]. This interpretation is discussed in Appendix ID.

[§] With the aid of this assumption, the quantity (FC) in Eqn. 12 can be evaluated. However, when the solvent is treated more realistically, namely, by permitting the solvent molecules to perform their actual complicated coupled anharmonic motions, no practical evaluation of (FC) has yet been made. Only a formal expression involving overlap integrals for the

$$k = \frac{2\pi}{\hbar} H_{AB}^2 \frac{1}{(4\pi\lambda RT)^{1/2}} e^{-(\Delta E^{0'} + \lambda)^2/4\lambda RT}$$
 (14)

where $\Delta E^{0'}$ is the energy (enthalpy) of reaction at the separation distance r in the prevailing medium and is negative for an exothermic reaction. On the other hand, when only some of the vibration frequencies are treated as relatively small and unchanged by reaction, with a contribution to λ denoted by λ_o , one obtains [47,49,50] from Eqn. 13:

$$k = \frac{2\pi}{\hbar} H_{AB}^2 \sum_{v_r, v_p} \frac{1}{(4\pi\lambda_o RT)^{1/2}}$$

$$\times e^{-(\Delta E^0 + \Delta \epsilon_v + \lambda_o)^2/4\lambda_o RT} (FC)_v p(v_r)$$
(15)

In applications of Eqn. 15, λ_o is typically intended to represent the orientational contribution from the solvent molecules outside the inner-coordination shells of the reactants and is usually assumed to be given by Eqn. 6. The quantity $\Delta \epsilon_v$ is the vibrational energy of the products minus that of the reactants for the set of quantum numbers (v_r, v_p) .

When assumption (ii) is made instead, Eqn. 13 can be rewritten [52,55,56] as Eqn. 16, which now contains $\Delta G^{0'}$ instead of the $\Delta E^{0'}$ in the exponent in Eqn. 15:

$$k = \frac{2\pi}{\hbar} H_{AB}^2 \sum_{v_r, v_p} \frac{1}{(4\pi\lambda_o RT)^{1/2}}$$

$$\times e^{-(\Delta G^{0'} + \Delta \epsilon_v + \lambda_o)^2/4\lambda_o RT} (FC)_v p(v_r)$$
(16)

The factor in the sum in Eqn. 16 multiplying $(FC)_v p(v_r)$ is equivalent to a Boltzmann-weighted sum of solvational Franck-Condon factors. Eqn. 16 has been recently derived [58] using semiclassical theory (e.g., Ref. 59) to treat the reorientational motion of the solvent. In that way, the reorientational contribution of the solvent to (FC) in Eqn. 12 was converted to a classical expression without introducing any harmonic oscillator or near-harmonic oscillator approximation for the molecular motion of the solvent. In the high-tem-

perature limit, Eqn. 16 reduces to

$$k = \frac{2\pi}{\hbar} H_{AB}^2 \frac{1}{(4\pi\lambda RT)^{1/2}} e^{-(\Delta G^{0'} + \lambda)^2/4\lambda RT}$$
 (17)

where λ is $\lambda_o + \lambda_i$. Eqn. 17 has the same form as Eqns. 1 and 2 ($w^p = w^r = 0$ in a fixed site intramolecular reaction), and provides by comparison a value when $\kappa \ll 1$ for the $\kappa \nu$ appearing later in Eqn. 31 for this fixed-site system.

Unlike Eqn. 16, Eqn. 15 has a major defect when there is a significant entropy of reaction ΔS^0 . In the presence of a polar environment there can be large entropic effects, not reproduced by Eqns. 13-15. These entropic effects result mainly from changes in freedom of the rotations and librations of the solvent molecules. An example of a reaction with a large ΔS^0 is $Fe(H_2O)_6^{2+}$ + Ru(bpy)₃³⁺ \rightarrow Fe(H₂O)₆³⁺ + Ru(bpy)₃²⁺: for this reaction ΔS^0 is $-180 \text{ J} \cdot \text{mol}^{-1} \cdot \text{deg}^{-1}$ (e.g., cited in Ref. 59). Under such conditions assumption (i) becomes grossly inadequate. If, however, there were no solvation outside the inner coordination shell, or if that solvent behaved as a collection of harmonic or anharmonic oscillators of fixed frequencies, one would expect $\Delta S^0 \approx 0$, and ΔE^0 $(\approx \Delta G^0 + T\Delta S^0) \approx \Delta G^0$, and then Eqn. 15 would be satisfactory. Because of this inadequacy of assumption (i) and hence of Eqn. 15, Eqn. 16 has frequently been used in articles to estimate quantum effects, e.g., Refs. 48, 52, 53, 56 and 61. In summary, when a large standard entropy change ΔS^0 accompanies a reaction, it influences the reaction rate and Eqns. 14 and 15 do not allow for this possibility, while Eqns. 16 and 17 do. Indeed, when there is a nonvanishing ΔS^0 , Eqns. 14 and 15 and their counterparts for the reverse reaction rate constant k_{rev} do not obey the correct equilibrium constant relation, $k/k_{rev} =$ $\exp(-\Delta G^{0'}/RT)$, but Eqns. 16 and 17 do.

Eqns 12-17 were obtained for reactions with $\kappa \ll 1$, as already noted. However, the main quantum correction in Eqn. 16 to the classical equation, Eqn. 17, proves to be one for nuclear tunneling, and the latter can occur regardless of whether κ is small or close to unity. Thus, one quantum correction to Eqn. 1, which has been used for $\kappa \approx 1$, is that given by [53,55,61]

tunneling is the WKB method referred to later. The ratio $k_{\rm quantum}/k_{\rm classical}$ has been calculated for a number of reactions using Eqn. 18: when λ_i is relatively small, as in the Ru(NH₃)₆^{3+/2+} self-exchange reaction, the calculated ratio of rate constants is close to unity at room temperature (it equals 1.2) [53]. At relatively large values of λ_i and appreciable vibration frequencies, for example, for the Fe(H₂O)₆^{3+/2+} self-exchange reaction, (the metal-ligand symmetric stretching frequency for Fe(H₂O)₆³⁺ and for Fe(H₂O)₆²⁺ is 490 and 389 cm⁻¹, respectively), the ratio is about 4 [53], and

for the $Co(NH_3)_6^{3+/2+}$ self-exchange it is about 6

[53,62]. The ratio of quantum-mechanical rate constants, $k_{12}/(k_{11}k_{22}K_{12}f_{12})^{1/2}$, for the reaction $Fe(H_2O)_6^{2+} + Ru(bpy)_3^{3+} \rightarrow Fe(H_2O)_6^{3+} +$

 $Ru(bpy)_3^{2+}$ was calculated to differ by only 6% from its classical value of unity [53]. I.e., these

quantum effects have relatively little effect on the

where $(FC)_{q}/(FC)_{c}$ is the ratio of the right-hand

sides of Eqns. 16 and 17, where the vibrational

motion is treated in a quantum manner but the

solvent motion is treated classically. Another way

of making the quantum correction for nuclear

 $k_{\text{quantum}} = k_{\text{classical}} (FC)_{\text{q}} / (FC)_{\text{c}}$

cross-relation in such cases. Where quantum effects on the nuclear motion can be quite large is in the inverted region or at low temperature: Refs. 48-52 and 61 present calculations for a number of examples.

In the case of a highly exothermic reaction and when a single vibration frequency ν is principally involved in the readjustment of nuclear coordinates associated with the electron transfer, the double sum in Eqn. 16 can be approximated by [27]:

$$k = \frac{2\pi}{\hbar} H_{AB}^2 \frac{e^{-S} S^v e^{-(\Delta G^{0'} + vh\nu + \lambda_o)^2/4\lambda_o RT}}{h\nu \Gamma(v+1)}$$
 (19)
where Γ is the Gamma function, S is $\lambda_i/h\nu$, $\Delta \epsilon_v$

where Γ is the Gamma function, S is $\lambda_i/h\nu$, $\Delta\epsilon_v$ in Eqn. 15 has become $vh\nu$, and v is approximately the solution of

the quantum result where tested ($h\nu = 1350 \text{ cm}^{-1}$, and λ_o not too small, varying from 500 to 2500 cm⁻¹) [27]. However, it is reasonably straightforward instead to evaluate the sum in Eqn. 16 numerically or by a 'saddle-point' approximation (e.g., Ref. 53), particularly for a single frequency

is given and used in Appendix II.

Equations 19 and 20 were in good agreement with

model. Another approximate quantum expression

An alternative treatment has been given for vibrational quantum effects, also known sometimes as the 'semiclassical' treatment [63,64]. The expression for the rate constant for electron transfer between fixed sites is of a form rather similar to Eqn. 14, namely: $k = \frac{2\pi}{\hbar} H_{AB}^2 \frac{1}{(4\pi\lambda RT\chi)^{1/2}} e^{-(\Delta E^0 + \lambda)^2/4\lambda RT\chi} \quad (21)$

where
$$\chi = \frac{h\nu}{2RT} \coth \frac{h\nu}{2RT}$$

(18)

The factor χ approaches unity when $h\nu/2RT$ becomes small. Eqn. 21 is readily extended to the

case of many vibrational coordinates and has the advantage of simplicity over Eqn. 12. Its disadvantage is one of accuracy [51,65]. For example for the Fe(H₂O)₆^{3+/2+} system, Eqn. 21 has an error of a factor of 15–20 at room temperature [65]. The source of its difficulty has been identified [65], namely in its treatment of nuclear tunneling through the potential-energy barrier in Fig. 1. Only when the products' surface P is extremely steep at the intersection does this particular error become small [65].

Another treatment, quite different from the above one but also labelled semiclassical, involves

use of a semiclassical ('WKB') approximation for the nuclear tunneling probability [66-68]. This treatment has been reviewed recently [21] and examples of its application are given in Ref. [69], both for the cases of small and large κ ($\kappa \approx 1$) in Eqn. 1. It has been developed mainly for singlefrequency systems and, more approximately, for

other systems. A more general version (many-dimensional) of semiclassical (WKB) theory was used, as already noted, in deriving Eqn. 16 [58]. The main quantum correction for the nuclear motion contained in the $(FC)_v$ factor in Eqn. 13, but absent in Eqn. 14 is, as previously noted, one for nuclear tunneling. The semiclassical (WKB) treatment provides one way of estimating this correction.

IIC. Electronic and steric aspects

The rate of electron transfer depends on the nature of the intervening medium and transfers through a series of conjugated bonds (π bonds) and through σ bonds have been discussed [28,70,71]. In extended Hückel calculations the dependence of κ on r was calculated [28,70] to be larger for σ - than for π -bonded systems. In the present article (Section IV) we shall use some averaged distance dependence for the protein medium; this dependence is suggested by other sources of experimental data (subsection IVC).

Recently, orientation effects on electron-transfer rates have been treated using a simplified cavity model for the the reactants (Ref. 72; Cave, R.J., Klippenstein, S.J., Marcus, R.A., unpublished data, and Cave, R.J., Siders, P. and Marcus, R.A., unpublished data). Orientation effects for π -electron systems are expected both because of the nonspherical nature of the electronic orbitals and because of the nonspherical shape of the molecules themselves. The relative contributions of the two effects have been investigated by making a calculation (Ref. 72 and Cave, R.J., Klippenstein, S.J. and Marcus, R.A., unpublished data) for spheroids, which includes both effects, and for spheres, which includes only the orbital shape effect. Applications are also being made to π -electron systems of biochemical interest (Cave, R.J., Siders, P. and Marcus, R.A., unpublished data). One can expect molecules such as the heme part of cytochrome c or the porphyrin part of bacteriochlorophyll to exhibit significant orientation contributions to H_{AB}^2 .

Steric effects in reactions can also occur, as is evident in the difference between the measured self-exchange rates of optically active d-helicene with its d-helicene anion, as compared with that for the reaction with the l-helicene anion [75]. Steric effects have been calculated [76] for the $Fe(H_2O)_6^{3+/2+}$ exchange: in this system the steric

effect originated from the (calculated) interpenetration of the H_2O ligands, to produce a greater H_{AB} at small r (approx. 0.525 nm) [77].

In this section we distinguish and discuss fur-

IID. Nuclear and electron tunneling

ther nuclear and electron tunneling. Nuclear tunneling was depicted in Figs. 3 and 5 ¶. The nuclear tunneling tends to occur at low temperatures, because then the probability is small that the reacting system has enough energy to surmount the barrier in Figs. 3 and 5. At sufficiently low temperatures, in fact, all of the reaction occurs by nuclear tunneling. The rate is then independent of temperature, since reaction is now occurring largely at an energy equal to that of the zero-point vibrational energy of the pair of reactants and surrounding system (provided the energy corresponding to the lowest point of the R surface equals or exceeds that for the P surface).

Electron tunneling occurs when the potential energy acting on the electron in the space between the reacting pair is higher than the energy of the electron in its initial localized site, as in Fig. 2. The nuclear rearrangement permits, as in Fig. 2b, the initial and final localized energy levels of the electron to be the same, and the electron can then tunnel through its barrier from one site to the other. This need for equality or near-equality of the electronic-energy levels is a consequence of the Franck-Condon principle and energy conservation: when the electronic energies are the same at the given nuclear configuration (Fig. 2b) then the total energy is conserved, since (using classical terminology and the F-C principle) the momenta of the nuclei and hence their kinetic energies do not change instantaneously during the electron transfer. Although electron tunneling is a convenient term for describing the process of electron transfer in Fig. 2b, it is, at best, only a physically intuitive one-electron description of the process. A more fundamental but more abstract approach

involves the many-electron quantum mechanical

Recalling that these figures are profiles of many-dimensional surfaces, one sees that this tunneling is also a many-dimensional one.

calculation of the H_{AB} in Eqns. 12–17.

Temperature affects the nuclear motion and thereby the Boltzmann probability of attaining any range of nuclear configurations such as near point a in Fig. 3 or near the intersection region in the same figure. Thus it affects the relative contribution of nuclear tunneling to the reaction rate. Temperature does not affect directly the value of H_{AB} , although H_{AB} will vary somewhat with nuclear configuration (a point neglected in Eqn. 12, where only the dependence of H_{AB} on the separation distance r is considered; the dependence of H_{AB} on nuclear configuration is discussed

in Appendix IC). Thus, temperature does not af-

fect the electron tunneling in Fig. 2 per se: the

existence of a temperature-independent rate constant at low temperatures and a temperature-de-

pendent one at higher temperatures [18] is, instead,

a manifestation of *nuclear* tunneling. Terms frequently used in the context of electron-transfer reactions are 'adiabatic' and 'non-adiabatic'. In an adiabatic electron transfer there is substantial electronic coupling of the reactant pair: there is enough splitting $2H_{\rm AB}$ of the dotted curves in Fig. 1 that this pair remains throughout largely or completely on the lowest potential-energy surface [25]. (In Fig. 1 this lowest surface is the one labelled R in the initial stages of reaction, then is the lower of the two dotted surfaces in the intersection region, and finally becomes the surface labelled P after the passage through that intersection that intersection is the lower of the passage through that intersection the intersection that intersection is the lower of the passage through that intersection the intersection that intersection the intersection that intersection the intersection that intersection the intersection that intersection the passage through that intersection the intersection that it is the inters

tion region.) For an adiabatic reaction the κ in

Eqn. 1 is unity, or approximately so.

In a nonadiabatic reaction the electronic coupling of the reacting pair is so weak, i.e., the H_{AB} in Eqn. 12 is so small, that the probability κ of proceeding from the R to the P surface when the system is in the vicinity of the intersection region in Fig. 1 is very small. Most systems reaching that intersection region now remain on the R surface, instead of going to the P surface. Eqns. 12-17 and 21 assume the reaction to be nonadiabatic. For reactions intermediate between adiabatic and nonadiabatic, and hence with intermediate k, expressions of the Landau-Zener type are available for calculating κ , e.g., cited in Refs. 25 and 67, but the latter is mainly a refinement only infrequently used. In the 'inverted region', namely in the case of Fig. 4, where the solid curves again denote the adiabatic surfaces, a reaction in which the system goes directly from the R surface to the P surface is necessarily nonadiabatic and there is no 'adiabatic' path [34]: the system must 'jump' from one solid curve to the other, unlike Fig. 1, in order to form directly the ground state products.

IIE. Relaxation-limited rates

Throughout this section and the previous one the rate expressions have been those which are appropriate to 'activation control', in which the probability of forming reactive configurations can be calculated from an equilibrium-type distribution function. That is, it is assumed that the probability of finding the system in the vicinity of the intersection region of Fig. 1 is a near-equilibrium one. However, under certain conditions the motion leading to the reactive configurations may partly involve very slow diffusive steps, either in the relative translational motion of the two reactants or in fluctuations of the environment itself, e.g., a slow diffusive-type dielectric relaxation. Then the overall reaction rate may be smaller than the above value. Indeed, there are a number of cases in the literature where the intrinsic rate of a bimolecular reaction is so high that the mutual diffusion together of the two reactants itself becomes the slow step. In this case, the preceding equations can be used in conjunction with an expression for a diffusion-controlled rate constant to obtain an expression for the overall reaction rate constant (Appendix IA). An example has also appeared (cited in Appendix ID) in which the slow step is a dielectric relaxation step, the 'activation controlled' step being very fast in that example. Several papers discussing such a possibility are

III. Comparisons of predictions with experimental measurements on nonbiological systems

IIIA. Cross-reactions

cited in Appendix ID.

The most frequently used form of the theoretical equations has been what is now known in the literature as the 'cross-relation', Eqn. 10. If the work terms in Eqns. 2 and 3 are not neglected, the cross-relation, Eqn. 10, becomes Eqn. 22. The latter

reduces to Eqn. 10, of course, when the work terms can be neglected:

$$k_{12} = (k_{11}k_{22}K_{12}f_{12})^{1/2}W_{12},$$
 (22)

where now

$$\ln f_{12} = \frac{1}{4} \frac{\left(\ln K_{12} + \frac{w_{12} - w_{21}}{RT}\right)^2}{\ln\left(\frac{k_{11}k_{22}}{Z^2}\right) + \frac{w_{11} + w_{22}}{RT}}$$
(23)

$$W_{12} = \exp\left[-\left(w_{12} + w_{21} - w_{11} - w_{22}\right)/2RT\right] \quad (24)$$

and Z was defined earlier (= $A\sigma_{11}\sigma_{22}$); w_{12} is the work term for reaction 7, w_{21} is that for the reverse reaction and w_{11} and w_{22} are that for reactions 8 and 9, respectively. As before, K_{12} is the equilibrium constant expressed in terms of concentrations in the prevailing medium; it varies with salt concentration.

The work terms w_{ij} can be of several types, not just electrostatic. One rather approximate (Debye-Hückel) version for the electrostatic contribution

to the work term is given by

$$w_{ij} = \frac{\frac{z_i z_j e^2}{2 D_s r} \left(\frac{\exp(B\sigma_i \sqrt{\mu})}{1 + B\sigma_i \sqrt{\mu}} + \frac{\exp(B\sigma_j \sqrt{\mu})}{1 + B\sigma_j \sqrt{\mu}} \right)}{\exp(Br \sqrt{\mu})}$$
(25)

where $B = (8\pi Ne^2/1000 D_s kT)^{1/2}$, $\sigma_i = a_i + a'_i$, $\sigma_j = a_j + a'_j$, where a' is the radius of the dominant ion of opposite charge in the ionic atmosphere, and z_i and z_j are the charges of the reactants. For water at 25°C B equals 3.29 nm⁻¹. Even if Eqn. 25 provides a first approximation to w_{ij} , a free-energy term, the corresponding entropy expression is probably quite poor, and should not be used to calculate the effect of ionic strength on the entropy of activation [78].

There have been numerous experimental tests of the cross-relation Eqn. 10, e.g., Refs. 79–83. Some representative examples are shown in Table I. The calculated rate constants were obtained neglecting the work terms. The entries illustrate the fact that the agreement between theory and experiment can be very good: the observed and

TABLE I
COMPARISON OF OBSERVED AND CALCULATED RATE CONSTANTS FOR OUTER-SPHERE ELECTRON-TRANSFER
REACTIONS

The values of $k_{12,\text{calcd}}$ were calculated from Eqn. 10 neglecting the work terms; the calculated values are from the references cited and from Refs. 79-83. The references in the last column should be consulted for the experimental data.

Reaction	$\log K_{12}$	$\begin{array}{c} k_{12,\text{obsd}} \\ (M^{-1} \cdot s^{-1}) \end{array}$	$\begin{array}{c} k_{12,\text{calcd}} \\ (\mathbf{M}^{-1} \cdot \mathbf{s}^{-1}) \end{array}$	Ref.
$Ru(NH_3)_6^{2+} + Ru(NH_3)_5 py^{3+}$	4.40	1.4·10 ⁶	4·10 ⁶	79
$Ru(NH_3)_5 py^{2+} + Ru(NH_3)_4 (bpy)^{3+}$	3.39	$1.1 \cdot 10^{8}$	$4 \cdot 10^{7}$	80
$Ru(NH_3)_6^{2+} + Co(phen)_3^{3+}$	5.42	$1.5 \cdot 10^4$	$1 \cdot 10^{5}$	81
$R_{11}(NH_{-}) \cdot nv^{2+} + Co(nhen)^{3+}$	1.01	$2.0 \cdot 10^{3}$	$1 \cdot 10^{4}$	81
$V_{cc}^{2+} + \text{Co(en)}_{3}^{3+}$	0.25	$5.8 \cdot 10^{-4}$	$7 \cdot 10^{-4}$	81
$V_{2}^{2+} + Ru(NH_3)_6^{3+}$	5.19	$1.3 \cdot 10^3$	$1 \cdot 10^{3}$	81
$V_{2}^{2+} + Fe_{2}^{3+}$	16.90	$1.8 \cdot 10^4$	$2 \cdot 10^{6}$	81
$V_{aq}^{(1)} + Co(en)_{3}^{3+}$ $V_{aq}^{2+} + Ru(NH_{3})_{6}^{3+}$ $V_{aq}^{2+} + Fe_{aq}^{3+}$ $Fe_{aq}^{2+} + Os(bpy)_{3}^{3+}$ $Fe_{aq}^{2+} + Fe(bpy)_{3}^{3+}$ $Ru(NH_{3})_{6}^{2+} + Fe_{aq}^{3+}$	1.53	$1.4 \cdot 10^{3}$	5·10 ⁵	81
$Fe_{2}^{2+} + Fe(bpy)_{3}^{3+}$	3.90	2.7 · 10 4	$6 \cdot 10^{6}$	81
$Ru(NH_3)_{4}^{2+} + Fe_{aa}^{3+}$	11.23	$3.4 \cdot 10^{5}$	2·10 ⁶	84
$Ru(en)_{3}^{2+} + Fe_{aq}^{3+}$	9.40	$8.4 \cdot 10^4$	$4 \cdot 10^{5}$	84
$Mo(CN)_8^{4-} + IrCl_6^{2-}$	2.18	$1.9 \cdot 10^6$	$8 \cdot 10^{5}$	85
$Mo(CN)_{8}^{4-} + MnO_{4}^{-}$	-4.07	$2.7 \cdot 10^{2}$	$6 \cdot 10^{1}$	86
$Mo(CN)_8^4 - + HMnO_4$	8.48	$1.9 \cdot 10^{7}$	$2 \cdot 10^{7}$	86
$Fe(CN)_{6}^{4-} + IrCl_{6}^{2-}$	4.08	$3.8 \cdot 10^{5}$	$1 \cdot 10^{6}$	85
$Fe(CN)_{6}^{4-} + Mo(CN)_{8}^{3-}$	2.00	3.0·10 ⁴	4·10 ⁴	85
$Fe(CN)_{6}^{4-} + MnO_{4}^{-}$	3.40	$1.7 \cdot 10^{5}$	6·10 ⁴	87

calculated rate constants generally agree within an order of magnitude. There are some relatively minor trends: the data in the table are typical in

that, with one exception, $k_{12,\text{obsd}} < k_{12,\text{calcd}}$. However, in almost all of these the $\text{Fe}(\text{H}_2\text{O})_6^{3+/2+}$

couple is one of the reactants (suggesting, as one possibility, that its mechanism for self-exchange may not be the same as that for the cross-reactions

[83]). Good agreement is generally found when the

couples involved are substitution inert. Unless corrected for work terms, deviations become larger

when the reactants are oppositely charged. They can also be large when the driving force is large [88]. In other cases nonelectrostatic contributions to the work terms may be important. This appears to be the case, for example, in the cross-reactions of complexes with aromatic ligands (bipyridine or phenanthroline) with aquo ions: in these systems

the calculated rates are 2-3 orders of magnitude faster than the observed rates. Noncancellation of the hydrophobic/hydrophilic interactions in the cross-relation (as well as departures from adiabaticity) have been considered as possible reasons for the breakdown in these unsymmetrical systems [31,79,85,89]. As mentioned above, in certain cases a change in mechanism (for example, from outer-

sphere to inner-sphere) may be involved [83,90].

IIIB. Self-exchange reactions

In the present section we apply the formalism to electron exchange rates of metal complexes in solution. Because the reactants are chemically sim-

ilar to the products (merely an interchange of an electron) there are no net thermodynamic changes associated with the electron transfer; conse-

quently, such reactions have an extra simplicity to them - no effect of ΔG^0 need be included. In addition, in exchange reactions the molecular properties of only one redox couple rather than of

two different redox couples need to be known. The rates of a variety of electron exchange reactions are summarized in Table II. These reactions have been selected because they illustrate the range of application of the theory and because, as will be seen later, several of the couples are used as probes of metalloprotein reactivity. As expected, the electron exchange rates depend upon the nature of the metal center as well as upon the nature of the surrounding ligands. For a given metal center the rates tend to increase with the size of

the ligands, reflecting the decrease in the magni-

tude of the solvent reorganization barrier with

Refs.

increasing size of the reactants, while, for a given set of ligands, the rate constants decrease with COMPARISON OF OBSERVED ELECTRON-EXCHANGE RATE CONSTANTS WITH THE VALUES CALCULATED

TABLE II

FROM THE OUTER-SPHERE, ADIABATIC MODEL The value of the nuclear tunneling factor Γ was calculated from

$$\Gamma = \exp\left\{-\frac{\lambda_{i}}{h\nu_{i}}\left(\tanh\frac{h\nu_{i}}{4kT} - \frac{h\nu_{i}}{4kT}\right)\right\},\,$$

 $Co(bpy)_3^{3+/2+}$ couple.

where v_i is the symmetric breathing frequency. See Ref. 21 for details of the calculations of k_{calcd} ; the values of $\lambda_i/4$ and $\lambda_o/4$ for the six reactions in Table II are 35.2, 28.9, 15.9, 28.9; 3.3, 28.0; 0, 13.8; 73.7, 28.5 and 57.3, 13.8 kJ mol⁻¹, respectively. The

references in the last column should be consulted for the experimental data. Couple $k_{\text{obsd}} (M^{-1} \cdot s^{-1})$ Δd_0 μ (M) (nm)

 $k_{\text{calcd}} (\mathbf{M}^{-1} \cdot \mathbf{s}^{-1})$ (nm) $Fe(H_2O)_6^{2+} + Fe(H_2O)_6^{3+}$ $Ru(H_2O)_6^{2+} + Ru(H_2O)_6^{3+}$ 0.55 0.014 0.65 3.2 78,91 (10^2) $4 \cdot 10^{3}$ 0.009 0.65 88.92 1.0 1.5 $Ru(NH_3)_6^{2+} + Ru(NH_3)_6^{2}$ $3 \cdot 10^{3}$ 0.004 $1 \cdot 10^{5}$ 78,84 0.10 0.67 1.0 $Ru(bpy)_3^{2+} + Ru(bpy)_3^{3+}$ $1 \cdot 10^{9}$ $4 \cdot 10^{8}$ 78,93 0.10 0.000 1.36 1.0

 $Co(NH_3)_6^{2+} + Co(NH_3)_6^{3+}$ 1.0 $\geq 10^{-7}$ 0.022 0.66 7.0 4.10^{-6} 78,94 $Co(bpy)_3^{2+} + Co(bpy)_3^{3+}$ 78,95 0.1 0.019 1.36 5.0 20 ^a The reduced force constant for the $Co(NH_3)_6^{3+/2+}$ couple was used in calculating the inner-sphere reorganization energy for the increasing difference in the metal-ligand distances in the two oxidation states.

Included in Table II are the rate constants calculated from the outer-sphere adiabatic model, but corrected (via Γ) for nuclear tunnelling effects [21]. Here, changes in the bond lengths and angles within the ligands themselves were not considered in calculating λ_i . The λ_0 values for the reactions were calculated assuming that r is equal to the sum of the radii of the two reactants: support for this assumption is provided by the fact that the logarithm of the (precursor corrected) rate constants for the $Ru(NH_3)_6^{3+/2+}$, $Ru(NH_3)_5(py)^{3+/2+}$, $Ru(NH_3)_4(bpy)^{3+/2+}$, $Ru(NH_3)_2(bpy)_2^{3+/2+}$, and Ru(bpy)3^{+/2+} exchange reactions, in which the sizes of the reactants are systematically increased, show a linear dependence upon 1/2a (Fig. 6) [96]. The slope is somewhat larger (30%) than the theoretical value of -7.6 nm (the agreement improves when inner-shell changes are included). The effect of change of metal-ligand bond lengths Δd_0 on ΔG^* is displayed in Fig. 7, where the rate constants for some thirteen self-exchange reactions are corrected via Eqn. 6 for the differences in $\lambda_{\rm o}$ and plotted as $\ln (k_{\rm ex}/K_{\rm A}) + (\lambda_{\rm o}/4RT)$ vs. $(\Delta d_0)^2$ [78], where K_A is an encounter equilibrium con-




Fig. 6. Plot of the logarithm of the precursor-corrected exchange rate constants as a function of 1/2a. (1), $Ru(NH_3)_6^{3+/2+}$; (2), $Ru(NH_3)_5(py)^{3+/2+}$; (3), $Ru(NH_3)_4(bpy)^{3+/2+}$; (4), $Ru(NH_3)_2(bpy)_2^{3+/2+}$; (5) $Ru(bpy)_3^{3+/2+}$. From Ref. 96.

stant (Appendix IA). The correlation is the one expected from Eqn. 5. The comparatively minor deviations are, in part, due to differences in the normal mode force constants and in Γ .

The dependence of the exchange rates on the solvent dielectric properties predicted by the model is found for some, but not all of the systems studied. Thus the predicted dependence on $(1/D_{op} - 1/D_s)$ is seen for the bis(biphenyl)chromium +/0 [97]. Ru(hfac) $_3^{0/-}$, Ru((CH $_3$) $_2$ bpy)(hfac) $_2^{+/-}$ (hfac = hexafluoroacetylacetonate ion) [98], and aromatic molecule-aromatic anion exchanges [99], but not for the Fe(cp) $_2^{+/0}$ exchange [100] (cp = cyclopentadienyl radical). An example of such a plot is given in Fig. 8. Specific ion-pairing and solvation effects or strong interaction of the reactants could be responsible for the absence of the predicted solvent dependence in the Fe(cp) $_2^{+/0}$ system (where the ligand carries a net negative charge).




Fig. 7. Plot of the sum of $\lambda_o/4RT$ and the logarithm of the precursor-corrected exchange rate constant as a function of $(\Delta d_0)^2$. (1), $\operatorname{Cr}(H_2O)_6^{3+/2+}$; (2), $\operatorname{Fe}(H_2O)_6^{3+/2+}$; (3), $\operatorname{Fe}(\operatorname{phen})_3^{3+/2+}$; (4), $\operatorname{Ru}(H_2O)_6^{3+/2+}$; (5), $\operatorname{Ru}(\operatorname{NH}_3)_6^{3+/2+}$; (6), $\operatorname{Ru}(\operatorname{en})_3^{3+/2+}$; (7), $\operatorname{Ru}(\operatorname{bpy})_3^{3+/2+}$; (8), $\operatorname{Co}(H_2O)_6^{3+/2+}$; (9), $\operatorname{Co}(\operatorname{NH}_3)_6^{3+/2+}$; (10), $\operatorname{Co}(\operatorname{en})_3^{3+/2+}$; (11), $\operatorname{Co}(\operatorname{bpy})_3^{3+/2+}$; (12), $\operatorname{Co}(\operatorname{sep})^{3+/2+}$; (13), $\operatorname{Co}(\operatorname{bpy})_3^{3+/2+}$. From Ref. 78.




Fig. 8. Plot of the logarithm of the rate constant for the bis(biphenyl)chromium $^{+/0}$ exchange reaction as a function of $(1/D_{\rm op}-1/D_{\rm s})$. From Ref. 97.

In view of the generally excellent agreement of

the observed and calculated exchange rate constants it is tempting to conclude that the model is

essentially correct and, in particular, that the assumption that $\kappa \cong 1$ is justified for the systems

considered. This assumption has been explored theoretically for a few systems. For example, in the case of ruthenium systems the spatial extension of the ruthenium 4d orbitals and the electron delocalization on to the ligands through π -backbonding effects suggest that κ for these exchanges is unlikely to be much less than unity [101]. Consistent with this conclusion, a value of $\kappa \approx 0.2$ has been calculated for the $Ru(NH_3)_6^{3+/2+}$ exchange reaction [77]. On the other hand, on the basis of direct 4d-4d overlap of the metal centers, the $Ru(bpv)_{3}^{3+/2+}$ exchange would be highly nonadiabatic at the separation distance defined by first contact of the bipyridine ligands of the two reactants ($\kappa \approx 10^{-4}$ based on a β value in $\exp[-\beta(r-r_0)]$ of 0.12 nm⁻¹). However, a κ value of about unity has been suggested for the Ru(bpy)₃^{3+/2+} exchange by estimating the $\pi^*-\pi^*$ interaction of the two reactants resulting from the

[101].

Possible nonadiabatic effects have also been explored in calculations for some of the other

delocalization of the metal $d\pi$ electron density

onto the π^* orbitals of the bipyridine ligands

systems. Recent calculations of $\kappa(r)$ for the Fe(H₂O)₆^{3+/2+} exchange indicate that the optimum r occurs at 0.525 nm, appreciably smaller than the separation distance 0.65 nm defined by the effective radii of the reactants [76]. The calculations yield $\kappa(r) \approx 0.2$ at r = 0.525 nm compared with $\kappa(r) \approx 10^{-2}$ at r = 0.65 nm. The dependence of $\kappa(r)$ on orientation at small r's (0.525 nm) introduces a steric factor there, as noted earlier.

In contrast to the ruthenium and iron ex-

changes, the cobalt exchanges are accompanied by a spin change and on this basis should proceed very slowly: κ for the cobalt exchanges has been estimated to be $10^{-3\pm1}$ at r = 0.68 nm [21]. Despite this, the good agreement in Table II of the observed rate with the rate calculated assuming $\kappa \approx 1$ leaves very little room for any special nonadiabaticity in the cobalt exchanges. The same conclusion has also been reached in a more extensive comparison of rates, where it was found that the data for a variety of exchanges (including cobalt systems) are consistent with $\kappa \ge 10^{-2}$ [78]. Since many of the cobalt systems studied involve changes in spin quantum number, these results suggest that spin multiplicity restrictions per se are generally not of major importance in determining electron-transfer rates between transition metal complexes. For systems where the spin-orbit coupling is small, for example in electron-transfer

Good agreement between the observed and calculated exchange rates is not always obtained: for example, the $Co(H_2O)_6^{3+/2+}$ exchange reaction proceeds about 10^7 -times more rapidly than predicted [21]. Clearly, there is a more favorable pathway for this reaction than the simple outersphere exchange considered. Various possibilities have been proposed including a pre-equilibrium spin change [101] and an inner-sphere water-

involving organic systems, the spin restrictions can be more important and would reduce the value of

IIIC. Relation between rates and equilibrium constants

bridged mechanism [90].

We next consider reactions in which the electron transfer is accompanied by a net chemical change. For such systems the formalism predicts a

linear relationship between log k and log K (denoted earlier by K_{12}) provided that $|\Delta G^{0'}| \ll \lambda$. More generally, plots of log k vs. log K are predicted from Eqns. 1-3 to be curved with a local slope $\bar{\alpha}$ (at any particular value of $\Delta G^{0'}$) given by

$$\alpha = \frac{1}{2} \left[1 + \Delta G^{0} / \lambda \right] \tag{26}$$

In a number of systems a linear relationship between log k and log K with a slope of 0.4-0.6 is indeed observed (Table III). In systems encompassing large driving forces plots of $\log k$ vs. \log K are found to be curved as expected from Eqn. 26. Often, however, they are essentially linear over the restricted driving force ranges usually used. An example is the family of RuL₃⁺-Eu_{aq}³⁺ reactions (different polypyridine ligands L) for which $-\Delta G^0$ ranges from 1.2 to 1.5 eV, and which is predicted to have a slope of 0.28; the value observed (0.12) is, however, even smaller. The decreased slope has been ascribed to effects of nonadiabaticity [21,101]. Good correlation between $\log k$ and $\log K$ has been found for a broad variety of metalloprotein electron-transfer reactions [111].

TABLE III SLOPE OF PLOT OF LOG k VS. LOG K FOR A SERIES OF RELATED ELECTRON-TRANSFER REACTIONS

Data for electrochemical systems are presented in Appendix ID. The asterisks in the first column indicate electronically excited states of the complexes, and HAs⁻ denotes the ascorbate anion, PTZ N-alkylphenothiazine derivatives, SnR₄ tetraalkyltin derivatives, P²⁺ bipyridinium derivatives, and ArNR₂ aromatic amines.

Reaction	Slope	Refs.
$FeL_3^{3+} + Fe_{aq}^{2+}$	0.51	102
$FeL_3^{2+} + Ce(IV)$	0.44	103
$RuL_3^{3+} + Fe_{aq}^{2+}$	0.51	89
$OsL_3^{3+} + Ti_{aq}^{3+}$	0.37	104
$*RuL_3^{2+} + Eu_{aq}^{3+}$	0.49	89
$*CrL_3^{3+} + Fe_{aq}^{2+}$	0.43	105
$FeL_3^{3+} + HAs^{-}$	0.50	106,107
$Fe_{aq}^{3+} + PTZ$	0.50	108
$Fe(phen)_3^{3+} + SnR_4$	0.50	109
$P^{2+} + *Ru(bpy)_3^{2+}$	0.50 a	110
*Ru(bpy) $_{3}^{2+}$ + ArNR ₂	0.50 a	110

a In acetonitrile.

IIID. Mixed-valence systems

Mixed-valence complexes in which two metal centers are connected by a bridging group provide systems that are very useful for the comparison of theoretical and experimental parameters. Not only are the theoretical problems associated with the formation of the precursor complex eliminated, but the distance separating the metal centers is also known. In addition, many mixed-valence systems feature an intervalence absorption band in the near-infrared region of the spectrum and $H_{\rm AB}$, $\lambda_{\rm i}$ and $\lambda_{\rm o}$ can be obtained from the intensity, energy and solvent dependence of this band [112,113].

The energy of the intervalence transition $E_{\rm op}$ is equal to $\lambda + \Delta G^{0}$, [114] ¶ , whence the ΔG^{*} in Eqn.

[¶] In Refs. 112 and 113 $E_{\rm op}$ is assumed to be $\lambda + \Delta E^0$, where ΔE^0 is the internal energy change. However, in Ref. 114 the transition energy is shown to be a free-energy change \(\lambda\) + ΔG^{0} . The equality of $E_{\rm op}$ and $\lambda + G^{0}$ is evident from Eqn. 14 of Ref. 114, where $F_{\rm e^-g}^{\rm op} - F_{\rm e^-g}$ is the polar contribution to λ and $F_{\rm e} - F_{\rm g}$ is the polar contribution to ΔF^0 (ΔG^{0}) in the present notation). In more physical terms, the reason why $E_{\rm op}$ equals $\lambda + \Delta G^{0}$ rather than $\lambda + \Delta E^{0}$ can be seen from the following argument. According to the Franck-Condon principle, the momenta and positions of the nuclei do not change during the vertical transition. Consequently, the instantaneous distribution of momenta and coordinates is the same immediately before and just after the optical transition and there is no change in the entropy during the vertical transition (apart from entropy changes associated with changes in electronic multiplicity). Let the vertical transition energy $E_{\rm op}$ be denoted by λ when $\Delta G^{0\prime} = 0$. At $\Delta G^{0\prime} = 0$, we have $\Delta H^{0\prime} = T\Delta S^{0\prime}$, and so $E_{\rm op}$ can be written equivalently as $\lambda + \Delta H^{0}' - T\Delta S^{0}'$. If, now, the products' potential-energy surface is vertically displaced downward, ΔS^{0} is unchanged, but ΔH^{0} is made more negative. The $E_{\rm op}$ is now λ plus this (negative) change in ΔH^{0} . In other words, when $\Delta G^{0} \neq 0$, we can still write E_{op} as $\lambda + \Delta H^{0} - T\Delta S^{0}$, i.e., as $\lambda + \Delta G^{0}$, as in the text. The \(\lambda \) defined above can be related to the free-energy barrier in the thermal electron transfer as follows: according to the key approximation made in deriving Eqns. 1 to 3, the free energy of each of the surfaces is a quadratic function (actually of a quantity m in Refs. 1-3), and it follows from this property that when $\Delta G^{0}' = 0$ the free-energy change for reaching the intersection is one fourth the free energy change for the vertical transition. Since, as we have already noted, there is no change in entropy during the vertical transition, the vertical change in free-energy equals that in energy, namely E_{op} . Thus, the free-energy barrier for the thermal transfer is $E_{op}/4$, i.e., $\lambda/4$.

which is now absent) $\Delta G^* = E_{\rm op}^2 / 4 \left(E_{\rm op} - \Delta G^{0} \right).$

(27a)The ΔH^* and ΔS^* for this first-order reaction are

defined by the standard thermodynamic relations
$$\Delta H^* = \left(\frac{\partial (\Delta G^*/T)}{\partial (1/T)}\right)_n \tag{27b}$$

2 becomes Eqn. 27 (apart from the w^r in Eqn. 2,

(27b) (27c)

$$\Delta S^* = -\left(\frac{\partial \Delta G^*}{\partial T}\right)_p = \frac{\Delta H^* - \Delta G^*}{T}$$
 (27c)
with ΔG^* given by Eqn. 2 or, for the present system, by Eqn. 27a. Eqn. 27a and b yield [115]:

$$\Delta H^* = \frac{E_{\rm op} \left[E_{\rm op} + 2(\Delta H^{0\prime} - \Delta G^{0\prime}) \right]}{4(E_{\rm op} - \Delta G^{0\prime})}, \qquad (27d)$$
and ΔS^* then equals $\frac{1}{2}\Delta S^{0\prime}$, $E_{\rm op}/(E_{\rm op} - \Delta G^{0\prime})$.
$$(\Delta H^* \text{ and } \Delta G^* \text{ have been confused in certain applications involving } E_{\rm op} \text{ in the literature, and } Eqn. 27d is new [115].$$
If the intervalence band has a Gaussian shape and the degree of ground-state delocalization is

and the degree of ground-state delocalization is small, then H_{AB} is related to the properties of the band by [112,113]:

band by [112,113]:
$$|H_{AB}| = \frac{2.06 \cdot 10^{-3}}{r} \left(\epsilon_{\max} \bar{\nu}_{\max} \Delta \bar{\nu}_{1/2}\right)^{1/2} \text{ cm}^{-1}(28)$$
 where r is in nm and ν in wavenumbers. At room

temperature the energy and half-width of a Gaus-

sian-shaped band (for the case $\Delta G^{0\prime} = 0$) are re-

lated by Refs. 112 and 113 (cf.Ref. 114): $\Delta \bar{\nu}_{1/2} = (2.31 \cdot 10^3 \ \bar{\nu}_{\text{max}})^{1/2} \ \text{cm}^{-1}$ (29)

(When $\Delta G^{0'} \neq 0$ the $\bar{\nu}_{\text{max}}$ in Eqn. 29 is replaced by $\bar{\nu}_{\text{max}} - \Delta G^{0'}$, where $\Delta G^{0'}$ is now in units of cm⁻¹.) Consequently, the half-width can be eliminated from Eqn. 28 which then contains only r, ϵ_{max} , and $\bar{\nu}_{max}$ as parameters. Another way of obtaining H_{AB} is from EPR, namely from the singlet-triplet splitting of a radical pair [64,116,117]: such a method yields a value for the electron-transfer matrix element H_{AB} when the splitting is predomitransfer state [116,117]. The latter affects the en-

ergy of the singlet, but not of the triplet, radical pair. The application of Eqns. 27a, 28 and 29 will be illustrated by considering the electron transfer

within mixed-valence diruthenium systems. There is reasonable agreement between $\bar{\nu}_{max}$ (9.7 · 10³ cm⁻¹) for the 4,4'-bipyridine-bridged system and the value calculated $(12.8 \cdot 10^3 \, \text{cm}^{-1})$ using a model in which each half of the binuclear ion is approximated by a sphere of radius 0.37 nm [96,118]. Similarly, the solvent dependence of the intervalence band (slope of $\bar{\nu}_{max}$ vs. $(1/D_{op} - 1/D_{s})$) is only 30% smaller than the value calculated from $E_{\rm op} = \lambda_{\rm i} + \lambda_{\rm o}$

(30)

nantly due to the presence of a singlet charge-

where λ_i and λ_0 are given by Eqns. 5 and 6, with a = 0.37 nm and r = 1.08 nm. The value of H_{AB} calculated from the intensity of the intervalence band [119,120] is 400 cm⁻¹, corresponding to $\kappa \approx 1$. Introducing a -CH₂-group between the two pyridine rings reduces the coupling of the metal centers, so that the electron transfer becomes somewhat nonadiabatic ($\kappa = 0.2$). Nevertheless, the coupling in the latter system is still very large compared with the through-space coupling of two ruthenium centers a distance 1.0-1.2 nm apart for which $\kappa \approx 10^{-4}$ has been estimated as discussed

In the Creutz-Taube ion [121] the two ruthenium

centers are connected by a pyrazine group. This group provides strong electronic coupling of the metal centers instead of weak coupling. The properties of the pyrazine-bridged system are those expected for a delocalized rather than a localized ground state, (and so Eqn. 6 becomes inapplicable). The near-infrared band of the Creutz-Taube ion does not show the solvent dependence expected for weak overlap intervalence systems and,

in contrast to the band of the bipyridine-bridged system, the near-infrared band of the pyrazinebridged complex is much narrower than predicted by Eqn. 29 [113]. This is an interesting system and the proper description of the Creutz-Taube ion is still a subject of debate [122]. A three-site electronic-nuclear model has been discussed recently

for such strongly delocalized bridged systems [123–125].

Although a bridging group is useful, it is not required for the observation of an intervalence absorption band: such bands are also observed in outer-sphere ion pairs and can be analyzed in terms of the above two-site formalism [126].

Experimental [127,128] and theoretical [71] studies of the dependence of absorption intensity of charge-transfer spectra on distance along fixed bridges have been described recently. A dependence of H_{AB} on the coordinates (e.g., in Figs. 1–4) has also been considered [129]. Such a dependence can cause the matrix elements for the thermally and optically induced electron transfer to differ [129]. However, when a realistic model is used instead of a delta function or a square-well potential, the difference between the matrix elements becomes much smaller [71].

A phenomenon related to charge-transfer spectra is the photoionization of ions in solution [130,131]. The photoionization spectrum of each ion is related to the value of λ plus the ΔG^0 for this process [131].

IIIE. Electrode reactions

There has been an extensive comparison of the self-exchange rates with the rates of the corresponding reactions at an electrode, determined at zero activation overpotential in order to test the relation predicted in Eqn. 11 (e.g., Refs. 17, 132,133 contain recent summaries). There is a good correlation at lower rates, but at the highest rates the available data for the electrochemical rates appear to show a levelling-off.

There have been experimental studies, e.g., cited in Appendix ID, of the theoretical analog [3,29-31] of Eqn. 26 for electrode reactions in which the activation overpotential $ne(E - E'_0)$ replaces $\Delta G^{0'}$. (E is the half-cell potential, E'_0 the 'standard' half-cell potential in the prevailing medium, and n is the number of electrons, of charge e, transferred between reactant and electrode in the elementary step.) Recent studies of the curvature of plots of $\ln k$ vs. $E - E'_0$ have also been made [134,135]. They yield approximate agreement of the theoretical and calculated $E - E'_0$ values for onset of significant deviation of α in Eqn. 26 from 0.5.

IV. Extension to biological electron transfers

IVA. Similarities and differences with small molecule electron transfers

We first note that biological electron transfers have certain features in common with the small molecule electron transfers, as well as features which are different. As in the simpler reactions, the reactants in biological systems can be expected to undergo vibrational and solvational (or environmental) changes during the reaction, and so the rates should depend on λ and ΔG^0 . However, substantial differences also exist. (1) Whereas it was possible in the solution case to find self-exchange reactions for predicting the rate of a crossreaction, the former may sometimes not even exist in the biological case. (2) Whereas the structural information (bond length changes) needed for the prediction of absolute rate constants of self-exchange reactions is becoming increasingly available in the case of reactions in solution (particularly with the introduction of EXAFS) as are the relevant vibrational frequencies, such information is at present only fragmentarily available for most biological systems. Indeed, in some cases even the identity of all the coordinated ligands may not be known. (3) The environment (solvent) for the electron-transfer reactions in solution, which affects λ , is at present better known than that, for example, in protein systems, which are structurally less homogeneous. (4) Whereas ΔG^0 is usually quite well-known for reactions in solution, knowledge of it for the case where the pair of reactants are either in, or attached to, a membrane is frequently less certain \P . (5) Quantities such as ΔG^0 now become dependent on any electric field which has developed across the membrane. (6) Protein-conformational changes may precede or follow the electron transfer in biological systems. For example, such conformational changes may modify relative binding free energy of the oxidized and the reduced

Further, in applications of the theoretical expressions to an electron transfer step (both in a biological and a nonbiological reaction) one also needs to be particularly careful that the ΔG^0 used refers only to that step and hence that any experimental value is corrected, when necessary, for any loss or gain of protons.

forms of a substrate to the protein. (7) Whereas most electron-transfer reactions in solution are expected to occur at or near contact between the reactants, with a reaction probability which depends on λ and ΔG^0 , reactants in a biological membrane are frequently fixed in position, and so are prevented from coming in contact. (In this respect the biological reactants are similar to those in long range non-biological intramolecular electron-transfer reactions.) Their rate is expected to vary roughly as $\exp(-\beta r)$, i.e., to decrease exponentially with separation distance r of the reactants and the rate now depends not only on the intrinsic (λ) and thermodynamic (ΔG^0) factors, but also on β and on the mutual orientation of the reactants. It also depends on r through the outer (λ_0) contribution to λ . Thereby, β and r become important parameters for any proper understanding of the rates.

Before proceeding to a discussion of some individual reactions, we first consider one general aspect of biological electron transfers: they often occur via a series of intermediates at fixed sites, for example in the charge transfer across a membrane for the photosynthetic reaction center (subsection VD). There are at least two factors which influence the choice of a mechanism involving several reaction intermediates versus one involving a single giant step. To illustrate these factors we consider a simple example of a system abc with one intervening site b. For transfer to occur in one hop (e.g., for mediation by b to occur via a superexchange mechanism without a well-defined intermediate b being formed), there is first a reorganization of the nuclear configuration of the end sites a and c, to make their energies equal, the system thereby reaching the intersection region in Fig. 1. The vertical energy difference of site b and end sites a and c is denoted by V. H_{abc} is the effective electronic matrix element coupling a and c and in a superexchange mechanism (Appendix IC) equals $H_{ab}H_{bc}/V$, when $V \gg H_{ab}$ and H_{bc} §.

The site b poses little additional barrier only when $H_{\rm bc}/V$ is not too small, and the transfer then occurs in a single hop. A single hop has the advantage for a downhill reaction of utilizing in full the overall $-\Delta G^{0}$, typically then requiring less reorganization free energy in the normal region, in accordance with Eqns. 1-3. On the other hand, when H_{bc}/V is very small, the transfer will occur in two successive hops provided that the formation of an actual intermediate b^- is not too 'uphill'. The matrix element for the first hop is now H_{ab} , and is H_{bc} for the second hop. The $-\Delta G^{0'}$ for at least one of these hops is now less favorable than the overall $-\Delta G^{0'}$, and so may require a somewhat larger free energy of reorganization. Thus, the actual choice of mechanism depends on the relative contributions of the $\Delta G^{0'}$ and $H_{\rm bc}/V$ factors discussed above.

IVB. Reorganization parameters

We next examine how the equations of Section II might be modified so as to treat reactions of metalloproteins, for instance those between cytochrome c and various inorganic redox reagents. In applying Eqns. 1-4 the detailed description of the λ_o and λ_i contributions to the biological system (cytochrome c in the present case) may differ from those for the ions in solution, and the magnitude of κ becomes a particularly important question if there is an appreciable distance between the two electron-transfer sites. There may also be contributions to w or λ_i arising from conformational changes.

The geometry of most metalloproteins, particularly as regards the dielectric situation, differs from that used to derive Eqn. 6 (although one might use Eqn. 6 with some mean D_s for the protein and/or membrane system). A more elaborate model to calculate λ_o [137] can, of course, be introduced. In the present article we explore the possibility of using Eqns. 1–3. This we do in Section V, by first exploring the applicability of the cross-relation, Eqn. 10. Eqn. 2 has been used for cytochrome c recently (apart from w^r) [137]. The authors made an estimate of the reorganization parameter λ (Eqn. 4) using a molecular model for the protein [137]. Electron transfer between most reactants in biological systems oc-

This result, which is used here for illustrative purposes only, applies when there is only one orbital per bridge. In general, there is a sum of these $H_{\rm ab}H_{\rm bc}/V$ terms over the orbitals of the bridge, and $H_{\rm ab}$, $H_{\rm bc}$ and V vary with the bridge orbital ν . See, for example, Eqn. 15 of Ref. 70, where $H_{\rm ab} = \eta_1 c_{1\nu}$, $H_{\rm bc} = \eta_2 c_{n\nu}$ and $V = b_{\nu} - a$, in the notation of Ref. 70.

(34)

(35)

The first-order rate constant k can then be written

curs between spatially fixed and oriented sites. In

this case the reaction is formally a first-order one.

$$k = \kappa(r)\nu \exp(-\Delta G_r^*/RT)$$
 (31)

$$\Delta G_r^* = \frac{\lambda}{4} \left[1 + \frac{\Delta G^{0'}}{\lambda} \right]^2$$
where λ again equals $\lambda_i + \lambda_o$, ΔG_r^* and $\Delta G^{0'}$ are

the free energy barrier and the free energy of reaction, respectively, when the reactants are a distance r apart in the prevailing medium, and ν

has dimensions of a frequency (s^{-1}) . In the adiabatic regime ν is an effective frequency for nuclear motion along the reaction coordinate in Fig. 1

(discussed in Appendix I). In the nonadiabatic regime ($\kappa \ll 1$), $\kappa \nu$ is actually independent of the frequency of nuclear motion: comparison of Eqn. 31 with Eqn. 17 shows that when $\kappa \ll 1$, $\kappa \nu$ is given by:

$$\kappa(r)\nu = 2\pi \frac{H_{AB}^2}{\hbar (4\pi\lambda RT)^{1/2}}$$
 (33a)

As discussed later H_{AB}^2 decreases exponentially with separation distance for many systems and so we shall write $\kappa(r)\nu$ as

$$\kappa(r)\nu = 1 \cdot 10^{13} \exp[-\beta(r - r_0)] \text{ s}^{-1}$$
where r_0 is the value of r at which $\kappa \nu$ equals some

preassigned value, $1 \cdot 10^{13} \text{ s}^{-1}$. Some experimental results in *nonbiological* systems place β at about 12 nm⁻¹ for aromatic molecules, and perhaps less if one of the states is electronically excited (cf.

Section IVC). The values of β are likely to vary from system to system somewhat, depending, for example, on the vertical ionization potential of the redox site and its determination for various model systems, for reactions of ground states, electronically excited states, anions, and cations, and its dependence on the intervening material present a

central experimental problem in this field. If the frequency for motion of the nuclei along the reaction coordinate is $1 \cdot 10^{13}$ s⁻¹, a typical

vibrational frequency, then in using Eqns. 31-33,

one is tacitly assuming that the reaction is 'adiamolecular reactions the separation distance where the reaction is adiabatic is frequently assumed to correspond to close contact of the two reactants.

batic' at the separation distance $r = r_0$, specifically that when $r = r_0$ the system moves from the R to the P surface upon every passage through the nuclear configuration defining the intersection region in Fig. 1 (cf Appendix IA). Strictly speaking, the κ in Eqn. 33 need not immediately fall exponentially with r for $r \ge r_0$: there could be some small interval for r, for which κ is roughly constant and then the $\kappa(r)$ changes smoothly to an approximately exponential decrease with r. In bi-

appears in Eqn. 32 is for the conditions prevailing in the membrane and as such, can vary with the nearby charges present and with the electric fields they produce. If the electric field produces a given potential ψ_A at the charge center A and ψ_B at charge center B, then ΔG^{0} can be written as [24]: $\Delta G^{0'} = \Delta G_0^0 + \Delta e_{\Delta} (\psi_{\rm R} - \psi_{\Delta}),$

where Δe_A is the (positive) charge transferred from site A to site B. This equation presupposes that there is no significant change of the potential ψ

itself when the electron transfer occurs. ΔG_0^0 is the

For membrane-bound systems the $\Delta G^{0'}$ which

value of $\Delta G^{0'}$ in the absence of ψ . Eqns. 31-33 have a number of applications. For example, one may vary $\Delta G^{0'}$ by systematically varying one of the reactants. If λ also varies, then an estimate of its variation is also needed (e.g., from self-exchange data). We discuss examples for

bimolecular reactions in Section VA. A second application is to the activation energy E_a of a first-order reaction. E_a/R is defined as the slope of the ln k vs. 1/T plot, i.e., it equals $-\partial$ ln $k/\partial(1/T)$. From Eqns. 27b, 31 and 32, E_a for a first-order reaction is therefore given approximately by:

$$E_{\rm a} \approx \Delta H_r^* \approx \frac{\lambda}{4} + \frac{\Delta H^{0'}}{2} \left[1 + \frac{\Delta G^{0'}}{\lambda} \right] - \frac{\left(\Delta G^{0'}\right)^2}{4\lambda}$$

when λ and $\kappa \nu$ are assumed independent of temperature. In the normal region the magnitude of $\Delta G^{0'}/\lambda$ in Eqn. 35 is less than unity. We see that when $\Delta H^{0'}$ is very negative, the activation energy can even be negative. We have discussed elsewhere [60] such an application of Eqn. 35.

The first-order rate constant k can be written as

$$k = A' \exp(-E_a/RT), \tag{36a}$$

where E_a is given by Eqn. 35. Using Eqns. 31 and 35 the preexponential factor A' is given by

$$A' = \kappa(r)\nu \exp(\Delta S_r^*/R), \qquad (36b)$$

where ΔS_r^* is obtained from the thermodynamic relation, Eqn. 27c (with r subscripts).

It is sometimes useful in discussions to introduce a term, the 'entropy of activation' ΔS^{\dagger} , which includes both ΔS_{\star}^{*} and an entropic contribution from $\kappa \nu$, namely $R \ln(A'/10^{13})$, where A' is in s⁻¹. Thereby,

$$\Delta S^{\dagger} = \Delta S_{\rm r}^* + \ln(\kappa \nu / 10^{13}), \tag{36c}$$

where $\kappa \nu$ is in s⁻¹. Using Eqns. 27c, 32 and 35, ΔS_r^* can be related to $\Delta S^{0'}$, the standard entropy of reaction when the reactants are a distance r apart in the prevailing medium, namely $(\Delta H^{0'} - \Delta G^{0'})/T$, as follows:

$$\Delta S_r^* = \frac{1}{2} \Delta S^{0'} (1 + \Delta G^{0'} / \lambda)$$
 (36d)

Eqn. 36c and d are applied later.

We shall have occasion to refer to the 'barrier-less regime', namely the region where $\Delta G^{0'}$ is sufficiently negative that $-\Delta G^{0'} \approx \lambda$. In this case, one sees from Eqns. 32, 35 and, with r subscripts, 27c that ΔG_r^* , ΔH_r^* and ΔS_r^* are each approx. zero. In particular, the activation energy E_a is approx. zero. (This point is also discussed in Appendix II.)

In the case of bimolecular reactions it is con-

venient to use a formalism for the rate constant based on the fixed site first-order rate constant k in Eqn. 36a and a pre-equilibrium constant K_A for a precursor complex, described in Appendix I and used in Section V.

IVC. Electron tunneling barriers. Distance dependence of electron-transfer rate in nonbiological systems

We next present a brief description of determinations of β in Eqn. 33 from nonbiological data. Several methods have been used:

(1) The first is based on the study of electron-transfer reactions in frozen media. Initially, there is a random distribution of the reactants. Since the closer pairs react first, the subsequent distribution is no longer random: consistent with Eqn. 33 the pairs at a distance r have a rate constant k(r) given by

$$k(r) = k_0 \exp[-\beta(r - r_0)]$$
(37a)

Using an expression for the fraction of pairs remaining at time t and at each r and then integrating over r, an expression for the fraction of unreacted material at time t is obtained. Fitting the experimental data to the expression yields values of k_0 and of β . In many reactions between solvated electrons and various acceptors β varied between 10 and 20 nm⁻¹ (data not listed in Table IV, but summarized in Ref. 138). In electron-transfer reactions of biphenyl anions with some aromatic molecules [139], the anions being formed by pulse radiolysis, β was calculated [138] from the data, and is given in the Table. Results for other systems are also given in Ref. 138, and most recently in Ref. 37 where a value of $\beta = 12 \text{ nm}^{-1}$ has been obtained (Table IV). Studies of hole transfer in frozen media were also reported [141], namely

ephenylenediamine (cf. Table IV).

(2) Monolayers of fatty acids have been deposited on an aluminum layer, and then another metal layer deposited to form a capacitor, cf. [147] and references cited therein. Conductivities were measured for monolayers of fatty-acid salts containing 14–23 carbon atoms, the second metal

being mercury [143], magnesium [144], gold [144],

from pyrene+ and biphenyl+ to tetramethylen-

[¶] A closely related definition of an 'entropy of activation' is frequently used in the literature. For example, the A' in Eqn. 36a is often equated to $(k_BT/h) \exp(\Delta S^\dagger/R)$, where k_B is Boltzmann's constant. However, since k_BT/h is approximately $1 \cdot 10^{13}$ s⁻¹ at room temperature, this definition is equivalent to that used in the text for reactions studied under typical conditions.

TE VALUES OF A IN THE LITERATURE

TABLE IV

multilayer system.

then.)

System	β (nm ⁻¹)	Comment	References
Biphenyl-aromatics a	12	frozen media	37
Biphenyl naphthalene	12	frozen media	138, 139
Biphenyl phenylethylene	10	frozen media	138,139
Biphenyl -acridine	11	frozen media	Miller, J.R., personal communication
Pyrene ⁺ -TMPD ^b	11.5	frozen media	141
Biphenyl ⁺ -TMPD ^b	11.5	frozen media	141
Biphenyl -cinnamoyl grp.	(<10)	steroid bridged	142
Al/C, /Hg, Al or Au	15	conduction	143,144
Al/C _n /Mg	10	conduction	144
Al/C, /Al	10.5-15 °	conduction, multilayer	145
Al/C _n , dye/Al	3-5	photoconduction, multilayers	146
dye/C _n /e ⁻ acceptor	≈ 3	fluorescence quenching	147
Al/C_n /anthracene	4.5	photoconduction	148
$Al/C_n/chloranil$	≈ 8	photoconduction	149

^a Many aromatic acceptors were used and gave an average value of $\beta = 12 \text{ nm}^{-1}$. ^b TMPD denotes tetramethylphenylenediamine.

studies with multilayers [144] gave a value 10.5–15 (only a few points however), consistent with the monolayer studies if hopping between traps separated by the monolayers was assumed. Photoconduction experiments performed in multilayered structures dyes and fatty acids gave a smaller $\beta \approx 3-5 \text{ nm}^{-1}$ [146]. A similarly small β was obtained in fluorescence quenching experiments in which a layer of dye and a layer of acceptor were

separated by a fatty-acid monolayer [147], and in

photoconduction experiments in which a (semi-

transparent) aluminum layer was separated from

an anthracene crystal [148] or a chloranil crystal

[149] by a fatty acid monolayer. Thus, β appears

or aluminum [145]. The results for β were 15

nm⁻¹, apart from that for magnesium ($\beta = 10$

nm⁻¹) and are included in Table IV. Analogous

^c Our estimate of the uncertainty in the slope of Fig. 2 in Ref. 145. A multilayered hopping formula wasused in Ref. 145 for the

in less than 0.5 ns [142]. Using a value for k_0 of $1 \cdot 10^{13} \text{ s}^{-1}$ and $\Delta G^* = 0$ yields $\beta < 10 \text{ nm}^{-1}$, while $k_0 = 1 \cdot 10^{14} \text{ s}^{-1} \text{ yields } \beta < 12 \text{ nm}^{-1}.$ Summarizing these results we see that the value

of the β in Eqn. 37a for the aromatic systems in Table IV is about 11-12 nm⁻¹, is higher when the transfer is from metals, and is less in both cases when photoexcited systems are involved. Incidentally, a flash photolysis study of electron transfer from an excited zinc porphyrin to a quinone ring bound a distance r of ≈ 1.0 nm above the porphyrin plane yielded $k \approx 1 \cdot 10^9 \text{ s}^{-1}$ [150], while the maximum calculated k assuming $\beta = 12 \text{ nm}^{-1}$

studies, including temperature dependences, are very desirable, and data for metalloprotein, polypeptide and other porphyrin systems are becoming available [38,151-156]. A rough interpretation of β is in terms of

 $r - r_0 = 0.7$ nm (see below) and $k_0 = 1 \cdot 10^{13}$ s⁻¹

(and $\Delta G^* = 0$) is about $1 \cdot 10^9$ s⁻¹. Clearly, further

to be smaller when one of the reactants is electronically excited. (There is a smaller barrier in Fig. 2b (3) Particularly interesting are intramolecular electron-transfer rate measurements in which the separation distance is varied. In an experiment in which aromatic reactants are separated from each other by a steroid molecule, at an edge-to-edge separation distance of about 1.0 nm the electron transfer for the most exothermic reaction occurred

electron tunneling through a square potential barrier, e.g., Ref. 143. (A more satisfactory model is discussed in Appendix IC.) Using a standard quantum-mechanical tunneling formula, β equals $(2/\hbar)(2mV_0)^{1/2}$, where m is the mass of the electron, V_0 the height of the potential barrier (Fig. 2)

and \hbar is $h/2\pi$. Values of β of 11 and 15 nm⁻¹

correspond to V_0 values of 1.1 and 2.1 eV, respectively. The V_0 is, in effect, a vertical ionization energy to the lowest unoccupied molecular orbital in the medium spanning the two redox centers, for the case where the nuclear configuration corresponds to the intersection region in Figs. 1 and 2. (Alternatively, it may be the corresponding quantity for a 'hole' [63].) The value of 14.4 nm⁻¹ proposed for β in Ref. [63] was based on an assumed barrier of 2 eV.

Strictly speaking, the pre-exponential factor k_0 in Eqn. 37a for a site-to-site electron transfer is also expected to have some polynomial dependence on 1/r, because of a geometrical factor appearing in the overlap of the electronic wavefunctions, see, e.g., Ref. 72. For the present purposes we shall regard this additional dependence on r as approximately included in the exponential factor.

In addition to the electronic distance effect βr , there is another distance dependence which arises when the environment is a polar one, namely the increase of λ with r, as in Eqn. 6: in the inverted region this additional effect causes the rate constant to increase initially with increasing r and then, because of the exponential decrease of the electronic factor with βr , to decrease [158]. The dependence of λ on r can cause a wrong value of β to be inferred from data on the distance dependence of intramolecular rate constants, unless allowance is made for it.

We next adapt Eqn. 33a to nonspherical molecules, such as the porphyrins occurring in biological systems. Structural information is normally given in terms of distances between atomic nuclei. However, a distance of particular interest for electron transfer is the distance between the closest atoms (of the two reactants) that are strongly coupled to their respective redox sites. When this distance is small enough $\kappa\nu$ equals some preassigned value, taken earlier as $1\cdot 10^{13}$ s⁻¹. Eqn. 33b can then be rewritten as

$$\kappa \nu = 1 \cdot 10^{13} \exp(-\beta d) \text{ s}^{-1},$$
 (37b)

where d is equal to the actual separation of the centers of the two closest (in the case of aromatics) carbon atoms of the two reactants minus some amount to allow for the extension of the π -elec-

tronic orbitals beyond the carbon nuclei. For concreteness we will use a value of 0.3 nm for the latter amount. (Thereby, if the frequency of motion along the reaction coordinate were about $1 \cdot 10^{13} \text{ s}^{-1}$, the reaction would be adiabatic ($\kappa = 1$) when these aromatic carbons are a distance 0.3 nm or less apart.) The value of $\kappa \nu$ also depends, of course, on the detailed geometry (e.g., Appendix IA), such as mutual orientation of the reactants, but for the present purpose we focus attention on the distance of closest approach.

IVD. Work terms for bimolecular reactions involving metalloproteins

There have been a number of studies of the effect of ionic strength on the rates of electron transfer between metalloproteins and metal complexes [159-167]. The charges on the reactants are varied in these studies by using metal complexes of different charge types and/or by varying the charges on the protein through pH variations or by modifying the protein. The results are frequently interpreted by approximating the metalloprotein by a sphere and calculating its radius either from the partial specific volume of the protein in aqueous solution or from crystallographic data. In this model the charge of the protein is assumed to be uniformly distributed, and is calculated from the amino acid composition; this charge is then compared with the value calculated from the ionic strength dependence of the rate constants.

Use of Eqn. 25 for the electrostatic work required to bring together a metalloprotein and a metal complex (or two metalloproteins) yields Eqn. 38 for the ionic strength dependence of the rate constant ¶:

$$\ln k = \ln \overline{k} - \frac{z_1 z_2 N e^2}{2 D_s \sigma_{12} RT} \left(\frac{\exp\left[-B(\sigma_{12} - \sigma_1)\sqrt{\mu}\right]}{1 + B\sigma_1 \sqrt{\mu}} + \frac{\exp\left[-B(\sigma_{12} - \sigma_2)\sqrt{\mu}\right]}{1 + B\sigma_2 \sqrt{\mu}} - 2 \right), \tag{38}$$

[¶] According to Eqns. 1-3 the ionic strength μ affects k via the work terms w^r and w^p and via ΔG^0 . The latter depends on μ because properties of the medium do. When $|\Delta G^{0'}/\lambda|$ is

where \bar{k} is the rate constant at infinite dilution, σ_{12} is the center-to-center separation of the two reactants, and σ_1 and σ_2 are defined as for Eqn. 25. In using Eqn. 38 simplifying assumptions for σ_1 and σ_2 are often made: (i) $\sigma_{12} = \sigma_1 + \sigma_2$ with $\sigma_i = a_i + 0.15$ nm [172] or (ii) $\sigma_{12} = \sigma_1 + \sigma_2$ with $\sigma_i = a_i$ [163,169–171]. The results obtained upon fitting the kinetic data to Eqn. 38 using the assumption (ii) and a least-squares routine in which \bar{k} and the total charge on the protein are allowed to vary are summarized in Table V. Included in Table V are the protein charges calculated from the amino-acid composition. The results are seen to be in satisfactory agreement, except for the result reported for the reaction of stellacyanin with Co(phen)₃³⁺, which apparently exhibits no ionic strength dependence §.

In another approach to ionic strength effects [172-175], it is assumed that the transition state of the reaction (denoted by AB[‡]) can be regarded as a single body rather than a pair of reactants. The rate constant can then be written as in Eqn. 39 using transition state theory

$$k = \overline{k} \gamma_{A} \gamma_{B} / \gamma_{AB^{\ddagger}}$$
 (39)

where γ_A , γ_B and $\gamma_{AB^{\ddagger}}$ denote the activity coefficients of A, B and AB[‡]. The quantity $\gamma_{AB^{\ddagger}}$ is assumed to be the activity coefficient of a sphere of radius σ_{12} and total charge $(z_1 + z_2)$. (Eqn. 39 was, in effect, obtained prior to transition state theory by Bronsted.) Expressing these activity coefficients in terms of Debye-Hückel theory, one obtains:

small, the dependence of $\ln k$ on μ is found from Eqns. 1-3 to be

$$\ln\ k = \ln\ \overline{k} + \frac{1}{2\,RT} \left(\,w^{\,\mathrm{t}} + w^{\,\mathrm{p}}\,\right) - \frac{1}{2} \ln \left(\,\gamma_1^{\,\mathrm{ox}} \gamma_2^{\,\mathrm{red}} / \gamma_1^{\,\mathrm{red}} \gamma_2^{\,\mathrm{ox}}\,\right)$$

for the case of reaction 7, where the γ 's denote activity coefficients. For self-exchange reactions w^r equals w^p , the activity coefficient ratio becomes unity, and use of Eqn. 25 then leads to Eqn. 38, where now $\sigma_1 = \sigma_2$. An equation more general than Eqn. 38 and arising from the above expression for $\ln k$ is given in the Appendix as Eqn. 60.

At the time this discrepancy was attributed, at least in part, to the presence of undefined amounts of carbohydrates in the preparation of the protein [159].

$$\ln k = \ln \bar{k} - \frac{Ne^2B\sqrt{\mu}}{2D_sRT}$$

$$\int z_s^2 \qquad z_s^2 \qquad (z_s + z_s)$$

$$\times \left(\frac{z_1^2}{1 + B\sigma_1\sqrt{\mu}} + \frac{z_2^2}{1 + B\sigma_2\sqrt{\mu}} - \frac{(z_1 + z_2)^2}{1 + B\sigma_{12}\sqrt{\mu}} \right)$$
(40)

At sufficiently low ionic strength Eqns. 38 and 40 both reduce to the same limiting form

$$\ln k = \ln \overline{k} + \frac{z_1 z_2 N e^2 B \sqrt{\mu}}{D_s R T} \tag{41}$$

In Eqns. 1-3, and thereby in Eqn. 38 and in Eqn. 60 of the Appendix, the interaction of the reactants A and B is treated as outer-sphere-like, i.e., they do not coalesce. For Eqn. 40, on the other hand, A and B are bound closely enough (or, instead, one ion is very small relative to the other) that they have in effect coalesced to form a sphere. In general, Eqns. 38 and 40 yield fairly similar results [170,172-175]: the difference between the expressions tends to become more marked at high ionic strengths, but Debye-Hückel-type theory also tends to break down under those conditions. The breakdown can be expected to be greatest for reactants of opposite charge because of specific ionic interactions [176], a point also noted later.

Eqns. 38 and 40 both assume that each reactant (and, in the case of Eqn. 40, AB[‡] also) is spherical and has a uniform charge distribution. When the charge distribution on a reactant is asymmetric, as is the case for most proteins, the departure from spherical symmetry has been modelled by introducing dipole (and higher order) interactions. In recent applications, Eqn. 38 has been corrected by adding an ion-dipole interaction energy to the ion-ion interaction energy (Eqn. 25) [174-177], while Eqn. 40 has been corrected by using the Kirkwood expression for the activity coefficient of a charged dipole, which involves adding another term [172] to the Debye-Hückel term. Use of these expressions requires information about the dipole moment of the protein.

The dipole moment of cytochrome c has been calculated to be ≈ 300 Debye [173]. For ionic strengths below 0.1 M, this dipole moment hardly

TABLE V

RESULTS OF CALCULATIONS OF THE CHARGE ON THE METALLOPROTEIN FROM THE IONIC STRENGTH DEPENDENCE OF THE ELECTRON-TRANSFER RATES

HAs denotes the ascorbate anion. The references cited in the last column contain both sets of calculations.

Protein	Reagent	рН	Charge		Refs.
			calculated from composition	calculated using Eqn. 38	
Cytochrome c (horse heart)	HAs ⁻	7.2	+8	+7.2	163
Cytochrome c (horse heart deriv.)	HAs ⁻	7.2	+6	+5.2	163
Cytochrome c-551 (Pseudomonas aeriginosa)	$Co(phen)_3^{3+}$	7.0	-3	-4.3	164
Cytochrome c-552 (Euglena gracilis)	HAs ⁻	7.2	-8	-7.3	163
Cytochrome c-552	$Fe(CN)_6^{4-}$	7.2	-8	-6.9	163
Cytochrome c-552	$Fe(CN)_6^{3}$	7.2	-9	-8.3	163
Cytochrome c (horse heart)	Fe(EDTA) ²⁻	7.0	+8	+10	169,170
Cytochrome c ₂ (Rhodospirillum rubrum)	Fe(EDTA) ²⁻	7.0	+ 3	+ 2.3	169,170
Azurin (Pseudomonas aeriginosa)	Fe(EDTA)2-	6.8	-1	-1.1	171
Plastocyanin (Phaseolus vulgaris)	Fe(EDTA) ²⁻	6.9	-8 a	-7.2	171
Stellacyanin (Rhus vernicifera)	$Co(phen)_3^{3+}$	7.0	+9	0.0	164

^a A charge of -9 is calculated from more recent sequence studies (subsection VF).

0.4 M the ionic strength dependence is mainly determined by the dipole interaction [174,175]. (The ion-ion interaction varies as the $\sqrt{\mu}$ while the ion-dipole interaction varies as μ .) If the charge on cytochrome c is +8, then the maximum correction, regardless of the ionic strength, to the charge on the protein was found to be ± 3 , the exact value depending on the cosine of the angle between the protein dipole and the interaction site on the protein surface [174,175].

In certain applications good agreement with the

makes any contribution to the ionic strength dependence of the rate, but at ionic strengths above

experimental ionic strength dependence has been obtained when the charge and radius of the presumed electron transfer site, rather than the total protein charge and radius, have been substituted into Eqns. 38 or 40 [166], or into more complex expressions [177]. In other applications the effective charge on the protein has been calculated from the ionic strength dependence of the binding of small anions or cations using the Bjerrum expression for the ion-pair formation constants. This procedure yielded good agreement between the effective charge and the charge of the binding site calculated from the protein structure [178], indicat-

ing that the binding constants in the systems

studied are determined by the local rather than by

the total protein charge (see discussion in Appendix IB).

In some cases the good agreement with the

experimental data found with one model or another

may be fortuitous, particularly if the local charge

has the same sign and happens to be of similar magnitude to the total protein charge. This is the case, for example, with horse-heart cytochrome c - the total charge on the oxidized protein is +8and the exposed heme edge (believed to be the electron transfer site, subsection VA) is surrounded by a ring of four positively charged amino acids. In the case of bean plastocyanin, the total charge of the oxidized protein is -8, while the electron-transfer site for the reaction of plastocyanin with negatively charged complexes is believed to be uncharged; nevertheless, the ionic strength dependence of the reactions of plastocyanin with Fe(EDTA)²⁻ [171] and Fe(CN)₆⁴⁻ [166] is consistent with reaction with a negatively charged protein. Presumably in these systems the approach of the negatively charged complex is

influenced by the protein charge nearest the elec-

tron-transfer site, which is a negatively charged

region in the case of plastocyanin (discussed fur-

ther in subsection VF), even though the presumed

electron transfer site itself is uncharged. On the

other hand, cytochrome f is an example of a

protein in which the charge on the binding site, as deduced from the effect of ionic strength on the reaction rate, is of opposite sign to the total protein charge [179].

As a general rule one expects that the total

protein charge will dominate the electrostatic interactions at large reactant separations, with the local protein charge becoming more important as the separation distance decreases. The 'cross-over' distance will depend upon the detailed charge distribution of the protein and is difficult to calculate, although some progress in modeling the electrostatic interactions is being made [177,180]. (In some systems, hydrophobic interactions may have to be included.) The results of recent calculates

distribution of the protein and is difficult to calculate, although some progress in modeling the electrostatic interactions is being made [177,180]. (In some systems, hydrophobic interactions may have to be included.) The results of recent calculations [177,180] for the reactions of a variety of c-type cytochromes support the view that a localized region of the cytochrome c surface is primarily responsible for the observed electrostatic interactions. This site is identified with the exposed heme region, an interpretation that is consistent with the results of other studies. These are described in the next section.

V. Applications to biological electron transfers

VA. Reactions of cytochrome c with metal complexes The reactions of cytochrome c with metal com-

plexes form an interesting bridge between the inorganic electron-transfer reactions in solution and purely biological electron transfers. Accordingly, we begin this section with a discussion of cytochrome c reactions and some of the relevant the-

ory.

TABLE VI

Cytochrome c is probably the best characterized metalloprotein [181–183]. The heme group lies in a crevice of the protein and is covalently bonded to the protein by thioether bridges. The iron atom is situated in the plane of the

porphyrin ring with the fifth and sixth coordina-

tion sites occupied by a ring nitrogen atom of

His-18 and the sulfur atom of Met-80. An im-

portant feature of the cytochrome structure is that

an edge of the porphyrin ring is located at the

surface of the metalloprotein. The rates of electron transfer between cytochrome c and a large number of metal complexes have been studied. The complexes used are substitution inert, ensuring an outer-sphere mechanism, and have known self-exchange rates [95,96,184–187] and redox potentials (Table VI). Some of the complexes are positively charged, others negatively charged; some are hydrophobic (e.g., Co(phen) $_3^{3+/2+}$), and others hydrophilic (e.g., Ru(NH $_3$) $_6^{3+/2+}$). The benzimidazole- and the pyridine-pentaammineruthenium complexes provide a large, aromatic ligand for probing hydro-

marized in Table VII, which also includes the values calculated from Eqn. 42, a common approximation to the cross-relation, Eqn. 10:

phobic protein regions such as the heme crevice.

The rate constants for the reactions are sum-

$$k_{12} \approx (k_{11}k_{22}K_{12})^{1/2}.$$
 (42)

The experimental value of $k_{11} = 1.2 \cdot 10^3 \,\mathrm{M}^{-1} \cdot \mathrm{s}^{-1}$ ($\mu = 0.1 \,\mathrm{M}$, pH 7) for the rate constant of the ferrocytochrome-ferricytochrome exchange reaction [193] and the values of k_{22} , the exchange rate

REDUCTION POTENTIALS, AVERAGE RADII, AND EXCHANGE RATE CONSTANTS OF METAL COMPLEXES AT $25^{\circ}\mathrm{C}$

The radii mentioned in column 3 are the average van der Waals radii of the reactants.

	E°	Radius	μ	k _{ex}	Refs.
	(V vs. NHE)	(nm)	(M)	$(\mathbf{M}^{-1} \cdot \mathbf{s}^{-1})$	
$\frac{\text{Co(phen)}_3^{3+/2+}}{\text{Co(phen)}_3^{3+/2+}}$	+0.36	0.70	0.1	45	167,95
$Ru(NH_3)_6^{3+/2+}$	+0.07	0.33	0.1	$3.2 \cdot 10^{3}$	96
$Ru(NH_3)_5 py^{3+/2+}$	+0.32	0.38	0.1	$1.1 \cdot 10^{5}$	168,184
$Ru(NH_3)_5BzIm^{3+/2+}$	+0.15	0.43	(0.1)	5 · 10 ⁴	185
$Fe(EDTA)^{1-/2-}$	+0.12	0.40	(0.1)	3 · 10 ⁴	186

TABLE VII

RATE CONSTANTS FOR HORSE-HEART CYTOCHROME c REACTIONS AT 25°C AND 0.1 M IONIC STRENGTH The K_{12} values are based on $E^{\circ} = +0.26$ V for the horse-heart cytochrome c couple [160]. The k_{12} calculates are rate constants

calculated in the prate constants do	present review from Eqn. 42 not include the W_{12} correct	l, using $k_{11} = 1.2 \cdot 10^3 \text{ M}$ tion. W_{12} is defined by E	⁻¹ ·s ⁻¹ for the cytochro cqn. 24; the individual w	ome c self-exchange work terms were co	ge [193]. The calculated from E	culated Eqn 25.
Reagent	K ₁₂	k _{12,obsd}	k _{12,calcd}	W ₁₂	Ref.	

Reagent	K ₁₂	$\begin{array}{c} k_{12,\text{obsd}} \\ (M^{-1} \cdot s^{-1}) \end{array}$	$\begin{array}{c} k_{12,\text{calcd}} \\ (M^{-1} \cdot s^{-1}) \end{array}$	W_{12}	Ref.
Co(phen) ₃ ³⁺	49	$1.5 \cdot 10^3$	1.6·10 ³	1.1	188
$Ru(NH_3)_6^{2+}$	$1.6 \cdot 10^3$	3.8 · 10 4	7.8 · 10 ⁴	1.9	189
$Ru(NH_3)_5 py^{3+}$	8.5	$9.6 \cdot 10^3$	$3.4 \cdot 10^{4}$	1.6	190
Ru(NH ₃) ₅ BzIm ²⁺	72	5.8 · 10 4	6.6 · 10 4	1.3	191
Fe(edta) ²	$2.3 \cdot 10^{2}$	2.6 · 10 4	$9.1 \cdot 10^4$	9.1	192
$Co(ox)_3^{3-}$	$1.7 \cdot 10^{5 \text{ a,b}}$	5.5 a	7.0 a,c	15 ª	194

^b Calculated using $E^{\circ} = +0.57 \text{ V}$ for the $Co(0x)_3^{3-/4}$ couple [194]. ^c Calculated using $k_{22} = 2.4 \cdot 10^{-7} \text{ M}^{-1} \cdot \text{s}^{-1}$ for the Co(ox) $\frac{3}{3}$ exchange at 0.5 M ionic strength. This value of k_{22} was calculated from the rate constant for the oxidation of $Co(terpy)_2^{2+}$ by $Co(ox)_3^{3-}$ reported in Ref. 190 using Eqn. 22.

constants for the metal complexes given in Table

VI, were used. The W_{12} correction, which was not

used, is also given, and is seen in the table to be

close to unity, except for the reaction of ferricy-

tochrome c with the negatively charged reactants

Fe(EDTA)²⁻ and Co(ox)³⁻, for which $W_{12} \approx 9$ and $W_{12} \approx 15$, respectively. The agreement of the observed and calculated exchange rates in Table VII is seen to be good (apart from that for the reactions with Fe(EDTA)²⁻ and Co(ox)³⁻ when W_{12} is included). The simplest explanation is that cytochrome c uses the same electron-transfer pathway with metal

complexes as it does in its self-exchange reaction, and that this pathway is the exposed heme edge. Additional support for this mechanism has re-

cently been obtained from protein modification

report [199] that the Fe(CN) $_{6}^{4-}$ -ferricytochrome c system exhibits an intervalence absorption band has also been shown to be erroneous: the observed band was due to Prussian-blue type impurities

and although rate saturation behavior has been

reported at high concentrations of the complex,

that interpretation is complicated by the need for

corrections for specific ion interactions as well as

for the reversibility of the reaction [196,197] \(^1\). At

this time there appears to be no good example of rate saturation in the electron transfer reactions of

cytochrome c with metal complexes [197,198]. A

[201].

VB. The cytochrome c self-exchange reaction

The cross-reaction studies discussed above are clearly consistent with the cytochrome c self-exchange rate of approx. $1 \cdot 10^3 \text{ M}^{-1} \cdot \text{s}^{-1}$ at 0.1 M

In terms of the model, the cytochrome c self-exchange rate constant is given by (Appendix IB)

studies [195]. One sees that there is no special rate enhancement with Ru(NH₃)₅BzIm²⁺ or ionic strength determined by NMR [193]. In this Ru(NH₃)₅py³⁺ ascribable to penetration of the section we interpret this rate constant in terms of heme crevice by the π -conjugated ligand (Table the quantities introduced in Section II. VIII). This last result is in contrast to an earlier

conclusion [190], which was based on the use of too small an exchange rate constant for the Ru(NH₃)₅py^{3+/2+} couple. The reaction of ferrocytochrome c with $Co(ox)_3^{3-}$ is reasonably wellbehaved at the high ionic strength (0.5 M) used for the studies. This is not the case for the reaction of cytochrome c with ferro- and ferricyanide. There

have been numerous studies of the latter reaction,

 $k = SK_A \kappa \nu \exp \frac{-\lambda}{4RT}$ (43) ¶ The redox potential and self-exchange rate of $Fe(CN)_6^{3-/4-}$ couple is highly medium-dependent [187] and this couple is

therefore not a good choice for cross-reaction studies.




Fig. 9. Computer-generated complexes formed from two cytochrome c molecules showing the coplanarity of the heme groups. Upper set: the two His-18 residues are cis relative to the Fe-Fe axis; lower set: the two His-18 residues are trans relative to the Fe-Fe axis. The figure was kindly provided by Dr. P.C. Weber of Genex Corporation.

where $\lambda = \lambda_i + \lambda_o$. A steric factor S has been introduced to allow for the fact that the cytochrome c molecule is not spherically symmetric, at least so far as the angular dependence of the electron-transfer probability is concerned. The quantity SK_A is the equilibrium constant for formation of a precursor complex A-B in which the active sites of A and B are in contact (Appendix IB).

We consider the various factors appearing in Eqn. (43): if it is assumed that the electron-transfer reactions of cytochrome c proceed through the exposed, but recessed heme edge, then a lower limit for the steric factor for the exchange is equal to the square of the ratio of the surface area of the exposed heme edge to the surface area of the whole molecule [22]. Cytochrome c is a prolate spheroid approximately $2.5 \times 2.5 \times 3.7$ nm in size, corresponding to a surface area of 26 nm². The effective area of the heme crevice is about 0.5×1.8 nm [200]; accordingly, the heme edge occupies about 3% of the surface of the molecule. We shall tentatively adopt a vlaue of 0.01 for S in subsequent calculations \P . The K_A for this large system ($\sigma = 2.84$ nm) is estimated with Eqns. 25 and 57 to be approx. 1 M⁻¹. As before, Eqn. 37b of Section IVC is used for $\kappa \nu$.

Mixing of the t_{2g} -orbitals of the Fe center in cytochrome c with the π^* -orbitals of the porphyrin ring effectively extends the metal d-electron density to the porphyrin edge, so that facile electron transport to and from the iron center via the exposed heme edge should be possible. However, this does not mean that the exchange reaction will be adiabatic \S , i.e., that $\kappa \approx 1$. The 'exposed' edge of the heme group of cytochrome c is surrounded by folds in the polypeptide chain: these folds may

prevent the close (edge-to-edge) approach of the hemes required for $\kappa \approx 1$. Computer modeling (Weber, P.C. (1984) personal communication to Sutin, N) of the approach of two cytochrome c molecules reveals that the heme groups are able to approach closest in a co-planar configuration. Two such complexes, in which the His-18 residues are cis and trans, respectively, relative to the Fe-Fe axis, are shown in Fig. 9. In these complexes the heme edge-to-edge (carbon-carbon) separation is 0.94 nm and the iron-iron distance is 1.77 nm. (Interestingly, this value is close to that found for two similarly docked cytochromes, 1.6 nm, in an independent study - an X-ray structure determination of a photosynthetic reaction center discussed in subsection VD.) Although there are more short heme-heme distances in the cis complex, this advantage is offset by the lower electrostatic repulsion in the trans complex. By movement of the side chains of the surface amino acids the separation of the heme groups can be decreased from 0.94 nm to approx. 0.9 nm, but cannot readily be reduced much further (Weber, P.C. (1984) personal communication to Sutin, N.). For purposes of calculation, we use the former value for the heme edge-to-edge (carbon-to-carbon) separation in the cytochrome c exchange, and use the 0.3 nm correction discussed in subsection IVB to obtain an electron-transfer distance d of 0.64 nm. With this value of d and $\beta = 12$ nm⁻¹, a

value of $\kappa\nu \approx 5\cdot 10^9~{\rm s}^{-1}$ would be estimated ¶. Next we consider the values of λ_i and λ_o . X-ray crystallographic studies (0.18 nm resolution) indicate minimal differences between the structures of the heme group in oxidized and reduced cytochrome c [181,182]. A crystal of ferricytochrome c (but not of ferrocytochrome c) contains two molecules per unit cell: the average Fe to

this does not mean that the exchange reaction will be adiabatic \S , i.e., that $\kappa \approx 1$. The 'exposed' edge of the heme group of cytochrome c is surrounded by folds in the polypeptide chain: these folds may

In general, it is necessary to integrate k(r) over all mutual orientations and distances of the reacting pair. We adopt a simplified model in which the electron transfer through the remainder of the protein molecule may be neglected in comparison with the rate through the heme edge. Because of electron transfer from other angles the actual S is expected to be closer to unity than the square of the ratio of effective areas $(0.03)^2$ and we have adopted the value of 0.01.

Nor does the fact that the cross-reactions satisfy Eqn. 42

Nor does the fact that the cross-reactions satisfy Eqn. 42 require that the exchange reaction be adiabatic; Eqn. 42 will still be satisfied provided $\kappa_{12} \approx (\kappa_{11}\kappa_{22})^{1/2}$ or, more generally, provided that $S_{12}\kappa_{12} \approx (S_{11}S_{22}\kappa_{11}\kappa_{22})^{1/2}$ [22].

Alternative electron-transfer pathways are not considered. For example, mediation by the sulfur atoms of Cys-14 and Cys-17, which are covalently bonded to the exposed heme edge, could be important [202]. In this regard the sulfur atom of Cys-17 is particularly well-situated (Fig. 9) and extended Hückel calculations (Hanson, L.R. (1984) personal communication to Sutin, N.) indicate that the Fe electron density on the cysteinyl sulfurs may be comparable to that on the heme-edge carbon atoms. Mediation by the sulfur would decrease the effective d and increase κν.

pyrrole-N distances in the two molecules are 0.208 \pm 0.011 and 0.203 \pm 0.011 nm, while in ferrocytochrome c this distance is 0.206 ± 0.007 nm; the Fe to His-18N distance is 0.196 ± 0.011 and $0.204 \pm$ 0.011 nm in the two ferricytochrome c molecules and 0.197 ± 0.007 nm in the reduced molecule, while the Fe to Met-80S distance is 0.228 ± 0.005 nm and 0.226 ± 0.005 nm in the two oxidized molecules and 0.232 ± 0.003 nm in the reduced molecule [182]. Although the iron-ligand distances in the two oxidation states appear to be similar the uncertainty is too large to permit a good estimae of λ ./4, the reorganization energy arising from changes in the metal ligand bond distances. A small value of $\lambda_i/4$ (approx. 4-8 kJ·mol⁻¹) is suggested by the rapid self-exchange rates of free (i.e., without bound protein) low-spin § hemes in solution, and is not inconsistent with the X-ray data. This value of λ_i for a pair of cytochromes is also consistent with the Stokes shift (8 kJ·mol⁻¹) [204] for the low-frequency modes (200-500 cm⁻¹ range) that are coupled to the heme π - π * transition in a single ferrocytochrome c. (If these lowfrequency modes provided the only contribution to λ_i for the self-exchange, such a Stokes shift would yield $\lambda_i = 8 \times 2 \text{ kJ} \cdot \text{mol}^{-1}$).

For the metalloprotein reactions there are, in general, two contributions to λ_0 , one arising from the change in polarization (orientations of protein dipoles) of the polypeptide and the second from the change in polarization of the surrounding solvent. In general, the relative contributions to λ_0 from these two sources will depend upon the nature of the particular metalloprotein. For cytochrome c, the contributions of the reorientation of the protein dipoles and of the surrounding water molecules to λ_0 (for the pair of cytochrome reactants) have been calculated from a microscopic model to be 10 and 12 kJ·mol⁻¹, respectively, and λ_i was estimated to be 4 kJ · mol⁻¹ [137]. The resulting value of 6.5 kJ \cdot mol⁻¹ for $\lambda/4$ seems too low. An alternative approach would be to assume that the protein/water surrounding the heme can be treated as a continuous dielectric with $D_s = 10$ and $n^2 = 2$. If Eqn. 6 is then used with a, the radius of the heme, equal to 0.5 nm and r, the iron-to-iron distance, equal to 1.8 nm, $\lambda_o/4$ becomes $20 \text{ kJ} \cdot \text{mol}^{-1}$. A total value of $25 \text{ kJ} \cdot \text{mol}^{-1}$ for $\lambda/4$ for the Cyt c-Cyt c self-exchange is consistent with the activation energy of $29 \pm 4 \text{ kJ} \cdot \text{mol}^{-1}$ at $\mu = 1 \text{ M} [193]^{\dagger}$ and will be assumed here, together with an S of 0.01 for that reaction and $\beta = 12 \text{ nm}^{-1}$. Considering the complexities of the systems and the current uncertainties in their molecular parameters, a factor of ten difference in calculated and observed rates (which is equivalent to a change of $5.8 \text{ kJ} \cdot \text{mol}^{-1}$ in the free-energy barrier) can be regarded as good agreement.

The particular data which will be considered using the above values \P of β , S and λ are: (a) the self-exchange rate constant of the cytochrome c system; (b) the self-exchange for the protein-free hemes; (c) the bimolecular electron transfer between ferricytochrome c and $Ru(NH_3)_5His^{2+}$; (d) the intramolecular (NH₃)₅RuHis-Cyt c electron transfer, its small activation energy and its very negative entropy of activation; (e) the maximum bimolecular rate constant for electron transfer from Cyt c to various metal complexes; (f) the cytochrome c-chlorophyll dimer electron transfer in a photosynthetic reaction center; (g) the relative rates of the Cyt c-Cyt b₅ and Cyt c-Cyt c reactions; and (h) electron transfer between cytochrome c and a number of other metalloproteins. The bulk of these results and several others are collected later in Table VIII.

With the above values for $S(\approx 10^{-2})$, $K_A(\approx 1 \text{ M}^{-1})$, $\kappa\nu(\approx 5\cdot 10^9 \text{ s}^{-1})$ and $\lambda/4$ ($\approx 25 \text{ kJ}\cdot \text{mol}^{-1}$), the self-exchange rate constant for cytochrome c is estimated from Eqn. 43 to be $10^{-2}\cdot$

[§] The value of λ_i/4 is likely to be larger for high-spin than for low-spin hemes because of the larger changes in metal to axial-ligand distances expected for high-spin systems.

[†] The activation energy at $\mu = 0.1$ M is twice as large [193]; perhaps there is a large temperature-dependent salt effect at low ionic strengths. Debye-Hückel type expressions appear to be more valid for calculating free energy changes (e.g., activity coefficients or work terms) than for calculating their derivatives, such as entropies or energies (cf. results in Ref. 76).

Predicted results in Section V, such as the Cyt c-Cyt c self-exchange rate or the cross-reaction rate at small $|\Delta G^{0'}/\lambda|$ do not depend on the individual values of S and λ (S_{11} and λ_{11}), but only on the combination S_{11} exp ($-\lambda_{11}/4RT$), [cf. Eqn. 43]. Predicted results such as an intramolecular electron transfer k or its activation energy depend on $\lambda_{11}/2$, but not on S_{11} , while the predicted k in the barrierless regime depends on $S_{11}^{1/2}$, but not on λ_{11} .

 $5 \cdot 10^9 \text{ e}^{-10}$ i.e., approx. $2 \cdot 10^3 \text{ M}^{-1} \cdot \text{s}^{-1}$. The experimental value is $1.2 \cdot 10^3 \text{ M}^{-1} \cdot \text{s}^{-1}$ [193].

The rate constant k for heme-heme self-exchange is $1 \cdot 10^6 - 1 \cdot 10^{10}$ M⁻¹·s⁻¹ depending on the nature of the axial ligands (through their effect on λ_i and λ_o) and the medium [205]. The value of $\lambda_o/4$ for heme exchange will depend on the nature of the solvent, but is expected to be small because of the relatively large size of the heme group. The factor of more than 104 difference in the cytochrome c and free heme self-exchange rates is then primarily due to the extra separation distance between the hemes in the protein-protein case §. The large separation was estimated above the introduce a factor of $5 \cdot 10^{-4}$ into the k for the Cyt c-Cyt c exchange. In general, the value of k for a hemeprotein self-exchange will be sensitive to the degree of exposure of the heme and, all other factors being equal, changing the exposure of the heme by changing the protein conformation or the number of amino acids is likely to dramatically affect the exchange rate of the cytochrome. For example, Pseudomonas aeruginosa cytochrome c-551 has only 82 amino acid residues compared with the 104 residues of horse-heart cytochrome c, and has a self-exchange rate of $1.2 \cdot 10^7 \text{ M}^{-1} \cdot \text{s}^{-1}$ [206], instead of $1.2 \cdot 10^3$ M⁻¹·s⁻¹. A part of this difference may be due to the smaller size of Cyt c-551, which makes the hemes more accessible, and part may be due to a difference in work terms, e.g., the electrostatic repulsion in the Cyt c-551 self-exchange is smaller (cf. Table V for the ionic

VC. Intramolecular electron transfer

charges).

Rate constants calculated with the above parameters can be compared with data on the intramolecular electron transfer between ferricy-tochrome c and a bound $Ru(NH_3)_3^2$ residue.

We begin with the value of the bimolecular rate constant for reduction of ferricytochrome c by $Ru(NH_3)_5His^{2+}$, $9 \cdot 10^4$ M⁻¹·s⁻¹ [207]. When

 $|\Delta G^{0'}|\lambda$ is small, the theoretical rate constant for this reaction is given by (Appendix IB):

$$k \cong SK_{A}\kappa\nu \exp\left(\frac{-\lambda}{4RT} - \frac{\Delta G^{\prime\prime}}{2RT}\right),$$
 (44)

where k is measured in $M^{-1} \cdot s^{-1}$ and where $S\kappa\nu$ $= 2 \cdot 10^{10} \text{ s}^{-1}$, based on $S = (0.01)^{1/2}$ and $\kappa \nu$ calculated from $1 \cdot 10^{13} \exp(-\beta d)$ with $\beta = 12$ nm^{-1} and an electron-transfer distance d ≈ 0.32 nm. (This distance is half the value used in subsection VB for the cytochrome c self-exchange; the electron is assumed to be delocalized to the edge of the electronic cloud of the hydrogen atoms on the periphery of the Ru(NH₃)₅His complex.) For $\sigma = 1.8 \text{ nm} \text{ and } \mu = 0.1 \text{ M}, K_A \text{ is about } 0.4 \text{ M}^{-1}$ (including only electrostatic contributions to w). Using the parameters for the cytochrome c exchange estimated above and Ru(NH₃)₅py^{3+/2+} [96] as a model for the $Ru(NH_3)_5His^{3+/2+}$ exchange, $\lambda/4$, and $\Delta G^0/2$ are calculated to be 27.2 and $-8.4 \text{ kJ} \cdot \text{mol}^{-1}$, respectively ¶. Substituting these values into Eqn. 44 gives $k_{\rm calc} \approx (2 \cdot 10^{10})$ (0.4) $e^{-7.5}$, i.e., $4 \cdot 10^6$ M⁻¹·s⁻¹. The observed value is $9 \cdot 10^4 \text{ M}^{-1} \cdot \text{s}^{-1}$ [207].

This experimental rate constant of $9 \cdot 10^4 \, \mathrm{M}^{-1} \cdot \mathrm{s}^{-1}$ may be converted to a first-order constant k_1 for electron transfer at the heme edge by setting k_1 equal to k/SK_A (Appendix IB). With $S \approx (0.01)^{1/2}$ and $K_A = 0.4 \, \mathrm{M}^{-1}$ for the cited σ (cf. Eqn. 57 of Appendix I with w^r , assumed for simplicity to be nonelectrostatic, given by Eqn. 25) the first-order rate constant k_1 equals $2 \cdot 10^6 \, \mathrm{s}^{-1}$. This k_1 is introduced into Eqn. 45 below.

By the use of imaginative synthetic techniques, a $Ru(NH_3)_5$ -residue has been covalently attached to His-33 of ferricytochrome c and the intramolecular electron-transfer rate from the ruthenium(II) to the heme iron(III) measured [151–157]. The closest distance from the imidazole group of His-33 to the imidazole group that is coordinated to the

The respective values of $S\sigma^2$ also enter. Because of orientation requirements the S for the free heme system may be less than unity. Consequently, we estimate that $S\sigma^2$ is 10^{-2} (3)² nm² and less than 1 nm² for the protein-bound and protein-free heme systems, respectively.

The value of λ for the cross-reaction was taken equal to the mean of the λ values for the corresponding exchange reactions. For the Cyt c-Cyt c exchange the value of $\lambda/4 \approx 25$ kJ·mol⁻¹ mentioned earlier was used, while $\lambda/4$ for the Ru(NH₃)₅py^{3+/2+} couple is about 29 kJ·mol⁻¹ [96]. The $\sigma = 1.8$ nm is the sum of the radii of the two reactants (1.42 nm plus 0.38 nm). ΔG^0 , defined in subsection IIA, depends on ionic strength, as do the w's.

heme iron is 1.18 nm [155] and the measured intramolecular rate constant is $30 \pm 3 \text{ s}^{-1}$ by flash photolysis [155] and $53 \pm 2 \text{ s}^{-1}$ by pulse radiolysis [156] techniques. The rate constant may be compared with $2 \cdot 10^6 \text{ s}^{-1}$ estimated above for 'intramolecular' electron transfer at the heme edge (the driving force, $-\Delta G^0$, for the two electron transfers is similar). It is seen that the adjacent heme-edge pathway is approx. 10^5 times more favorable than the remote His-33 pathway, which involves electron transfer through the protein.

Assuming that λ , ΔG^0 , and β are the same for the adjacent and remote pathways, the rate constant k for the remote pathway can be estimated from k_1 and the change in separation distance:

$$k = k_1 \exp(-\beta \Delta d) \tag{45}$$

where k is measured in s⁻¹ and $k_1 \approx 2 \cdot 10^6$ s⁻¹. The difference in the electron-transfer distances for the adjacent and remote pathways $\Delta d \approx 0.7$ nm § and $\beta = 12$ nm⁻¹. Substitution into Eqn. 45 gives $k_{\rm calc} \approx 500$ s⁻¹, which is to be compared with the observed value of 30-53 s⁻¹.

In addition to the rate constant, the temperature dependence of the intramolecular ferricytochrome c to Ru(NH₃)₅His²⁺ rate was also measured. The flash photolysis results give $\Delta H^* \leq 6.3$ $kJ \cdot mol^{-1}$ and $\Delta S^{\dagger} = -200 \pm 8 J \cdot deg^{-1} \cdot mol^{-1}$ [155], while the pulse radiolysis studies yield ΔH^* = 14.6 \pm 0.8 kJ·mol⁻¹ and $\Delta S^{\dagger} = -163 \pm 4$ J· deg⁻¹·mol⁻¹ [156]. Although there is some disagreement, ΔH^* for both studies is relatively small while the entropy of activation ΔS^{\dagger} is large and negative. The low value of ΔH^* derives, we believe, largely from the negative ΔH^0 for the reaction: The reorganization enthalpy at the separation distance r is related to the standard enthalpy change of the reaction $\Delta H^{0'}$ at that separation distance in the prevailing medium by [60] #:

$$\Delta H_{12}^* = \frac{\Delta H_{11}^*}{2} + \frac{\Delta H_{22}^*}{2} + \frac{\Delta H^{0'}}{2}, \tag{46}$$

where ΔH_{11}^* and ΔH_{22}^* , the reorganization enthalpies of the component exchange reactions, are approximately equal to $\lambda_{11}/4$ and $\lambda_{22}/4$, respectively. From the value of $\Delta H^0 = -49.8 \text{ kJ} \cdot \text{mol}^{-1}$ [155] * and the values used earlier for the λ 's $(\Delta H_{11}^* = 25 \text{ kJ} \cdot \text{mol}^{-1} \text{ and } \Delta H_{22}^* = 29 \text{ kJ} \cdot \text{mol}^{-1}$ [96]), one finds $\Delta H_{12}^* = 2.1 \text{ kJ} \cdot \text{mol}^{-1}$, consistent with the small observed [155,156] value. A similar analysis for the corresponding azurin-Ru(NH₃)₅His²⁺ molecule has just been published [208]. Finally, in terms of the present model, the entropy of activation ΔS^{\dagger} for the intramolecular electron transfer can be interpreted using the expression, obtained from Eqns. 36c, 36d and 37b:

$$\Delta S_{12}^{\dagger} \cong \frac{\Delta S^{0'}}{2} - R\beta d, \tag{47}$$

neglecting the $\Delta G^{0'}/\lambda$ term. The value of $\Delta S^{0}/2$ is $-52.3~\mathrm{J\cdot deg^{-1}\cdot mol^{-1}}$ [155] ¶ and $-R\beta d$ equals $-100~\mathrm{J\cdot deg^{-1}\cdot mol^{-1}}$ (calculated, as above with $\beta=12~\mathrm{nm^{-1}}$ and $d=1.0~\mathrm{nm}$). One thus finds $-152.8~\mathrm{J\cdot deg^{-1}\cdot mol^{-1}}$ for the calculated ΔS_{12}^{\dagger} , while the observed value is -163 [156] or -200 [155] $\mathrm{J\cdot mol^{-1}\cdot deg^{-1}}$.

It is also useful to examine experimental data on the maximum value of the rate constant observed in the barrierless $(-\Delta G^{0'} \approx \lambda)$ regime for the electron transfer from Cyt c to various reactants and to compare this rate constant with calculated values. For the reaction of ferricytochrome c with the charge-transfer excited state of Ru(bpy)₃²⁺, $(\Delta G^0 \approx -1.1 \text{ V})$ the observed rate constant is $2.5 \cdot 10^8$ M⁻¹·s⁻¹ at 0.1 M ionic strength [209]. For the reaction of ferrocytochrome c with $Ru(bpy)_3^{3+}$, $Ru(phen)_3^{3+}$, $Os(bpy)_3^{3+}$ and Os(phen)₃³⁺, all reactions having large negative $\Delta G^{0'}$'s, the rate constants are 1.2, 0.65, 1.2 and 0.76, respectively, in units of $10^8 \text{ M}^{-1} \cdot \text{s}^{-1}$. The maximum value of the 'activation-controlled rate constant', $SK_A\kappa\nu$ (Appendix IB) can be calculated from the parameters estimated earlier to be approx. $8 \cdot 10^9$ M⁻¹·s⁻¹. The diffusion-controlled rate constant (k_D in Eqn. 53b of Appendix IA) for the reaction of the cytochrome c's with large +3

This value assumes the shortest path for electron transfer to be a linear one. Another possibility is a more tortuous path along a polypeptide backbone.

[#] Eqn. 46 follows from Eqn. 35 when $|\Delta G^0'/\lambda|$ is small compared to unity. In the present case ΔG^0 is $-17 \text{ kJ} \cdot \text{mol}^{-1}$ and λ is $108 \text{ kJ} \cdot \text{mol}^{-1}$.

[¶] The difference between ΔH^0 and the $\Delta H^{0'}$ in Eqn. 46 is negligible in the present system for the remote pathway, as is that between ΔS^0 and the $\Delta S^{0'}$ in Eqn. 47.

TABLE VIII ABSOLUTE RATE CONSTANTS OR ACTIVATION PARAMETERS FOR SEVERAL CYTOCHROME $\it c$ REACTIONS

Reaction	$\frac{\Delta G^0}{(kJ \cdot mol^{-1})}$	Parameter	Observed	Calculated	Subsection
Cyt c^{II} + Cyt c^{III}	0	$E_a (\mu = 1 \text{ M}) (k \text{J} \cdot \text{mol}^{-1})$	29 ± 4	25 a	VB
ejte , ejte	0	$k (\mu = 0.1 \text{ M}) (M^{-1} \cdot \text{s}^{-1})$	$1 \cdot 10^{3}$	$2 \cdot 10^{3}$	VB
Cyt c^{III} + Ru(NH ₃) ₅ His ²⁺	-17	$k \left(\mathbf{M}^{-1} \cdot \mathbf{s}^{-1} \right)$	9.104	$4 \cdot 10^6$	VC
Cyt $c^{III} \sim \sim \text{Ru}(NH_3)_5^{2+}$	-17	$k(s^{-1})$	30-50	500	VC
Cyr c 166(.1113)5	-50 (≤4) b	$\Delta H^* (kJ \cdot mol^{-1})$	146,56	2	VC
	$-105(-202)^{\circ}$	$\Delta S^* (\mathbf{J} \cdot \deg^{-1} \mod^{-1})$	-163,-20 0	-152	VC
Cyt c^{II} + ML_3^{3+d}	-59 to -84	$k_{\max} \left(\mathbf{M}^{-1} \cdot \mathbf{s}^{-1} \right)$	≈ 10 ⁸	3 · 10 9 e	VC
Cyt $c^{II} \sim \sim BChl_2^+$	≈ -16	$k (s^{-1})$	$4 \cdot 10^6$	$4 \cdot 10^{7}$	VD

^a This is the value estimated for $\lambda/4$.

and +2 metal complexes is about 3.2^i and $3.4 \cdot 10^9$ M⁻¹·s⁻¹, respectively, at 0.1 M ionic strength, using Eqn. 25 for the work term. One sees that the maximum experimental value (approx. 10^8 M⁻¹·s⁻¹) is substantially less than the diffusion-controlled value ($3 \cdot 10^9$ M⁻¹·s⁻¹). Perhaps it represents an experimental result for the value of $SK_A\kappa\nu$ for these Cyt c-metal complex systems (the estimated theoretical value above was approx. $8 \cdot 10^9$ M⁻¹·s⁻¹). Some of the rate constants and other kinetic parameters that have been discussed in this section are summarized in Table VIII.

VD. Photosynthetic reaction centers

A *c*-type cytochrome is the electron donor to the oxidized bacteriochlorophyll dimer in the reaction centers of photosynthetic bacteria [211,212]:

$$c^{2+} + (BChl)_2^+ \rightarrow c^{3+} + (BChl)_2$$
 (48)

We first consider kinetic data on reaction (Eqn. 48) for *Chromatium vinosum* and then consider the *Rhodopseudomonas viridis* reaction center complex, for which a crystallographic study has just become available, and for which there are kinetic data.

The rate of oxidation of ferrocytochrome c by $(BChl)_2^+$ in the reaction center of *Chromatium vinosum* is temperature-independent in the range 100-4 K ($k=3\cdot 10^2 \text{ s}^{-1}$); at 300 K the electron

transfer rate constant is about 10^6 s^{-1} , the pre-exponential factor appears to be $6 \cdot 10^8 \text{ s}^{-1}$, and the activation energy to be 17.6 kJ·mol⁻¹ [19,212]. $\Delta G^{0'}$ for the reaction is $-43.5 \text{ kJ} \cdot \text{mol}^{-1}$ [19,213]. From EPR measurements the center-to-center distance of Cyt c and $(B\text{Chl})_2^+$ is at least 2.5 nm, corresponding to an edge-to-edge separation of 1.2–1.8 nm [214]. These two sets of results are examined next.

First we consider the high-temperature pre-exponential factor for the reaction in terms of the above model. This factor may be obtained from Eqn. 36b and hence equated (approximately) with $10^{13} \exp(-\beta d)$, neglecting any $|\Delta G^{0'}/\lambda|$ compared with unity in Eqn. 35 and neglecting $\Delta S^{0'}/R$ in Eqn. 36b; substitution of $\beta = 12 \text{ nm}^{-1}$ leads to d = 0.8 nm, and so a nearest carbon to nearest carbon distance of 1.1 nm is calculated using the correction in subsection IVB. The value of 1.1 nm is close to the lower limit of the range of the separation distances (1.2–1.8 nm) implicated by the EPR measurements [214] $^{\$}$, it would be useful to determine ΔS^0 for reaction 48, it having been neglected in the above calculation.

^b Value of ΔH^0 in kJ·mol⁻¹.

^c Value of ΔS^0 in J·deg⁻¹·mol⁻¹.

^d Tris polypyridine complexes of osmium(III) and ruthenium(III).

 $^{^{\}circ}$ Estimated diffusion-controlled value. Estimated activation-controlled value is $8\cdot 10^9~M^{-1}\cdot s^{-1}$.

In a quantum mechanical calculation an H_{AB} in Eqn. 12 was inferred from the experimental pre-exponential factor to be $1090 \text{ kJ} \cdot \text{mol}^{-1}$ [215,216]. The calculation in Refs. 215, 216 also explored a fitting of the detailed temperature dependence of the rates. In the model used [215,216], both medium (solvent) and intramolecular modes were included.

We turn next to the photosynthetic reaction center of Rhodopseudomonas viridis. The arrangement of the components has recently been determined from an X-ray crystal structure at 0.3 nm resolution [217]. In addition to protein, the crystallized reaction center complex contains four bacteriochlorophyll-b and two bacteriopheophytin-b molecules, four c-type cytochromes, one non-heme iron and one menaquinone. Two of the four BChl's are present as (BChl)₂ ('special pair') [218-220]), in which their pyrrole rings I are stacked on top of each other about 0.3 nm apart. The distance between BChl Mg atoms in the special pair is approx. 0.7 nm and the angle between the ring planes is about 15°. To each BChl of the special pair is associated another BChl, whose Mg atom is 1.3 nm from the Mg atom of the special pair member, and the angle between the porphyrin planes of that member and the BChl is approx. 70°. Each of these monomeric BChl molecules is in turn also associated with a BPh molecule: the distance between the centers of their rings is 1.1 nm, and the angle between the planes of their rings is approx. 64°. (This arrangement, in which there is a site between the special pair and the BPh had been predicted earlier so as to explain the effect § of a small magnetic field on the reaction kinetics for the closely related Rhodopseudomonas sphaeroides [221].) One of the BPh's is also in contact with a side chain of the menaquinone *. The latter is, in turn, close to the non-heme iron.

A rectangular distribution of medium modes with a mean energy and width of 200 cm⁻¹ was used, corresponding to $\lambda_o = 23.9 \text{ kJ} \cdot \text{mol}^{-1}$. The frequency used for the intramolecular mode was 500 cm⁻¹ with $\lambda_i = 222 \text{ kJ} \cdot \text{mol}^{-1}$. These parameters provided a good fit of the temperature dependence of the rate constants, except at the transition to temperature independence, which is more abrupt than calculated. However the λ_i value seems unreasonably high – certainly much higher than λ_i for the reactions of Cyt c with typical metal complexes.

The four BChl's, two BPh's, the menaquinone and the non-heme iron are believed to be situated within the photosynthetic membrane of the bacterial cell, with the non-heme iron near the inner membrane surface and the special pair near the outer surface. The four cytochrome c molecules are located outside the membrane and are positioned roughly in a straight line in the crystal. The heme-iron of the cytochrome c closest to the special pair is approx. 2.1 nm from each of the two magnesium centers of the BChl's of the (BChl)₂. The four heme groups (of the cytochromes) can be viewed as forming two pairs, the pair closest to the BChl is related to the outer heme pair by rotation about an approximately 2-fold rotation axis. Within each pair the angle between the heme planes (i.e., the angle between the planes of the first and second hemes as well as the angle between the planes of the third and fourth hemes) is 78.5° and the iron-iron distance is approx. 1.4 nm. The angle between the planes of the second and third hemes is 17.4° and the distance between their iron centers is approx. 1.6 nm.

Flash-induced redox titrations have shown that two different c-type cytochromes, designated cytochromes, c-553 and c-558 to indicate the positions of their α -band maxima, are present in the Rps. viridis reaction center [222]. The midpoint potentials of the cytochromes are -12 mV and +330 mV, respectively [222,223]. The midpoint potential of the BChl-b dimer is +500 mV at pH 8 [223]. Both types of Cyt c can reduce the (BChl b) $_{2}^{+}$ at room temperature but only c-553 is capable of reducing it at 77 K [211,223]. The immediate electron donor to the oxidized dimer at room temperature is the Cyt c-558 [223] *.

We first consider electron transfer within the linear arrangement of Cyt c molecules. The separation of the iron centers of adjacent cytochromes

[§] An alternative explanation has been given [129]; it presumes that the decay constant β in Eqn. 37 is quite sensitive to orbital energies, a result inferred from calculations with delta function and square-well potentials. However, as discussed in subsection IIID, more recent calculations [71] show little difference between the optical and thermal matrix elements.

^{*} A second quinone molecule expected for the other BPh molecule was not present and may have been lost during the preparation of the crystal [217].

We are grateful to Drs. J. Deisenhofer and H. Michel for this information.

^{*} If the approximately linear arrangement of the cytochromes in the crystal of the reaction center also obtains in the bacterial cell, then the high-potential cytochrome (Cyt c-558) may be the only direct electron donor to the (BChl)½; the apparent reduction of (BChl)½ by the low-potential cytochrome (c-553) then results from very rapid reduction of the Cyt c-558 by the Cyt c-553. This sequence has recently been entertained for Chromatium vinosum (see Ref. 20, p. 46).

in the chain (1.4-1.6 nm) is similar to the separation of the iron centers found for the optimum docking of two Cyt c molecules (1.77 nm, see subsection VB). Assuming that the $\lambda/4$ and β parameters for electron transfer in the Cyt c chain are similar to those used for the Cyt c exchange in solution (25 kJ·mol⁻¹ and 12 nm⁻¹, respectively) and using the correction in subsection IVB to obtain d, k is calculated to be about 10^6-10^7 s⁻¹ for electron transfer between two adjacent Cyt c-553 or Cyt c-558 molecules and about 107-108 $\rm s^{-1}$ for downhill electron transfer from Cyt c-553 to a neighboring Cyt c-558 molecule. These estimates do not allow for the effect of the low dielectric constant of the medium, which will be to lower λ and increase k, and departure of the heme groups from co-planarity, which will tend to decrease $\kappa \nu$ and decrease k. These effects tend to cancel. It is interesting to note that the estimated rate constants for intrachain electron transfer are comparable to the rate constants for electron transfer from the terminal Cyt c-558 to the oxidized BChl-b dimer, a reaction discussed next.

The rate constant for reduction of the oxidized special pair by ferrocytochrome c-558 is $4 \cdot 10^6$ s⁻¹ at room temperature [224]. The reduction of the special pair is presumably by the closest Cyt c molecule. If the approx. 2.1 nm separation between the magnesium of the (BChl)₂ and the closest heme-iron in the crystal is also appropriate to the rate measurements, then the edge-to-edge separation of the chlorophyll and heme rings in the rate measurements is approx. 1.1 nm and, using the correction in subsection IVB, d is approx. 0.8 nm. The driving force $(-\Delta G^{0'})$ associated with the rate constant of $4 \cdot 10^6$ s⁻¹ is 170 mV or 16.3 kJ·mol⁻¹ and the reaction is expected to have only a low ΔG_r^* in Eqn. 31. The value of λ for the reaction is presently not known though a rough estimate of about 12 kJ·mol⁻¹ (based on a barrierless result) is made for (BChl)2, BChl and BPh in Appendix II. Using the value for $\lambda/4$ of 25 $kJ \cdot mol^{-1}$ for the Cyt c-Cyt c exchange reaction given earlier the estimated λ for the (BChl)⁺₂-Cyt c reaction is the averaged of these, namely about 56.5 kJ·mol⁻¹. Using the above driving force, the ΔG_r^* in Eqn. 31 is estimated to be 7.1 kJ·mol⁻¹ §. From Eqns. 31-33a and 37b the predicted rate constant is $1 \cdot 10^{13} e^{-12 \cdot 0.8} e^{-1.6/RT}$ or $4 \cdot 10^7 s^{-1}$,

using $\beta = 12 \text{ nm}^{-1}$ and d = 0.8 nm. This value may be compared with the experimental value [224] of $4 \cdot 10^6 \text{ s}^{-1}$ found for reaction 48 in *Rps. viridis*.

Picosecond flash-photolysis techniques have been used to study the primary electron-transfer reactions in reaction centers and in a variety of model photosynthetic systems [226–229]. An important result of the studies with model systems is that the forward electron transfer from an excited porphyrin monomer or dimer to an electron acceptor (usually a quinone) can be very rapid while the back reaction to reform ground-state reactants, which is even more exothermic, can be relatively slow, depending on the system and solvent used. The forward reactions generally lie in the normal free-energy region, and it has been proposed that the back reactions are slow because they lie in the inverted region.

In reaction centers of Rps. sphaeroides electron donation from the excited singlet state of (BChl)₂ to BPh occurs in 4 ps (Kirmaier, C., Holten, D. and Parson, W.W., personal communication) * and the driving force for the reaction has been reported to be 0.13 eV though perhaps it may be somewhat larger ¶. Some spectroscopic evidence (a 'shoulder') has been offered that the reaction involves the formation of BChl as a distinct

The resulting activation energy is 7.1 kJ·mol⁻¹, if ΔS^0 can be neglected. Its experimental value has not yet been determined apparently, but the experimental value for *Rps. gelatinosa* is 9.6 kJ·mol⁻¹ [225].

^{*} Current work (Kirmaier, C., Holten, D. and Parson, W.W., personal communication) on this rate constant is consistent with the earlier value [230]. Very recently, evidence for possible formation of BChl⁻ in 1 ps and BPh⁻ in 5 ps has been reported [231].

The driving force is based on estimated potentials [228,234,235] for reduction of (BChl)₂⁺ to excited (BChl)₂ and of BPh to BPh⁻. Recently, delayed fluorescence measurements gave a driving force of 0.16 eV [236]. In *Rps. viridis* the estimated potentials [223,228,237] lead to a driving force of 0.12 eV. However, the in vivo measurements for BPh/BPh⁻ in each case appear to be made with the Q present. If it were present as Q⁻ rather than as QH, the Q⁻ would destabilize the BPh⁻ and so make the calculated driving force too small. The dielectric constant D_s of the membrane has been reported to be about 3.8 [238] and has been estimated to be 30% less (Dutton, P.L., personal communication). Two negative charges a distance 1.5 nm apart in a medium of D_s = 3 have a repulsive interaction of 0.33 eV.

for the BChl to BPh rings (from the Mg in the BChl to the acetyl group in the BPh). Since the reaction is barrierless (negligible activation energy [232]) we shall take $-\Delta G^{0'} \cong \lambda$ (Appendix II) §. Then, using $\beta = 12$ nm⁻¹ and d = 0.1 nm (after subtracting 0.3 nm from the 0.4 nm distance as in subsection IVB) a half-life of the order of 0.2 ps is calculated, as in Appendix II.

The half-life for the electron transfer from BPh

to quinone in Rps. sphaeroides is 200 ps at room

temperature and 100 ps, independent of tempera-

ture, below 100 K [231]. Thus, there is a negligible

intermediate [232]. (However, such evidence does

not appear to be definitive.) From the crystal

structure the closest approach (carbon-to-carbon)

of the porphyrin rings in (BChl), and BChl ap-

pears to be about 0.35-0.4 nm, and about 0.4 nm

activation energy. The driving force for the reaction at room temperature is, incidentally, approx. 0.61 eV [228,241]. The corresponding values in Rps viridis are $4 \cdot 10^9$ s⁻¹ and approx. 0.47 eV at room temperature [224,223]. When the calculated k is written as $1 \cdot 10^{13} \exp(-\beta d)$ s⁻¹, with $\beta \approx 12$ nm⁻¹ and d estimated from the crystal structure to be in the vicinity of 1.0 nm minus the 0.3 nm correction (subsection IVB), the calculated k is about $5 \cdot 10^9$ s⁻¹, corresponding to a half-life of about 140 ps. The calculated value at 0 K (Ap-

In other experiments on the rate constant for the BPh⁻+ Q reaction, the nature of Q has been varied, causing the $-\Delta G^0$ of the reaction to change by about 0.2 eV [242]. The resulting variations in the rate constant were only a factor of two or so

[242] and, perhaps due to other changes caused by

The potentials for the reduction of (BChl)₂⁺ to excited (BChl)₂

were calculated from the energy for the ground state to singlet transition $E_{0\rightarrow0}(0\rightarrow0$ levels) and the reduction

potential for the ground state $(BChl)_{2}^{+}/(BChl)_{2}$ couple. E_{0-0}

pendix II) is only slightly greater than this.

the substitution of Q, were somewhat random. From Eqns. 1-3 it can be seen that in the vicinity of the maximum rate, i.e., when $-\Delta G^0 \approx \lambda$, a change $\delta \Delta G^{0'}$ causes the rate constant to change by a factor of $\exp[-(\delta \Delta G^{0'})^2/4\lambda RT]$. With a value of λ of about 0.5 eV, the reaction is barrierless at $-\Delta G^{0'} \approx 0.5$ eV and the above rate changes only by a factor of two when $|\delta \Delta G^{0'}| = 0.2$ eV, a result consistent with the data. Again, when a second quinone Q_B was present as Q_B^- , the rate constant for the BPh⁻+ Q_A electron transfer was smaller by almost a factor of two [243]. The presence of Q_B^- is calculated to reduce the driving force $-\Delta G^{0'}$ by perhaps several tenths of a volt, due to the coulombic repulsion of the Q_A^- and Q_B^-

in the low dielectric constant medium. Such an

effect is calculated, as above, to reduce k by a

maximum rate constant for the reduction of the

In a model porphyrin-quinone study [38] the

factor of near 2 from its value at $-\Delta G^{0'} \approx \lambda$.

quinone by the excited porphyrin (occurring at $-\Delta G^{0'} \approx \lambda$) was reported as $2 \cdot 10^{11} \text{ s}^{-1}$. The carbon-carbon distance at closest approach of the reacting partners was 0.6-0.7 nm [244], giving a d of about 0.4 nm after using the 0.3 nm correction in subsection IVB. The calculated maximum rate constant, $1 \cdot 10^{13} \exp(-\beta d)$, is about $8 \cdot 10^{10} \text{ s}^{-1}$.

which is close to the reported value of 2 · 10¹¹ s⁻¹. The various results in this section are summarized in Table IX.

VE. Reactions of cytochrome c with cytochrome b_5 and cytochrome c oxidase

In this section we consider the electron-transfer reactions of cytochrome c with cytochrome b_5 and with cytochrome c oxidase. The first reaction involves electron transfer between cytochrome c and a heme protein of opposite charge and the second involves electron transfer of cytochrome c with its natural partner.

It will be recalled that cytochrome b_5 differs from cytochrome c in several respects. The fifth and sixth coordination sites of the heme in cy-

equals the energy of the absorption maximum $h\nu_{\rm max}$ minus one-half the Stokes shift [240]. Values for the Stokes shift (450 cm⁻¹) for *Rps. sphaeroides* are given in Ref. [234]. The self-exchange rate constants k for the Chl^{+/0} and BPh^{0/-} pairs are about $2-4\cdot10^6$ M⁻¹·s⁻¹ in a somewhat polar solvent (methanol/methylene chloride) [233]. If k is written as $1\cdot10^{11}$ exp^{- $\lambda/4RT$} M⁻¹·s⁻¹, then $\lambda/4$ is estimated to be about 14.6 kJ·mol⁻¹. In the less polar membrane system, $\lambda/4$ would be less. The value of λ is estimated in Appendix II to be about 25 kJ·mol⁻¹ or less.

As pointed out in an earlier footnote, two negative charges a distance 1.5 nm apart in a medium of estimated dielectric constant of about 3 has a repulsive interaction energy of about 0.33 eV.

TABLE IX ABSOLUTE RATES FOR ELECTRON-TRANSFER REACTIONS IN PHOTOSYNTHETIC REACTION CENTERS

Reaction ^a	$\Delta G^{0 \text{ b}}$ (kJ·mol ⁻¹)	$ au_{ m obsd}$	^T calcd
Cyt c ^{II} ~~~ BChl ₂ ⁺	-16.3	0.2 μs	0.02 μs
*(BChl), ~~~ BChl ~~~ BPh	-21(?)	$\approx 1-4$ ps °	> 0.2 ps ^d
BPh ⁻ ~~~ O	- 50	100-200 ps ^e	140 ps
$BPh^- \sim Q_A, Q_B^-$ present	≈ 0.2 eV ^f	increases factor of 2	increases factor of 2
BPh ~~~ Q, Q varied	$\approx 0.2 \text{ eV}^{\text{g}}$	changes factor of 2	changes factor of 2
*Porphyrin ~~~ Q	$-\lambda$	$\tau_{\rm min} \approx 4 \text{ ps}$	$\tau_{\rm min} \approx 9 \text{ ps}$

^a Asterisks indicate electronically excited states. ^b The ΔG^0 values are approximate.

tochrome b₅ are occupied by N atoms of histidine

imidazole groups and the porphyrin ring is not

covalently bonded to the protein (other than

through the coordinated histidines). In addition,

the net charge of ferricytochrome b_5 , calculated from its amino acid composition, is -9 [245] and there are four negatively-charged carboxyl groups in the heme crevice region. By contrast, the net

charge of ferricytochrome c is +8 and the heme is surrounded by four positively charged lysine residues. Therefore, unlike the corresponding exchange reactions, interaction (of the hemes) of cytochrome c and cytochrome b_5 is attractive. By

shown that ferricytochrome c and ferrocytochrome b_5 can form a complex in which the heme groups of the two proteins are nearly coplanar with their edges separated by 0.84 nm at first

contact of the surface amino-acid side chains. Additional support for the formation of this complex has been obtained from protein modification stud-

using computer modeling, Salemme [246,247] has

ies [248,249] (although the more recent results implicate interactions between five complementary pairs, rather than just four as originally proposed) and from energy transfer measurements using

The second-order rate constant for the oxidation of ferrocytochrome b_5 by ferricytochrome c is

zinc-substituted cytochrome c [154].

phosphate ($\mu = 0.04$ and 0.20 M) (pH 7), respectively [249]. (The latter value is very similar to that reported earlier [250].) The association involves two oppositely-charged reactants, and so the equilibrium constant for this association is, as expected, very dependent on ionic strength [251]. The experimental binding constant (expressed in concentrations rather than activities) for the reactants is approx. 10^5 M^{-1} at $\mu = 0.01 \text{ M}$ and approx. $5 \cdot 10^7 \text{ M}^{-1}$ at $\mu = 0$ [251]. The extrapolation to higher ionic strengths is highly uncertain,

 $3 \cdot 10^8$ and $3 \cdot 10^7$ M⁻¹·s⁻¹ at 0.02 and 0.10 M

Like cytochrome c, the structures of oxidized and reduced cytochrome b5 are essentially identical [252]. In view of the similarity of the electron transfer distance d and the structures of the heme sites, the value of Skv is unlikely to be very

A binding constant calculated from S = 0.01 and a K_A

obtained from Eqns. 25 and 57 is only about 2-3 M⁻¹ for these oppositely charged reactants. The contrast between this

value and the extrapolated experimental binding constant

and we tentatively use a value of approx. $10^3 \, \mathrm{M}^{-1}$

for the binding constant at $\mu = 0.04$ M ¶.

^c Assignment of time constants is uncertain.

^d The value of τ_{calcd} is larger than this value if smaller λ 's are used, i.e., τ_{calcd} is about 1 ps when $\lambda = 5 \text{ kJ} \cdot \text{mol}^{-1}$ (see also Appendix

^e Value depends on temperature. This is the approximate decrease in $|\Delta G^0|$ due to the presence of Q_n^- .

⁸ This is the approximate variation in $|\Delta G^0|$ as Q is varied.

suggests that for oppositely charged reactants in contact the local electrostatic charges may dominate the interaction and require that the delocalized-charge expression Eqn. 25 be replaced by a more complex one such as that in Refs. 174-177.

different for the cytochrome c exchange reactions and the ferricytochrome c-ferrocytochrome b_5 cross-reaction. Thus, the 10^5 -fold difference in the rate constants (at $\mu=0.1$ M) for these two reactions may reflect several factors, including a difference in driving force $(-\Delta G^0=0.26 \text{ eV} [160,253])$, which contributes a factor $\exp(-\Delta G^0/2RT)$ or 150, and a difference in the stability constants of the precursor complexes (a factor of 10^3 if the experimental K_A for Cyt c-Cyt b_5 is used).

The ferricytochrome c-ferrocytochrome b_5 bimolecular rate constant and binding constant may be used (Appendix IB) to obtain a first-order rate constant for a complexed pair, namely $3 \cdot 10^8 / 10^3$ or $3 \cdot 10^5$ s⁻¹ at $\mu = 0.04$ M and pH 7, or if an S = 0.01 is also to be introduced, $3 \cdot 10^7$ s⁻¹. This value may be compared with that calculated from Eqns. 31-33 with $|\Delta G^{0'}/\lambda| \ll 1$: the rate constant calculated using (as above) $\lambda/4 \approx 25$ kJ·mol⁻¹, $\beta = 12$ nm⁻¹, $d \approx 0.54$ nm and $-\Delta G^0 = 0.30$ eV at $\mu = 0.04$ M (0.02 M phosphate buffer) [160,253] equals 10^{13} e^{-12·0.54} e⁻¹⁰ e^{5.8} or $2 \cdot 10^8$ s⁻¹, which should be compared with the value of $3 \cdot 10^5$ s⁻¹ or $3 \cdot 10^7$ s⁻¹ obtained above. There are, however, some puzzling features §.

Next we consider the reaction of cytochrome c with its physiological partner, cytochrome c oxidase. The functional unit of mammalian cytochrome oxidase, it will be recalled, contains two heme-A residues, namely cytochromes a and a_3 , differing only in their axial ligands, and two copper

atoms [256], one of which, termed 'copper A', is magnetically isolated (although the existence of a weak magnetic dipolar interaction between the Cu_A and cytochrome a has been recently proposed [257]). The Cu_A is detectable by EPR in the cupric state. The first step in the anaerobic oxidation of ferrocytochrome c by cytochrome oxidase is the very rapid oxidation of ferrocytochrome c by cytochrome a with a rate constant of $(8-9) \cdot 10^6$ $M^{-1} \cdot s^{-1}$ at pH 7.4 and 25°C; the estimates of the equilibrium constant for this reaction range from 1 to 10² [258-260]. For a binding constant of approx. 10⁵ M⁻¹ [261], the equivalent first-order rate constant for this step is $(8-9) \cdot 10^6 / 10^5$ s⁻¹; i.e., approx. 10² s⁻¹ (Appendix IB). The reaction of ferrocytochrome c with cytochrome a is followed by a slower step $(k \approx 40-100 \text{ s}^{-1})$, presumably intramolecular electron transfer from the reduced cytochrome a to copper A, for which a separation distance of 1.3-2.6 nm has been estimated [257]:

$$c^{2+} + [a]^{3+} [Cu_A]^{2+} \rightleftharpoons c^{3+} + [a]^{2+} [Cu_A]^{2+}$$
 (49)

$$[a]^{2+}[Cu_A]^{2+} \rightarrow [a]^{3+}[Cu_A]^{+}.$$
 (50)

The very rapid oxidation of ferrocytochrome c by cytochrome a is noteworthy in the light of the small driving force for the reaction (< 12 kJ·mol⁻¹ [258-260]) and the large distance between the heme groups, which has been estimated to be approx. 2.5 nm [262]. If the latter distance permits a heme-to-heme edge separation of 1.2-1.5 nm, then the above rate constants can be rationalized using the typical parameters derived earlier. For example, for modest driving forces (approx. 10.5 kJ·mol⁻¹) and small $\lambda/4$ parameters (approx. 25 kJ·mol⁻¹), a rate constant of the order of 50 s⁻¹ is consistent with an electron transfer distance d of approx. 1.5 nm if $\beta \approx 12$ nm⁻¹.

VF. Reactions of blue copper proteins

We consider next the reaction of cytochrome c with copper proteins and also other protein-protein reactions. It will be recalled that many copper-containing proteins contain a 'type 1' or 'blue' copper center which imparts an intense blue color to these substances in their copper(II) forms. Such copper centers are present in the electron

[§] Pulse radiolysis experiments on the Cyt c-Cyt b₅ complex itself show a low $(1.4 \cdot 10^3 \text{ s}^{-1})$ value for k (McLendon, G.L. and Miller, J.R., unpublished results). Also puzzling is the value of $5 \cdot 10^3$ s⁻¹ reported [154] for the k for quenching (presumably by electron transfer) of a triplet in a zinc-substituted Cyt c with ferricyt b_5 . The driving force of this reaction is about 1 eV [255], d is believed to be small (approx. 0.5-0.6 nm) and yet k is relatively small. In a plot of $\ln k$ vs ΔG^{0} for related systems the maximum k was 5.105 s⁻¹ which is also relatively low for the small value of d inferred above from computer modeling studies. Possible interpretations include a prior conformational change or a wrong docking of the two proteins. In this connection it is of interest to note that the oxidation of ferrocytochrome b₅ by metal complexes and the competitive inhibition of the reaction by redox-inactive complexes are consistent with electron transfer through the exposed heme edge and an effective charge of -4 at the oxidant binding site [245].

carriers plastocyanin, azurin and stellacyanin, which each contain one copper atom per molecule [263]. X-ray structural work has shown that the copper sites in poplar plastocyanin and in Pseudomonas aeruginosa azurin are very similar [264,265]. In each protein the blue copper is bound to two nitrogen and two sulfur atoms: the copper atom is coordinated by two histidine imidazolate groups, a cysteine thiolate, and a methionine thioether group. Although its structure has not been determined, stellacyanin must have at least one copper ligand different from that in plastocyanin and azurin, since the protein does not contain methionine. It has been suggested that in stellacyanin the site normally occupied by methionine is filled by a disulfide group [266].

After cytochrome c, plastocyanin is probably the best characterized metalloprotein, at least so far as electron-transfer properties are concerned. NMR studies have implicated two sites for the reactions of plastocyanin with metal complexes [267]. The first site is an uncharged region situated near the copper. This site may provide an electron transfer pathway via the exposed imidazole ring of His-87 and is believed to be the site used by negatively charged oxidants such as $Fe(CN)_6^{3-}$ [166]. The second site is a negatively charged region near Tyr-83 which is about 1.2 nm from the copper [267]. Electron transfer to and from this region may occur via the π -systems of aromatic amino acids and may be the site used by positively-charged oxidants such as Co(phen)₃³⁺ [164,197,268-272].

Reactions of the blue copper proteins with a variety of metal complexes have been studied [164,190,197,268–274]. Some representative examples together with the rate constants calculated from Eqn. 42 are presented in Table X. The measured self-exchange rate constants for the proteins are: azurin, $2 \cdot 10^6 \text{ M}^{-1} \cdot \text{s}^{-1}$ at 50°C [276], plastocyanin, up to $2 \cdot 10^4 \text{ M}^{-1} \cdot \text{s}^{-1}$ at 50°C [277]; stellacyanin, $1.2 \cdot 10^5 \text{ M}^{-1} \cdot \text{s}^{-1}$ at 20°C [278]. In order to calculate the k_{12} values, the azurin and plastocyanin self-exchange rates were corrected to 25°C on the assumption that the activation energies for these exchanges are comparable to that for cytochrome c. The self-exchange rates for azurin and plastocyanin at 25°C are then calculated to be $10^5 \text{ M}^{-1} \cdot \text{s}^{-1}$ and up to $10^3 \text{ M}^{-1} \cdot \text{s}^{-1}$,

respectively. We have used the upper limit for the plastocyanin exchange rate constant in calculating the k_{12} values in Table X. Comparison of the observed and calculated k_{12} values shows that plastocyanin and stellacyanin are better behaved than azurin, at least as regards their cross-reactions [272,273,279] ¶.

In the case of azurin, the calculated values of the rates of the cross-reactions with cationic metal complexes are at least comparable with those predicted from the cross-relation. However, anomalously low values for the rates are obtained when the reaction is with anionic metal complexes (Table X). This anomaly is also present, but to a lesser extent, for plastocyanin and in a milder way for cytochrome c (Table VII). Although only a limited amount of data is available, it does appear that these reactions with anionic metal complexes are anomalous. Perhaps there is specific binding of the anions. Anions tend, for example, to be more readily bound at interfaces than do cations of comparable size, a known result in electrochemistry, and there is expected to be an interplay between binding and electrostatics. There may even be multiple binding sites. Precursor complexes of ferrocyanide with azurin [280] and complexities for the cytochrome c-Fe(CN) $_6^{3-/4-}$ reaction [195,196] have been described. Different electron-transfer distances may also be involved [281].

It has proved possible to exploit binding at the Tyr-83 site of plastocyanin to obtain information about the distance dependence of electron transfer from this site to the copper center. In a recent study $Cr(phen)_3^{3+}$ bound near the Tyr-83 of reduced plastocyanin was excited using a laser flash-photolysis technique. The $Cr(phen)_3^{3+}$ excited state produced is a very powerful electron acceptor and was found to oxidize the reduced plastocyanin with a concentration-independent rate constant of $3 \cdot 10^6 \text{ s}^{-1}$. If it is assumed that the electron transfer is barrierless because of the high driving force of the reaction, then a value of

As noted earlier in connection with the cytochrome c studies, an incorrect self-exchange rate for the Ru(NH₃)₅py^{3+/2+} couple has been used in certain other cross-relation applications. This has resulted in some erroneous conclusions regarding the validity of the cross-relation.

TABLE X

COMPARISON OF OBSERVED AND CALCULATED RATE CONSTANTS FOR THE REACTIONS OF BLUE COPPER PROTEINS WITH METAL COMPLEXES

Reactions at 25°C, pH 7 and 0.5 M ionic strength unless otherwise noted. The rate constants were calculated using Eqn. 42 and the following exchange rates and redox potentials [272] for the copper proteins: azurin, $1 \cdot 10^5$ M⁻¹·s⁻¹, 0.31 V; plastocyanin, $1 \cdot 10^3$ M⁻¹·s⁻¹, 0.36 V; stellacyanin, $1 \cdot 2 \cdot 10^5$ M⁻¹·s⁻¹, 0.19 V. Net charges on the proteins (oxidized/reduced) are: azurin (-1/-2), plastocyanin (-9/-10), stellacyanin (0/0) [194,273]. The zero net charge on stellacyanin is based on the ionic-strength independence of its reaction with Co(phen)³⁺₃ [164]. For additional data, see Table VII.

Protein	Reagent	K_{12}	$\begin{array}{c} k_{12,\text{obsd}} \\ (M^{-1} \cdot s^{-1}) \end{array}$	$k_{11,\text{calcd}} $ $(\mathbf{M}^{-1} \cdot \mathbf{s}^{-1})$	Ref.
Azurin	Co(phen) ₃ ³⁺	7.0	4.4·10 ³	5.6·10 ³	194
(R. aeruginosa)	$Ru(NH_3)_5 py^{3+}$	1.5	$2.0 \cdot 10^{3} a$	1.3 · 105	190
	Fe(EDTA) ²⁻	$1.6 \cdot 10^3$	$1.3 \cdot 10^3$	$2.2 \cdot 10^6$	270
	$Co(ox)_3^{3-}$	2.5 · 10 4	$2.9 \cdot 10^{-2}$	$2.3 \cdot 10^{1}$	194
Plastocyanin	Co(phen) ₃ ³⁺	1.0	$1.0 \cdot 10^{3}$	$2.1 \cdot 10^{2}$	194
(spinach)	$Ru(NH_3)_5 py^{3+}$	0,21	$7.1 \cdot 10^{3} \text{ a}$	$4.8 \cdot 10^{3}$	190
	Fe(EDTA) ²⁻	1.1 · 10 4	8.2 · 10	$5.8 \cdot 10^{5}$	270
	$Co(ox)_3^{3-}$	$3.5\cdot10^{3}$	$2.4 \cdot 10^{-1}$	$9.1 \cdot 10^{-1}$	194
Stellacyanin	$Co(phen)_3^{3+}$	$7.5 \cdot 10^{2}$	1.8 · 10 5	6.3 · 10 4	194
(R. vernicifera)	$Ru(NH_3)_5 py^{3+}$	$1.6 \cdot 10^{2}$	1.9·10 ⁵ a	$1.5 \cdot 10^6$	190
	Fe(EDTA) ²	$1.5 \cdot 10^{1}$	4.3·10 ⁵	$2.3 \cdot 10^5$	270
	$Co(ox)_3^{3-}$	$2.6 \cdot 10^{6}$	$7.3 \cdot 10^{2}$	$2.7 \cdot 10^{2}$	194

a Ionic strength, 0.1 M.

 $\beta \approx 12 \text{ nm}^{-1}$ and an intramolecular electron-transfer distance d of 1.0 nm yields a rate of $6 \cdot 10^6 \text{ s}^{-1}$ [282].

cyanin (Cu^I and Cu^{II}) with various metal com-

plexes have been measured for reactions having

Bimolecular reaction rate constants of plasto-

very negative ΔG^0 's, and it is interesting to compare the results with those of cytochrome c. The reaction of plastocyanin Cu^I with the excited state of $Ru(bpy)_3^{2+}$, of Cu^{II} with $Cr(phen)_3^{2+}$, of Cu^{II} with the excited state of $Cr(phen)_3^{3+}$ and of Cu^{II} with excited $Ru(bpy)_3^{2+}$, for which $-\Delta G^{0'}$ is 0.48, 0.64, 1.06 and 1.20 V, respectively, have rate constants of 1.9, 2.3, 3.5 and 4.2, in units of 10^9 $M^{-1} \cdot s^{-1}$, respectively [281]. The diffusion-controlled rate constants k_D calculated (for spherically symmetric reactants) using Eqns. 25 and 53b are all about $6 \cdot 10^9$ $M^{-1} \cdot s^{-1}$. The above values

are perhaps somewhat less than this, but the dif-

ference is small, and one possible explanation is

mentioned briefly in Appendix IB. Another possibility is that the calculation of a diffusion-con-

trolled rate constant corrected for electrostatic effects is approximate.

We next consider the reactions of the blue

copper proteins with other proteins as well as some other protein-protein reactions. Observed and calculated rate constants for these systems are compared in Table XI. Except for the reaction of plastocyanin with cytochrome c, in which the reactants are of very different charge types, the electrostatic work term corrections (based on total protein charges) are small and have not been included. It can be seen from Table XI that good agreement between observed and calculated rate constants is obtained. It is of interest that the rate constants for the reactions of azurin with other proteins appear to be in better agreement with the cross-relation than are the rate constants for its

In concluding this section, it is useful to compare the self-exchange rate constant found for Cyt $c (1.2 \cdot 10^3 \text{ M}^{-1} \cdot \text{s}^{-1})$ at 25°C with that found for azurin $(2 \cdot 10^6 \text{ M}^{-1} \cdot \text{s}^{-1})$ at 50°C [276]). The

reactions with small metal complexes.

b Ionic strength, 0.2 M.

TABLE XI

COMPARISON OF OBSERVED AND CALCULATED RATE CONSTANTS FOR PROTEIN-PROTEIN REACTIONS

Unless otherwise indicated, the self-exchange rate constants and reduction potentials are from Tables VII and VIII and the calculated rate constants do not contain the W₁₂ correction. The references in the last column should be consulted for the experimental rate

Oxidant	Reductant	$\frac{k_{12,\text{obsd}}}{(\text{M}^{-1}\cdot\text{s}^{-1})}$	$\frac{k_{12,\text{calcd}}}{(M^{-1} \cdot s^{-1})}$	Ref.	
Cyt c	Azurin	$1.6 \cdot 10^3$	3.104	284	-
Plastocyanin	Cyt c	$1.0 \cdot 10^6$	5·10 ^{5 a}	284	
Stellacyanin	Cyt c	$3.5 \cdot 10^{2}$	$4 \cdot 10^3$	283	
Azurin	Cyt <i>c</i> -551	$6.0 \cdot 10^6$	8·10 ^{5 b}	284	
Plastocyanin	Cyt <i>c</i> -551	$7.5 \cdot 10^{5}$	5·10 ^{5 b}	284	
Cyt c	Cyt <i>c</i> -551	$6.7 \cdot 10^{4}$	1·10 ^{5 b}	284	

^a The W_{12} correction is considerably larger for this reaction than for the other systems considered, and was included in the calculation

self-exchange [193] yields a k of about $2 \cdot 10^4$ M⁻¹·s⁻¹ at 50°C. Oxidized and reduced azurin carry small net negative charges (Table X) and hence have negligible coulombic repulsion, while ferro- and ferricytochrome c have substantial positive charges (Table VII). This difference in coulombic repulsions contributes a factor of about 5 to the ratio of the rate constants. Again, the values of the electron-transfer distance d differs for the two reactions. In the crystal of azurin from

activation energy of 54 kJ·mol⁻¹ for the Cyt c

VG. Cytochrome c peroxidase

the reorganization term $(\lambda/4)$ needed for electron

We conclude this review with an example which illustrates the kinetic complexity which can arise in an oxidation-reduction scheme and for which an effort was made to extract a rate constant for an electron-transfer step. We use as an example the cytochrome c peroxidase system, which catalyzes

transfer [286,266].

Alcaligenes denitrificans pairs of azurin molecules are related by an approximate 2-fold axis, with their hydrophobic patches in contact and with a distance of 0.7 nm between the edges of the two


the oxidation of ferrocytochrome c by hydrogen peroxide and alkyl hydroperoxides. The reaction of yeast cytochrome c peroxidase (CcP) with cytochrome c [287–289] and with metal complexes involves several intermediates [290-292]. Scheme I summarizes some [291,293] (but not all [293]) of

the features that have been proposed for the reaction of cytochrome c with metal complexes. The first step is the formation of a CcP-peroxide complex CcP-OOH which then undergoes a bimolecular reaction with the added reductant to form RFe^{IV} or an intramolecular proton-catalyzed redox reaction to form CcPI. The latter is described as an Fe^{IV} (ferryl) species possessing an oxidized

histidine imidazole rings [285]. This distance is less than the 0.94 nm estimated earlier for the edge-toedge distance of the two histidines in the Cyt c case. This difference is estimated to contribute a factor of exp $(12 \cdot 0.24)$ or about 20 to the value of k. These three factors (temperature, coulombic repulsion and distance) account for the difference in self-exchange rates of Cyt c and of azurin, if the values of $\lambda/4$ for both are comparable. Interestingly, the geometry of the ligands in azurin is that of a distorted tetrahedron, intermediate between the tetrahedron common for 4-coordinate Cu^I complexes and the square planar or tetrahedron common for 4-coordinate CuII, thereby reducing

protein residue (methionine-172 or tryptophan-51) [292-294] ¶. There is some disagreement concerning the formulation of the subsequent steps indicated by the solid [291] vs. dashed [293] lines in

b Calculated using 1.2·10⁷ M⁻¹·s⁻¹ for the self-exchange rate constant [206] and +0.28 V for the reduction potential of the cytochrome c-551 couple [275]. The net charge on the protein is -2/-3 [284].



Scheme I.

the scheme, where $(RFe^{IV})_u$ denotes a species which is not in its most stable conformation.

The kinetics of the reactions of yeast cytochrome c peroxidase with metal complexes has, so far, only been interpreted in terms of the mechanism indicated by the solid lines [291]. In this interpretation the rate law permits the assignment of a rate constant for the reduction of R+Fe^{IV} to R+FeIII and/or for the reduction of RFeIV to RFe^{III} (the kinetic data do not allow a unique assignment). Both reactions involve the reduction of the Fe^{IV} peroxidase to the Fe^{III} state and can be represented by $Fe^{IV}O + e^{-} = Fe^{III}O$. On the basis of the assumption that the stable form of the Fe^{III} product is Fe^{III}OH (i.e., that the Fe^{III} species contains a coordinated hydroxide ion), the E^0 for Eqn. 50 has been estimated to be +0.7 V [291]. Application of the cross-relation without any work corrections then yields a rate constant of approx. $10^2 \text{ M}^{-1} \cdot \text{s}^{-1}$ for the exchange between the Fe^{IV} and Fe^{III} peroxidase. This value is higher by a factor of 1 · 106 than another estimate [290] in which proton-coupled changes were not considered \S . As discussed above, electron transfer through the exposed heme edge is implicated for the cytochrome c system, and it is likely that a similar mechanism obtains for the peroxidase exchange as well [291].

The bimolecular rate constant for the reaction of ferrocytochrome c with the peroxidase is approx. $5 \cdot 10^8 \text{ M}^{-1} \cdot \text{s}^{-1}$ at pH 6.0 [287]. This very rapid rate, which is 10^3-10^4 -times more rapid than the reaction of the peroxidase with metal complexes of comparable redox potential and exchange rates, has been ascribed [291] to the stability of the precursor complex formed between cytochrome c and the (negatively charged) peroxidase (binding constant, approx. 10^5 M^{-1} [45]). The X-ray structure of yeast cytochrome c peroxidase has been determined [296] and computermodeling [297,298] of the complex formed between cytochrome c and the peroxidase implicates a structure in which the two heme groups are almost parallel with a perpendicular separation between their planes of 0.68 nm and a heme edge-to-edge separation of 1.65 nm. Although coupling of the two heme groups via aromatic (specifically histidine-181 [299]) and conjugated groups has been invoked as an additional rate-enhancing factor [297,298], no 'special' electronic interaction of the reactants may be necessary to account for a first-order rate constant of approx. 10^3-10^4 s⁻¹. Thus ΔG^* is small ¶ and, for $\beta = 12 \text{ nm}^{-1}$ and d = 1.35 nm (after using the 0.3 nm correction in subsection IVB) the value of $1 \cdot 10^{13} \exp(-\beta d)$ equals $1 \cdot 10^6$ s⁻¹.

It has recently been proposed (Ref. 295 and references cited therein) that the formation of CcPI from the resting state of the enzyme CcP (PFe^{III}R) proceeds via an iron(IV) porphyrin cation-radical intermediate (P⁺ Fe^{IV}R) and that the imidazole coordinated to the iron center then mediates the oxidation of the nearby methionine-172 by the porphyrin radical to form CcPI (PFe^{IV}R⁺).

[§]The estimate of the self-exchange k in Ref. 290 is based on a fitted redox potential $E^{0}'=1.1$ V for the CcPII/CcP couple at pH = 5.3 and $\mu=0.1$ M, where CcPII is the one-electron reduction product of CcPI: E^{0}' was deduced from applications of the cross-relation and based on deviations of the f factor there from unity, a very uncertain procedure. The E^{0}' in [291] corrects the estimated redox potential for assumed pH effects and suffers from a similar uncertainty. Different assumptions about the nature of the proton transfers that are coupled to the electron transfer will lead to a different exchange rate.

[¶] The ΔG^{0} , for the reaction has been estimated to be very favorable [290,291] and $\lambda/4$ for a heme-heme system is expected to be small.

VI. Conclusions

The theory of electron-transfer reactions has been particularly useful for correlating a wide body of experimental data on reactions of metal complexes in solution, particularly via the cross-relation. It has also been useful in understanding the absolute values of the reaction rates, and indeed

gives reasonable agreement with the experimental data (Table I). It has been used for predictions,

including that of the recently observed inverted effect. One large area of study, only briefly mentioned in this review, but which has played a significant role in the interplay of theory and experiment, has been that of electron transfers at electrodes. Here theory has been useful in predicting correlations between homogeneous and electrochemical electron transfers.

In the application to biological systems new

features arise. Frequently, the reactants may be fixed some distance apart, and the attenuation of the coupling of their electronic orbitals with separation distance then becomes an additional significant factor; not infrequently the (edge-to-edge) separation distance of the redox centers is only approximately known. This additional factor is more readily studied when the reaction is in the 'barrierless' regime, i.e., in the regime where the driving force $-\Delta G^{0'}$ has overcome the intrinsic

reorganizational barrier λ .

There are several additional features which arise in the application of the theory to systems of biological interest for which increased information would be very useful. They include distances and orientations of reactants, and the reorganizational response of the protein to changes in charges of the redox centers. Nevertheless, the correlation between predicted and calculated rates for reactions involving redox proteins can be very good, as illustrated for the reactions of cytochrome c with various inorganic reagents (Table VII). In the case

of intramolecular electron transfers (subsection

VC), the current results on biological systems are consistent with an electron-transfer rate that falls

off exponentially with distance, with an exponent

near those found for nonbiological systems (Table

IV). In applying the equations to the biological systems we have, for concreteness, estimated some of the numerical values of the rate-determining

parameters even when they were not known independently. These estimates, made to illustrate the magnitudes of the parameters, are of course to be regarded as tentative at this time. Nevertheless, it is encouraging that the values used lead to a consistent treatment of the reactions of the metalloproteins (mainly cytochrome c) with a variety of substrates and of reactions in the photosynthetic system.

Appendix I. Expressions for activation- and diffu-

IA. No steric effects

sion-controlled reaction rates

The physical events involved in a bimolecular collision in solution between two molecules A and B involve a random diffusion of the molecules toward and away from each other, becoming in some instances a close-contact encounter complex $A \cdot B$ and then diffusing apart. We may write this process as

$$A + B \underset{k_{-D}}{\rightleftharpoons} A \cdot \cdot B \tag{51}$$

The electron transfer can occur at any stage of this process, with shorter distances weighted more favorably because of the nonadiabatic $\exp(-\beta r)$ factor. In the absence of reaction, the rate constant for diffusion k_D multiplied by the average number of repeated encounters y will equal the mean collision frequency Z of the pair in solution. Since $k_{\rm D}$ for small uncharged species with no specific

systems. If during the encounters, electron transfer occurs with a first-order rate constant k_1 , and if the reverse electron transfer rate is negligible relative to the rate of separation of A⁺ and B⁻,

interactions is of the order of 1/10 to 1/100 the

value of Z,γ is of the order of 10 to 100 for such

 $A \cdot \cdot B \stackrel{k_1}{\rightleftharpoons} A^+ B^- \rightarrow A^+ + B^-,$ (52)

the overall rate constant for electron transfer is given by Eqn. 53 in a steady state approximation

for the concentration of $A \cdot \cdot B$ [300,301], for the case of spherical reactants:

$$\frac{1}{k_{\text{obsd}}} = \frac{1}{k_{\text{D}}} + \frac{1}{K_{\text{A}}k_{1}},$$
 (53a)

where K_A , which equals k_D/k_{-D} , is the equilibrium constant for formation of $A \cdot B$ in Eqn. 51. The force field used to calculate K_A and also k_D (both depend on long-range electrostatic forces) is taken to be spherically symmetric. The value of k_D is given by [302]:

$$k_{\rm D} = 4\pi D / \int_{r=g_{12}}^{\infty} \exp(w^{\rm r}/RT) r^{-2} \, \mathrm{d}r$$
 (53b)

where D is the sum of the diffusion constants of

the two reactants and the work term w^r is a function of r and, in the presence of an ion atmosphere, is assumed to be given by its equilibrium value, e.g., by Eqn. 25 in a Debye-Hückel approximation. (If the ionic atmosphere were not quite keeping pace with these diffusing reactants, a different effective w^r would be needed.)

The rate constant k appearing in Eqn. 1 is the same as the $K_A k_1$ in Eqn. 53a. In other words, when the reaction is partially diffusion-controlled, Eqn. 53a is used with $K_A k_1$ given by Eqn. 1, and hence by

$$K_{A}k_{1} = \kappa A\sigma^{2}\exp(-\Delta G^{*}/RT). \tag{54}$$

Several models for $A\sigma^2$ are available. For example, since $K_A k_1$ is equal to the rate constant when diffusion control is absent, one sees that it is equivalent to some mean (equilibrium) collision frequency in solution multiplied by the reaction efficiency. The effective collision frequency in solution is the value $Z_{\rm soln}$ in the absence of the work term w^r , multiplied by an exponential factor $\exp(-w^r/RT)$, which influences the probability of close encounters. The reaction efficiency is $\kappa \exp(-\Delta G_r^*/RT)$, where ΔG_r^* is the free energy barrier to reaction between fixed sites at the separation distance r and is given by Eqn. 32. Thereby:

$$\kappa A \sigma^2 \exp(-\Delta G^*/RT) = \kappa Z_{\text{soln}}$$

$$\times \exp\left[-\left(w^{r}+\Delta G_{r}^{*}\right)/RT\right].$$
 (55)

If Z_{soln} has its gas phase value $(8\pi RT/m)^{1/2}\sigma^2$, where m is the reduced mass of the collision partners $m_{\text{A}}m_{\text{B}}/(m_{\text{A}}+m_{\text{B}})$ and σ is the separation distance r in the encounter complex $A \cdot \cdot B$, then, since ΔG^* equals $(w^{\text{r}} + \Delta G^*_r)$,

$$A = (8\pi RT/m)^{1/2} \tag{56}$$

Another model [18,21,303] is to take the equilibrium constant K_A as the effective volume occupied by the separation coordinate over which reaction occurs, $4\pi r^2 \delta r$ (the r's over which electron transfer can still occur significantly lie between r and $r + \delta r$) , multiplied by the additional free energy factor $\exp(-w^r/RT)$ describing the additional probability of finding the pair in the reaction zone:

$$K_{\rm A} = 4\pi\sigma^2\delta\sigma \exp(-w^{\rm r}/RT), \tag{57}$$

where r has been set equal to σ .

The first-order rate constant k_1 in reaction 52 is taken to be an effective frequency ν for motion along a 'reaction coordinate' (vibrational mainly, but also with solvational contributions) multiplied by the reaction efficiency, mentioned earlier $\kappa \exp(-\Delta G_r^*/RT)$,

$$k_1 = \kappa \nu \, \exp(-\Delta G_r^*/RT). \tag{58}$$

We then have from Eqns. 54, 57 and 58 and the relation between ΔG^* and ΔG_r^* that

$$A = 4\pi\nu\delta\sigma\tag{59}$$

Eqns. 56 and 59 yield a value for A within a factor of 10 or so of each other, depending, of course, on the size of $\delta\sigma$ which is typically of the order of $1/\beta$ or 0.08 nm [21,304]. This is the value used in the calculations in the present paper.

In the case of an adiabatic reaction κ is unity and the frequency ν in Eqn. 58 is given as a

This δr also appears in the theory described in Refs. 3 and 58, via a term ρ there which is estimated to be of the order of unity and hence omitted from the present Eqn. 1; ρ is the ratio of this δr to the width of the 'equivalent equilibrium distribution' along the reaction coordinate (defined in Refs. 2 and 3).

weighted sum of nuclear frequencies ν_i , namely:

$$\kappa \nu = \nu = \left(\frac{\sum_{j} \lambda_{j} \nu_{j}^{2}}{\sum_{j} \lambda_{j}}\right)^{1/2} \qquad \text{(adiabatic case)} \qquad (60)$$
when there are several vibrational motions con-

tributing to ν and to λ ($\lambda = \sum_{i} \lambda_{i}$) [305]. In the case of a nonadiabatic reaction ($\kappa \ll 1$) the product $\kappa \nu$ is given instead by Eqn. 32, and is seen to be independent of the frequencies ν_i of the nuclear motion!

case of a nonadiabatic reaction (
$$\kappa \ll 1$$
) the product $\kappa \nu$ is given instead by Eqn. 32, and is seen to be independent of the frequencies ν_j of the nuclear motion!

An estimate of the value of $\kappa \nu$ at contact of the reactants can be made from values of H_{AB} in the literature. The estimate of H_{AB} in Ref. 63 for edge-to-edge contact of the hemes and $\lambda \approx 1$ V yields $\kappa \nu \approx 3 \cdot 10^{12}$ s⁻¹, which is close to the value of 10^{13} s⁻¹ assumed in Section V. An estimate of H_{AB} in Ref. 306 for a face-to-face contact of hemes gives a value for $\kappa \nu$ even larger than $1 \cdot 10^{13}$

adiabatic reaction, and the latter equals a typical vibration frequency approx. •1013 s-1, one concludes that $\kappa \approx 1$ for the latter contact. We next give the equation for ionic strength effects obtained from Eqns. 1-3 by including the effects on the work terms (Eqn. 25) and on the activity coefficients affecting the $\Delta G^{0'}$ in the prevailing medium. Using the expression given in the footnote to Eqn. 38 for small $|\Delta G^{0'}/\lambda|$, and using the Debye-Hückel expression for the activity coefficients, we obtain Eqn. 61 for the case of reaction 7. In this reaction z_1 increases by unity and z_2

 s^{-1} . Since $\kappa \nu$ shouldn't exceed the value for an

$$+ \frac{Ne^2B\sqrt{\mu}}{D_sRT} \left(\frac{\left(z_1 + \frac{1}{2}\right)}{1 + B\sigma_1\sqrt{\mu}} - \frac{\left(z_2 - \frac{1}{2}\right)}{1 + B\sigma_2\sqrt{\mu}} \right)$$
(61)
where C is the factor which multiplies $-z_1z_2$ in

latter one notes that in Eqn. 25 $\sigma_{12} = \frac{1}{2}(\sigma_1 + \sigma_2)$.)

decreases by unity in a one-electron transfer:

 $\ln k = \ln \bar{k} + \left[\left(z_1 + \frac{1}{2} \right) \left(z_2 - \frac{1}{2} \right) - \frac{1}{4} \right] C$

IB. Steric effects included

A steric factor S, not present in the above equations, was introduced in subsection VB to take into account the fact that only some limited range of orientations of A and B at a given r may be appropriate for electron transfer in protein systems. We denote by A - B and AB pair in such a properly oriented configuration; A - B is formed from the encounter complex A · B by suitable reorientation of A and/or B:

$$A \cdot \cdot B \rightleftharpoons A - B \rightarrow A^{+} + B^{-} \tag{62}$$

For the present review it is convenient to separate approximately two contributions to the equilibrium constant (denoted by K_e) for the formation of A - B from A and B. This constant K_e differs from that for forming $A \cdot \cdot B$ from A and B in two respects. (1) The work term appearing in K_{Δ} (Eqn. 57) is now for a fixed orientation of A

and B and is based on the actual charge distribu-

tion in the transition state rather than on total

charges; (2) There is a steric factor S to correct for

the fact that the 4π in K_A assumes spherical symmetry, i.e., it is obtained by integrating uniformly over the angular coordinates. This factoring of the equilibrium constant K_e into S and K_A is, as already noted, approximate \(^{\mathbb{I}}\). A first-order rate constant k_1 for a properly docked pair (active sites in contact) can be defined

as in Eqn. 58, where ΔG_r^* is given by Eqn. 32. However, now the w^r and w^p appearing in ΔG^{0} there refer to work terms for forming A - B and $A^+ - B^-$ in the properly docked configuration (the same for both). The (activation-controlled) bi-

 \P While the factoring of K_e can be avoided by using statistical

mechanics, the resulting expression would be more com-

cal localization of A and B into a small range of r's, the localization of their relative angular coordinates into an interval, and the electrostatic and other nonelectrostatic contributions. In contrast, the w^{r} in Eqn. 57 is only one of these.

The quantity W^{r} is a work term which includes the geometri-

Eqn. 38. In a self-exchange reaction $z_1 + 1$ equals z_2 , and σ_1 equals σ_2 , so that the last term in Eqn. 61 then vanishes. For any reaction Eqn. 61 reduces to Eqn. 41 at sufficiently low μ . (To obtain the

plicated. The K_e that is used above (SK_A) actually has three contributions to the formation of A-B from A and B, a geometrical one in K_A (the $4\pi\sigma^2\delta\sigma$), a nongeometrical one $[\exp(-w^{r}/RT)]$ and a reorienting-into-a-docking one S. In the more rigorous statistical mechanical expression all contribute to a single free-energy expression W^r , $K_e =$ $\exp(-W^{r}/RT)$, though in a complicated, entangled way.

molecular rate constant can be written as SK_Ak_1 with K_A given by Eqn. 57, where w^r is now for the docked configuration. Hence,

$$k = SK_{A}\kappa\nu \exp(-\Delta G_{r}^{*}/RT). \tag{63}$$

Eqns. 43 and 44 immediately follow from Eqns. 63 and 32. Further, the maximum value of the activation-controlled k is obtained by setting ΔG_r^* equal to zero in Eqn. 63, and yields the value given in subsection VC, $SK_{\Delta}\kappa\nu$.

An experimental value for k_1 can be obtained from the bimolecular rate constant k and the equilibrium constant K_e (measured kinetically or spectroscopically) for formation of A - B from A and B, if the 1:1 complex principally exists as A - B rather than $A \cdot B$, namely $k_1 = k/K_e$. If K_e is not known, its calculated value SK_A may be used instead, and we have $k_1 = k/SK_A$. Both results are used in subsection VC.

Since the work terms in Eqn. 63, namely, the w^{r} and w^p in K_A and ΔG_r^* , are appropriate to the docked configuration, they require a detailed knowledge of the charge distribution, particularly in the vicinity of the active sites of the proteins. Simpler calculations, discussed in subsection IVD, are typically adopted and one of them is used here, for convenience, in the form of Eqn. 25. It should be noted, however, that the binding constants for natural partners, which generally have opposite charges, tend to be considerably higher than the values calculated from Eqns. 25 and 57, presumably because of the breakdown of the Debye-Hückel type theory for such cases, a result anticipated by Bronsted in his theory of specific ionic interaction [176] (see subsection VE). In the cases discussed in this article, the actual binding constants were used for these cases.

When the rate constant k given by Eqn. 63 becomes sufficiently high the reaction may, as in the spherically symmetric case (Appendix IA), become diffusion-controlled. Whereas the problem in the latter case involved the solution of a differential equation for diffusion and reaction, leading in the steady state to Eqn. 53, the differential equation to be solved now also involves a rotational-diffusive process in the reactants. For simplicity, we shall continue to assume Eqn. 53, with $K_A k_1$ now given by the k in Eqn. 63. If rotational

diffusion were sufficiently slow, however, the maximum rate might now be below k_D , even if $k_D \gg K_A k_1$. However, the rate constants for the plastocyanin reactions show that for these systems the rate constant is close to the translational diffusion-controlled rate constant k_D .

We consider, for completeness, rate saturation effects which sometimes occur. We examine first reactions 51 and 52. In Eqn. 53 it was tacitly assumed that $(A \cdot B) \ll (A)$. Under these conditions a steady-state assumption for $(A \cdot B)$ is appropriate. More generally the reaction is biphasic in time with a fast time constant being associated with the establishment of a near steady-state for $(A \cdot B)$ and with the slow time constant being given by

$$\frac{1}{k_{\text{obsd}}} = \frac{1}{K_{\text{A}}k_{1}(\text{B})} + \frac{1}{k_{\text{D}}(\text{B})} + \frac{1}{k_{1}},\tag{64}$$

where $k_{\rm obsd}$ is measured in s⁻¹. When (B) becomes very small the $1/k_1$ can be neglected and Eqn. 64 reduces to Eqn. 53. On the other hand, when (B) becomes large, $k_{\rm obsd}$, measured in s⁻¹, becomes k_1 and, thereby, independent of (B). Rate saturation has occurred.

Rate saturation effects may also occur in a reaction mechanism involving the formation of a dead-end complex. The slow step can now even be the dissociation of the dead-end complex rather than the electron transfer. Some discussion of these effects is given in Refs. 197 and 281, and is omitted here in the interest of brevity.

IC. Dependence of H_{AB} on nuclear configuration

In the treatment in Section II, it was assumed that the thermal electron-transfer matrix element H_{AB} was independent of nuclear configuration (the Condon approximation). Actually, there may be some dependence, e.g., Ref. 307. The detailed analysis of this effect depends, in part, on whether the nuclear motion is largely classical (minor nuclear tunneling) or highly quantum mechanical (extensive nuclear tunneling).

For the present purpose a very simple treatment is adopted in which the nuclear motion is treated classically, so that the electron transfer occurs at the intersection of the curves in Fig. 1[¶]. It can then be shown [§] that for a system at the intersection the potential energy of the reductant minus its potential energy after loss of an electron, equals $(-\lambda_1 \Delta G^{0'}/\lambda)$, where λ_1 is the contribution of the reductant (reactant 1) to λ ($\lambda = \lambda_1 + \lambda_2$), and that the corresponding value is λ_1 when the reactant is in its equilibrium configuration. Thus, if for reactant 1 in its equilibrium configuration of an electron to an adjacent bridging unit is V_0^c , the corresponding value V_0 , when the system is at the intersection is

$$V_0 = V_0^e - \frac{\lambda_1 \Delta G^{0'}}{\lambda} - \lambda_1. \tag{65}$$

Hence, also, V_0 equals V_0^e when $-\Delta G^{0\prime} = \lambda$. If $\Delta G^{0\prime}$ is varied at fixed λ_1 and λ_2 by varying only the oxidant, V_0 changes only because the energy of the intersection in Fig. 1 changes. The magnitude

of the thermal matrix element $|H_{AB}|$ is given by

$$|H_{AB}| = B \left(\frac{B}{V_0}\right)^N, \tag{66}$$

where V_0 is given by Eqn. 65, N is the number of intervening sites (identical bridging units) between the two reactants and B is the exchange integral between adjacent sites [307,309-312] . (B was taken here for simplicity to be the same for all adjacent pairs.) Thereby, the β in Eqn. 37 is given by:

$$\beta = 2l^{-1} \ln \frac{V_0}{R} \tag{67}$$

where l is the length of a bridging unit.

If, instead of varying the oxidant, the reductant were varied at fixed λ_1 , λ_2 and $\Delta S^{0'}$, and if V_0^e again denotes the value of V_0 for the case where $\Delta G^{0'} = -\lambda$, the value of V_0 at any other $\Delta G^{0'}$ would now be given by Eqn. 68. A change in $\Delta G^{0'}$ now changes V_0 not only by the nuclear configuration term but also by changing the electronic energy level of the reductant itself.

$$V_0 = V_0^{e} - \frac{\lambda_1 \Delta G^{0'}}{\lambda} - \lambda_1 - (\Delta G^{0'} + \lambda)$$
 (68)

We next apply Eqns. 65 and 67 to experiments in which $-\Delta G^{0'}$ is varied from 0 to λ in the normal region and from λ to 2λ in the abnormal region by varying the oxidant (as was the case in the experiments in Ref. 36.) For concreteness, we treat a solvent-mediated electron transfer and take $2\lambda_1 = \lambda = 1$ eV and $V_0^c = 5$ eV, the value for a vertical low energy ion-to-solvent charge-transfer transition. The change in β when $|\Delta G^{0'}|$ is changed by λ is estimated from Eqn. 67 to be about 0.4 nm⁻¹. If the separation distance d is 1.0 nm, then βd changes by about 0.4 when $|\Delta G^{0'}|$ changes by λ , while the classical free-energy barrier in the

In a nuclear tunneling situation, a broader interval of coordinates is involved for tunneling from any initial quantum state of the reactants.

[§] The argument used is based, for the present purpose, on a harmonic oscillator analysis and, to simplify the discussion, allots a solvent contribution outside the coordination shell to each ion, so that $\lambda = \lambda_1 + \lambda_2$. (A more detailed treatment of the solvent would not pose any difficulty and would be based on arguments related to those used in Ref. 12.) The potential energy in coordinate q of the reactant, specifically the reductant, is $f(q-q^r)^2/2$, where q^r is the equilibrium value of q and f is a force constant, assumed here to be the same in the ionized and ground states. If the electron is suddenly lost, this potential energy of this reactant is now $f(q-q^p)^2/2$, where q^p is the new equilibrium value of q after the electron is lost. Setting $q = q^r$, the vertical ionization energy V_0 for an ion in its equilibrium nuclear configuration is seen to contain a term $f(q^r - q^p)^2/2$, making it less than it would otherwise be. Summed over all q's of the reductant this term $\sum f(q^r$ $q^{\rm p}$)²/2 is denoted by λ_1 . We next make a similar analysis when the reductant is in a configuration at the intersection in Fig. 1. A treatment in Ref. 308 is used. In particular, at the intersection we note that a coordinate q equals q^{*} , namely $(m+1)q^r - mq^p$ [308]. The contribution to the vertical ionization potential is now $\sum f(q^{\neq} - q^{p})^{2}/2 - \sum f(q^{\neq}$ $(2m+1)^2/2$, which equals $(2m+1)\sum f(q^r-q^p)^2/2$, i.e., $(2m+1)\sum f(q^r-q^p)^2/2$ 1) λ_1 . But 2m + 1 equals $-\Delta G^{0'}/\lambda$ [308], and so one obtains the result cited in the text.

The superexchange mechanism via an atom was considered earlier in Ref. 313. The expression for the typical case that there are several orbitals in a bridge involves a multiple sum over these orbitals for each bridge, and each B/V_0 depends on the orbital. See, for example, the footnote in subsection IVA and Eqn. 25 of Ref. 73, and the extension of the latter to more than two bridges [73].

expression $(\lambda/4RT)$ $(1 + \Delta G^{0'}/\lambda)^2$ at room temperature equals 9.6, 0 and 9.6 when $-\Delta G^{0'} = 0$, λ and 2 λ , respectively. The effect of variation of β in the normal region or in the abnormal region, while not negligible, is seen to be minor in comparison with the variation in $(\lambda/4RT)$ $(1 + \Delta G^{0'}/\lambda)^2$. To the extent that nuclear tunneling is important, the effective sampling of extremes of potential energy and the thermal barrier would both be less.

ID. Miscellaneous

Eqn. 53, or its equivalent, was originally derived by solving the diffusion equation subject to the boundary condition that reaction occurs with a certain efficiency at the boundary $r = \sigma$. Recently, this diffusion equation, or really a reaction-diffusion equation, was solved numerically for the case that reaction can occur at any r with a reaction probability which varies as $\exp(-\beta r)$ [55]. The results showed that within this 'continuum' framework for diffusion, Eqn. 53 provides a good approximation for the conditions studied, e.g., as in Table V of Ref. 55.

Not considered in the present review is the effect of the dynamics of solvent dielectric relaxation on electron-transfer rates, experimental data on this subject being almost absent. A recent experimental study on an intramolecular electron transfer in the barrierless regime has shown agreement between the rate constant of the electron transfer and the reciprocal of a ('constant-charge' corrected) longest measured dielectric relaxation time [314]. Several recent theoretical studies on solvent dynamics in charge-transfer reactions include references (Refs. 315 and 316, and references cited therein; also Sumi, H. and Marcus, R.A., unpublished data).

We conclude this section with some comments on the slope α of plots of $-RT \ln k$ vs $\Delta G^{0'}$ at $\Delta G^{0'} = 0$, or, in the electrochemical case, versus the activation overpotential η at $\eta = 0$. It follows

from the analysis given in Eqns. A-14, 67 and A-11 of Ref. 3 that this α at $\Delta G^{0'} = 0$ or $\eta = 0$ is given in the classical treatment by

$$\alpha = \frac{1}{2} + \frac{\lambda_i}{4\lambda} \sum_s \frac{f_s^r - f^r p_s}{f_s^r + f_s^p} \frac{\lambda_{si}}{\lambda}, \tag{69}$$

where λ_i is the vibrational ('inner-coordination shell') contribution to λ (only the force constants f in the inner-coordination shell were assumed to change) and where λ_{si} is the contribution of coordinate q_s to λ .

We apply Eqn. 69 next to consider one model that has recently been assumed [43] for the solvation of singly charged organic ions. In this model the organic ion-solvent interaction is represented by a harmonic term with a force constant (assumed to be the same for cations as for anions) which exceeds considerably the force constant for the interaction of the solvent with an uncharged organic molecule [43]. Such a model clearly does not obey the quadratic free-energy relation used in obtaining Eqns. 1-3 and would predict from Eqn. 69 that the slope of a $-RT \ln k$ vs. $\Delta G^{0'}$ plot at $\Delta G^{0'} = 0$ would be quite different from 0.5 for reactions between uncharged molecules to form ions, such as $A + B \rightarrow A^+ + B^-$ or for the reverse reaction. (In reactions such as $A^+ + B \rightarrow A + B^+$ or $A + B^- \rightarrow A^- + B$, there is a compensating effect in Eqn. 69, so even quite different force constants $f_s^{\rm r}$ and $f_s^{\rm p}$ can lead [3] to a slope near 0.5.) Similar remarks apply to electrochemical oxidations and reductions at $\eta = 0$.

The above result, namely of deviations from 0.5 arising from the model in Ref. 43 can be shown by adapting Eqn. 69 so as to include that model. There were only the ion-solvent coordinates in Ref. 43, and we will write $\lambda = \lambda_i$. Using the model the last term in Eqn. 69 then becomes $\frac{1}{4}(f^r - f^p)/(f^r + f^p)$. The value of $f^r/(f^p - f^r)$ assumed in Ref. 43 for a reaction such as $A^* + B \rightarrow A^- + B^+$ was between 0.1 and 0.3 ¶. It yields an α ,

[§] Use of the square-barrier model for β mentioned in subsection IVC would be more artificial and yields a β varying as $V_0^{1/2}$ instead of as $\ln V_0$. Also, hole transfer might occur, but for brevity of presentation only electron transfer is considered.

However, the treatment in Refs. 318 and 43 differed from the present one. It was classical for the ion-solvent coordinate and quantum for the other modes in Ref. 318 while in Ref. 43 it was 'semiclassical' in the sense [63,64] discussed in subsection IIB for the ion-solvent coordinate with a high (800 cm⁻¹) assumed frequency for that coordinate and with no other modes.

based on Eqn. 69, of between 0.66 and 0.71 for such a homogeneous reaction or, in an electrochemical reaction, for the formation of an ion from an uncharged molecule. (It yields a value of

 α between 0.34 and 0.29 for the reverse reaction.)

These α 's differ considerably from the experimental α 's (approx. 0.5) given in Table III at $\Delta G^{0'} = 0$ for the organic compounds and differ from the

typical values of approx. 0.5 at $\eta = 0$ for the electrochemical reduction or oxidation of organic molecules [319-325] §. The solvent model proposed in Ref. 43 was

introduced to explain why the inverted effect was not found [326] for the intermolecular electron transfer $A^* + B \rightarrow A^{\pm} + B^{\mp}$. Recently, however, the inverted effect was found for the intramolecular $A^* - B \rightarrow A^+ - B^-$ reaction [38], complementing the recent intramolecular and frozen-solvent observations of the inverted effect [36,37] for reactions of the type $A^- - B \rightarrow A - B^-$. Moreover, the bridges thus introduced are not expected to have any significant effect on the previously postulated

the inverted effect in Ref. 326 lies in some direction other than specific ion-solvent forces. For example, the quenching of the fluorescence of A* may arise not only from an electron transfer, but also (particularly if the electron transfer becomes slow) by competitive pathways, such as exciplex formation followed by a radiationless transition

(e.g., Ref. 55 or references cited in Ref. 318). A variety of other possibilities exist [55]. However, a

further analysis of the incidence and consequences

of specific ion-solvent interactions would be de-

sirable, as well as further pursuit of intramolecular examples of reactions such as $A^* - B \rightarrow A^+ - B^-$.

[43,318] specific ion-solvent forces. All of the re-

sults suggest that the explanation for the lack of

Appendix II. Franck-Condon factors and absolute rates

The calculation of Franck-Condon factors has been discussed by many authors, both for radia-

given by Eqn. 69 differs slightly (approx. 0.04) from 0.5 when typical differences of vibrational force constants in the

tionless transitions and for electron transfers. Of interest in the present case are the calculations for the case where many coordinates are present (the

environment). For a reaction proceeding from the lowest state of the reactant(s) to yield quantum states of a set of N degenerate vibrational modes of the product(s), each having the same frequency

$$\nu$$
, the rate constant is found from Eqn. 13 to be given by ¶ :
$$k = \frac{2\pi}{\hbar} H_{AB}^2 \sum_{v=0}^{\infty} \frac{e^{-S}S^v}{\Gamma(v+1)} \delta(\Delta E^{0'} + vh\nu). \tag{70}$$

(70)

In Eqn. 70, ν is assumed to be the same for the products as for the reactants. The λ appearing in $S(S = \lambda/h\nu)$ is N-times the value of λ for an individual vibrational mode; δ is the Dirac delta function; v is the principal quantum number for the degenerate states of the N vibrationally-excited modes. Eqn. 70 is intended to be an approximation to the actual situation where the frequencies of the vibrational modes are not equal. In the latter case, there are many largely nondegenerate levels thermally occupied rather than only a relatively small number of highly degenerate levels. The sum in Eqn. 70 can then be replaced by an integral over v, and, on making use of the delta function, Eqn. 71 is then obtained:

$$k = \frac{2\pi}{\hbar} H_{AB}^2 \frac{e^{-S} S^v}{h \nu \Gamma(v+1)}, \qquad (71)$$

where ν is a mean frequency (defined for example

in Ref. 327); v now denotes $-\Delta E^{0'}/h\nu$ and need

no longer be an integer. Since any change in ν is

obtained, for example, in Ref. 327, Eqn. 9.16. This reference

also contains the idea of using a mean frequency when there

Ref. 27. When the latter is used and the limit $\lambda_0 \rightarrow 0$ is

[§] Numerous electrochemical results on metal complexes are also available and typically have slopes of about 0.5 at $\eta = 0$ [132,133]. The theoretical value for these metal complexes

oxidized and reduced forms of the reactant are taken into account [3]. The Franck-Condon factors which appear in Eqn. 13 are

are a number of contributing vibrational modes. One derivation of the equations for the case of one coordinate is given in Ref. 328 for a spectral band shape. The Franck-Condon factor in Eqn. 70 is obtained by setting $h\nu = 0$ and $h\nu_{ab} =$ ΔE^{0} in Eqn. 4-115 of Ref. 328. Eqn. 70 is related to Eqn. 19, or really to a predecessor of that equation, namely, Eqn. 5 of

introduced, one obtains Eqn. 70. (One uses the fact that when $a \to 0$, $(1/a \sqrt{\pi}) \exp(-x^2/a^2)$ becomes $\delta(x)$.) However, Eqn. 70 can be derived more directly, as in Ref. 328.

neglected, as is any other contribution to $\Delta S^{0'}$, this v also equals $-\Delta G^{0'}/hv$.

When higher vibrational levels of the reactants are also involved, e.g., at finite temperatures, the expression for k in Eqn. 13 contains Boltzmann-weighted Franck-Condon factors and can be written as \$:

$$k = \frac{2\pi}{\hbar h \nu} H_{AB}^2 e^{\nu y - S \coth y} I_{\nu}(S \operatorname{cosech} y)$$
 (72)

where $y = h\nu/2RT$, I_v is a modified Bessel function, and the remaining symbols have their previous meaning.

We consider next some properties of these equations. At sufficiently high temperatures Eqn. 72 reduces to Eqn. 14[†], while at sufficiently low temperatures it reduces to Eqn. 71. Further, the k given by Eqn. 71 has a maximum as a function of v (i.e., of $-\Delta G^{0'}/h\nu$) which occurs when $v \approx S$ (cf. Ref. 51). This result is seen by introducing Stirling's formula for $\Gamma(v+1)$, namely $\sqrt{2\pi v} (v/e)^v$, maximizing the expression in Eqn. 71, and neglecting the variation in \sqrt{v} relative to that in $(v/e)^v$. Thus, the maximum occurs at $\lambda \approx$ $-\Delta G^{0}$, just as it did for the classical expression, Eqn. 14. Further, it was seen earlier from Eqn. 35 that the activation energy for the classical expression is zero at $\Delta G^{0'} = -\lambda$. This same choice for λ also leads to a similarly low activation energy when Eqn. 71 is used for the value of k at 0 K and Eqn. 14 for the values at high temperatures, or hence when Eqn. 72 is used over the entire temperature range.

We next apply Eqns. 71-72 and 14 to the kinetic data given in subsection IVD for the photosynthetic barrierless regime. In that regime, as already noted, $-\Delta G^{0'} \approx \lambda$. It will be assumed, for the present purposes, that there is an averaged vibrational frequency ν for the contributing modes of about 200 cm⁻¹ (i.e., $h\nu$ is 2.5 kJ·mol⁻¹). (Larger quantum-mechanical corrections occur if ν is larger than the frequency used here [131].) If the driving force $-\Delta G^{0'}$ in the first step (or two steps)

for electronically excited $(BChl)_2 \rightarrow BChl \rightarrow BPh$ is about 0.25 eV total, then the value per step is about 0.13 eV or approx. 12.5 kJ·mol⁻¹, and so, within the framework of the above analysis, a typical value of λ for these reactants is expected to be in the neighborhood of 12.5 kJ·mol⁻¹ if the reaction is to be barrierless. We consider several value of λ below in the range 5-38 kJ·mol⁻¹, together with the above values for $h\nu$ and $-\Delta G^{0'}$.

We first compare at 25°C the classical Eqn. 14 with the quantum Eqn. 72 by comparing the factor multiplying $2\pi H_{AB}^2/\hbar h \nu$ in those equations. (In Eqn. 14 this factor is $h\nu/\sqrt{4\pi\lambda RT}$ -times the exponential.) For the above factor such a comparison at 25°C yields, respectively, 0.063 and 0.063 ($\lambda = 5.0 \text{ kJ} \cdot \text{mol}^{-1}$), 0.13 and 0.12 ($\lambda = 8.4 \text{ kJ} \cdot \text{mol}^{-1}$), 0.10 and 0.10 ($\lambda = 17 \text{ kJ} \cdot \text{mol}^{-1}$), and 0.048 and 0.050 ($\lambda = 25 \text{ kJ} \cdot \text{mol}^{-1}$). Thus, the classical Eqn. 14 and the quantum Eqn. 72 agree closely when $h\nu/2RT$ is not large. (Here, it is 0.5 at 25°C.) Analytically, Eqn. 14 is the 'high-temperature' limit of Eqn. 72, as already noted.

The temperature dependence of k is seen by comparing the above results at 25°C with those obtained at 0 K using Eqn. 71: at 0 K the above factors are found to be 0.036, 0.12, 0.14 and 0.038 for $\lambda = 5.0$, 8.4, 17 and 25 kJ·mol⁻¹, respectively. Thus, the results for $\lambda \le 25$ kJ·mol⁻¹ show negligible temperature dependence for the assumed value of $-\Delta G^{0'}$. At $\lambda = 38$ kJ·mol⁻¹ there is, however, a significant temperature dependence: the factor mentioned above calculated from Eqn. 14 is 0.014 at 25°C and calculated from Eqn. 71 is 0.0019 at 0 K. Larger λ 's will show an even larger temperature dependence.

We consider next the absolute value of k. Since Eqns. 14 and 72 are seen above to give similar values for k at room temperature in the general neighborhood of $\lambda = -\Delta G^{0'}$ we shall simply use the classical equations 31-33b for the present calculation of k. When $\Delta G^{0'} \approx -\lambda$, $d \approx 0.1$ nm and $\beta \approx 12$ nm⁻¹, Eqns. 31-33b yield a value of $3 \cdot 10^{12}$ s⁻¹, or a lifetime of 0.3 ps for the reaction, while for $\lambda = 5.0$ kJ·mol⁻¹, $k = 1 \cdot 10^{12}$ s⁻¹, which correspond to a lifetime of 1 ps. A λ larger than 25 kJ·mol⁻¹ will yield, for the given $\Delta G^{0'}$, a smaller k but will also give a significant temperature coefficient. The experimental value for the overall transfer is, as noted in the text, 4 to 5 ps.

[§] Cf. previous footnote.

[†] At high temperatures S coth y and S cosech y both become $2\lambda RT/(h\nu)^2$. Use of the high temperature form [328] of $I_v(x)$, namely $(2\pi x)^{-1/2} \exp(x - v^2/2x)$, where $x = 2\lambda RT/(h\nu)^2$, then yields Eqn. 14 from Eqn. 72.

are high frequency modes (in addition to the others) which contribute a Franck-Condon factor, such as the one in Eqn. 71, that is temperature-independent over the entire investigated temperature

Thus, perhaps the assumed $1 \cdot 10^{13}$ s⁻¹ in Eqn.

33b is too high a value for the orientational config-

uration of the molecules involved, or perhaps there

range, 5-300 K. The latter factor makes the rate lower by effectively contributing a temperature-independent factor to the rate constant expression. We turn next to the k for the BPh⁻+ Q reac-

tion. With a driving force in the vicinity of 0.5 to 0.6 eV (subsection IVD) and a λ of about this value (subsection IVD), the reaction is again barrierless. The values obtained from Eqns. 14 and 71, or 72 may again be compared: with $\Delta G^{0'} \cong$ $-\lambda = -0.5$ eV, and $h\nu \approx 200$ cm⁻¹ the factor multiplying $2\pi H_{AB}^2/\hbar h\nu$ in Eqn. 71 is 0.09, and in Eqn. 14 it is $h\nu/(4\pi\lambda RT)^{1/2}$, namely, 0.06. Thus,

once again the temperature coefficient is minor, as

expected. Use of Eqns. 31-33b with $d \approx 1.0$ nm

and $\beta \cong 12 \text{ nm}^{-1}$ (subsection VD), yields the value

of $5 \cdot 10^9$ s⁻¹ in the text, corresponding to a

lifetime of 200 ps. Various authors have treated the rate constants in the photosynthetic system and used a wide variety of vibration frequencies and models (see for example, Refs. 306, 329 and 330 and see also Kirmaier, C., Holten, D. and Parson, W.W., unpublished results). The negative temperature effect

for the BPh $^-$ + Q reaction of a factor of two in khas been interpreted by the latter authors in terms of differences of vibration frequencies of reactants and products. However, in the present calculations, factors of two are not the focus of attention and possible effects of differences in vibration frequencies of reactants and products have been

Acknowledgements

omitted.

We wish to acknowledge helpful discussions with Drs. J. Fajer, H.B. Gray, G. McLendon, T.L. Netzel, I. Pecht, A.G. Sykes and J.K. Yandell. We are also grateful to Dr. D. Holten for making

kinetic data available prior to publication and Drs.

contract DE-AC02-76CH00016 with the U.S. Department of Energy and supported by its Division of Chemical Sciences, Office of Basic Energy Sciences, while that at Caltech (R.A.M.) was supported by the National Science Foundation and the Office of Naval Research (contribution No. 7095 from Caltech).

References

1 Marcus, R.A. (1956) J. Chem. Phys. 24, 966-978 2 Marcus, R.A. (1960) Disc. Faraday Soc. 29, 21-31

3 Marcus, R.A. (1965) J. Chem. Phys. 43, 679-701 4 Marcus, R.A. (1968) J. Phys. Chem. 72, 891-899

5 Kresge, A.J. (1975) Acc. Chem. Res. 8, 354-360

6 Albery, W.J. (1980) Annu. Rev. Phys. Chem. 31, 227-263 7 Albery, W.J. and Kreevoy, M.M. (1978) Adv. Phys. Org. Chem. 16, 87-157 8 Donnella, J. and Murdoch, J.R. (1984) J. Am. Chem. Soc. 106, 4724-4735

9 Lewis, E.S. and Hu, D.D. (1984) J. Am. Chem. Soc. 106, 3292-3296

tional Laboratory (N.S.) was carried out under

10 Roberts, R.M.G., Ostovic, D. and Kreevoy, M.M. (1982) Faraday Disc. Chem. Soc. 74, 257-265 11 Kreevoy, M.M. and Lee, I.-S.H. (1984) J. Am. Chem. Soc. 106, 2550-2553 12 Magnoli, D.E., Murdoch, J.R. (1981) J. Am. Chem. Soc.

103, 7465-7469 13 Pellerite, M.J. and Brauman, J.I. (1980) J. Am. Chem. Soc. 102, 5993-5999 14 Wolfe, S., Mitchell, D.J., and Schlegel, H.B. (1981) J. Am. Chem. Soc. 103, 7694-7696

15 Chen, M.Y. and Murdoch, J.R. (1984) J. Am. Chem. Soc. 106, 4735-4743 16 Murdoch, J.R. (1982) Faraday Disc. Chem. Soc. 74,

297-306 17 Cannon, R.D. (1980) Electron Transfer Reactions, Butterworths, London 18 Reynolds, W.L. and Lumry, R.W. (1966) Mechanisms of Electron Transfer, Ronald Press, New York

19 DeVault, D. (1980) Q. Rev. Biophys. 13, 387-564 20 DeVault, D. (1984) Quantum-Mechanical Tunnelling in Biological Systems, 2nd Edn., Cambridge University Press, New York

21 Sutin, N. (1983) Progr. Inorg. Chem. 30, 441-498 22 Sutin, N. (1973) in Inorganic Biochemistry (Eichhorn, G.L., ed.) Vol. 2, pp. 611-653, Elsevier, New York 23 Newton, M.D. and Sutin, N. (1984) Annu. Rev. Phys.

Chem. 35, 437-480 24 Marcus, R.A. (1979) Life Sci. Res. Rep. 12, 15-43 25 Marcus, R.A. (1964) Annu. Rev. Phys. Chem. 15, 155-196

26 Marcus, R.A. (1957) J. Chem. Phys. 26, 867-871 27 Marcus, R.A. (1982) Faraday Disc. Chem. Soc. 74, 7-15

28 Larsson, S. (1983) J. Chem. Soc. Faraday Trans. 2, 79, 1375-1388 29 Marcus, R.A. (1957) ONR Tech. Rept. No. 12, Project

P.C. Weber and F.R. Salemme for modeling the

docking of two cytochrome c molecules and for providing Fig. 9. Research at Brookhaven Na-

NR-051-331; reproduced (1977) in Special Topics in Elec-

- trochemistry (Rock, P.A. ed.), pp. 180-209, Elsevier, New York
- 30 Marcus, R.A. (1959) Can. J. Chem. 37, 155-163
- 31 Marcus, R.A. (1963) J. Phys. Chem. 67, 853-857, 2889
- 32 Landau, L.D. and Lifshitz, E.M. (1965) Quantum Mechanics, 2nd Edn., pp. 177-180, Pergamon Press, New York
- 33 Marcus, R.A. (1965) J. Chem. Phys. 43, 2654-2657
- 34 Marcus, R.A. (1970) J. Chem. Phys. 52, 2803-2804
- 35 Wallace, W.L. and Bard, A.J. (1979) J. Phys. Chem. 83, 1350-1357
- 36 Miller, J.R., Calcaterra, L.T., and Closs, G.L. (1984) J. Am. Chem. Soc. 106, 3047-3049
- 37 Miller, J.R., Beitz, J.V. and Huddleston, R.K. (1984) J. Am. Chem. Soc. 106, 5057-5068
- 38 Wasielewski, M.R., Niemczyk, M.P., Svec, W.A., Pewitt, E.B. (1985) J. Am. Chem. Soc. 107, 1080-1082
- 39 Allen, A.O. Gangwer, T.E. and Holroyd, R.A. (1975) J. Phys. Chem. 79, 25-31
- 40 Hush, N.S. (1961) Trans. Faraday Soc. 57, 557-580
- 41 George, P. and Griffith, J.S. (1959) in The Enzymes (Boyer, P.D., Lardy, H. and Myrbäck, K., eds.), 2nd Edn., Vol. 1, pp. 347-389, Academic Press, New York
- 42 Orgel, L.E. (1956) Inst. Intern. Chim. Solvay Conseil Chim. Brussels 10, 289-338
- 43 Kakitani, T. and Mataga, N. (1985) J. Phys. Chem. 89, 8-10
- 44 Sutin, N. (1966) Ann. Rev. Phys. Chem. 17, 119-172
- 45 Levich, V.G. and Dogonadze, R.R.(1959) Dokl. Acad. Nauk SSSR 124, 123-126 (Proc. Acad. Sci. Phys. Chem. Sect. 124, 9-13)
- 46 Levich, V.G. (1966) Adv. Electrochem. Electrochem. Eng. 4, 249-371
- 47 Dogonadze, R.R., Kuznetsov, A.M. and Vorotyntsev, M.A. (1972) Phys. Stat. Sol. B 54, 125-134 and 425-433
- 48 Van Duyne, R.P. and Fischer, S.F. (1974) Chem. Phys. 5, 183-197
- 49 Kestner, N.R., Logan, J. and Jortner, J. (1974) J. Phys. Chem. 78, 2148–2166
- 50 Ulstrup, J. and Jortner, J. (1975) J. Phys. Chem. 63, 4358-4368
- 51 Jortner, J. (1976) J. Chem. Phys. 64, 4860-4867
- 52 Efrima, S. and Bixon, M. (1976) Chem. Phys. 13, 417-460
- 53 Siders, P. and Marcus, R.A. (1981) J. Am. Chem. Soc. 103, 741–747
- 54 Schmidt, P.P. (1978) Electrochemistry Spec. Period. Rep. 6, 128-241
- 55 Marcus, R.A. and Siders, P. (1982) J. Phys. Chem. 86, 622-630
- 56 Warshel, A. (1982) J. Phys. Chem. 86, 2218-2224
- 57 Ulstrup, J. (1979) Charge Transfer Processes in Condensed Media, Lecture Notes in Chemistry, No. 10, Springer-Verlag, New York
- 58 Marcus, R.A. (1984) J. Chem. Phys. 81, 4494-4500
- 59 Marcus, R.A. (1973) Faraday Disc. Chem. Soc. 55, 34-44
- 60 Marcus, R.A. and Sutin, N. (1975) Inorg. Chem. 14, 213–216
- 61 Siders, P. and Marcus, R.A. (1981) J. Am. Chem. Soc. 103, 748-752

- 62 Buhks, E., Bixon, M., Jortner, J. and Navon, G. (1981) J. Phys. Chem. 85, 3759-3762
- 63 Hopfield, J.J. (1974) Proc. Natl. Acad. Sci. USA 71, 3640–3644
- 64 Hopfield, J.J. (1977) in Electrical Phenomena at the Biological Membrane Level (Roux, E., ed.), pp. 471-492, Elsevier, New York
- 65 Marcus, R.A. (1982) in Oxidases and Related Redox Systems (King, T.E., Mason, H.S. and Morrison, M., eds.), pp. 3-19, Pergamon Press, Oxford
- 66 Holstein, T. (1979) in Tunneling in Biological Systems (Chance, B., DeVault, D.C., Frauenfelder, H., Marcus, R.A., Schrieffer, J.B. and Sutin, N., eds.), pp. 129-136, Academic Press, New York
- 67 Holstein, T. (1978) Philos. Mag. B, 37, 49-62
- 68 Scher, H. and Holstein, T. (1981) Philos Mag. B, 44, 343-356
- 69 Brunschwig, B.S., Logan, J., Newton, M.D. and Sutin, N. (1980) J. Am. Chem. Soc. 102, 5798-5809
- 70 Larsson, S. (1981) J. Am. Chem. Soc. 103, 4031-4040
- 71 Beratan, D.N. and Hopfield, J.J. (1984) J. Chem. Soc. 106, 1584–1594
- 72 Siders, P., Cave, R.J. and Marcus, R.A. (1984) J. Chem. Phys. 81, 5613-5624
- 73 Reference deleted
- 74 Reference deleted
- 75 Chang, R. and Weismann, S.I. (1967) J. Am. Chem. Soc. 89, 5968
- 76 Tembe, B.L., Friedman, H.L. and Newton, M.D. (1982) J. Chem. Phys. 76, 1490-1507
- 77 Newton, M.D. (1982) Am. Chem. Soc. Symp. Ser. 198, 255-279
- 78 Brunschwig, B.S., Creutz, C., Macartney, D.H., Sham, T.K. and Sutin, N. (1982) Faraday Disc. Chem. Soc. 74, 113-127
- 79 Chou, M., Creutz, C. and Sutin, N. (1977) J. Am. Chem. Soc. 99, 5615–5623
- 80 Weaver, M.J. and Yee, E.L. (1980) Inorg. Chem. 19, 1936-1945
- 81 Pennington, D.E. (1978) in Coordination Chemistry (Martell, A.E., ed.) Vol. 2, Am. Chem. Soc. Monograph No. 174, pp. 476-590
- 82 Frese, Jr., K.W. (1981) J. Phys. Chem. 85, 3911-3916
- 83 Hupp, J.T. and Weaver, M.J. (1983) Inorg. Chem. 22, 2557-2564
- 84 Meyer, T.J. and Taube, H. (1968) Inorg. Chem. 7, 2369-2379
- 85 Campion, R.J., Purdie, N. and Sutin, N. (1964) Inorg. Chem. 3, 1091–1094
- 86 Thomas, L. and Hicks, K.W. (1974) Inorg. Chem. 13, 749-752
- 87 Rawoof, M.A. and Sutter, J.R. (1967) J. Phys. Chem. 71, 2767-2771
- 88 Böttcher, W., Brown, G.M. and Sutin, N. (1979) Inorg. Chem. 18, 1447-1451
- 89 Lin, C.-T., Böttcher, W., Chou, M., Creutz, C. and Sutin, N. (1976) J. Am. Chem. Soc. 98, 6536-6544
- 90 Endicott, J.F., Durham, B. and Kumar, K. (1982) Inorg. Chem. 21, 2437–2444

- 91 Silverman, J. and Dodson, R.W. (1952) J. Phys. Chem. 56, 846-851
- 92 Bernhard, P., Bürgi, H.B., Hauser, J., Lehmann, H. and
- Ludi, A. (1982) Inorg. Chem. 21, 3936-3941
 - 93 Young, R.C., Keene, F.R. and Meyer, T.J. (1977) J. Am. Chem. Soc. 99, 2468-2473
 - 94 Hammershoi, A., Geselowitz, D. and Taube, H. (1984) Inorg. Chem. 23, 979-982
 - 95 Neumann, H.M., quoted in: Farina, R. and Wilkins, R.G.
- (1968) Inorg. Chem. 7, 514-518
- 96 Brown, G.M. and Sutin, N. (1979) J. Am. Chem. Soc. 101,
- 883-892 97 Li, T.T.-T. and Brubaker, C.H., Jr. (1981) J. Organomet.
 - Chem. 216, 223-234 98 Chan, M.-S. and Wahl, A.C. (1982) J. Phys. Chem. 86,
 - 126-130
 - 99 Brandon, J.R. and Dorfman, L.M. (1970) J. Chem. Phys. 53, 3849-3856
- 100 Yang, E.S., Chan, M.-S. and Wahl, A.C. (1980) J. Phys. Chem. 84, 3094-3099 101 Creutz, C. and Sutin, N. (1985) in Inorganic Reactions and
 - Methods (Zuckerman, J.J., ed.), Springer-Verlag, in the
- 102 Ford-Smith, M.H. and Sutin, N. (1961) J. Am. Chem. Soc.
- 83, 1830-1834
- 103 Dulz, G. and Sutin, N. (1963) Inorg. Chem. 2, 917-921 104 Brunschwig, B.S. and Sutin, N. (1979) Inorg. Chem. 18,
- 105 Brunschwig, B.S. and Sutin, N. (1978) J. Am. Chem. Soc. 100, 7568-7577
- 106 Pelizzetti, E., Mentasti, E. and Pramauro, E. (1978) Inorg. Chem. 17, 1181-1186 107 Macartney, D.H. and Sutin, N. (1983) Inorg. Chim. Acta,
- 74, 221-228
- 108 Pelizzetti, E. and Mentasti, E. (1979) Inorg. Chem. 18, 583-588
- 109 Fukuzumi, S., Wong, C.L., and Kochi, J.K. (1980) J. Am.
- Chem. Soc. 102, 2928-2939
- 110 Bock, C.P., Connor, J.A., Gutierrez, A.R., Meyer, T.J.,
- Whitten, D.G., Sullivan, B.P. and Nagle, J.K. (1979) J. Am. Chem. Soc. 101, 4815-4824
- 111 Meyer, T.E., Przysiecki, C.T., Watkins, J.A., Bhattacharyya, A., Simondsen, R.P., Cusanovich, M.A. and Tollin, G. (1983) Proc. Natl. Acad. Sci. USA 80, 6740-6744 112 Hush, N.S. (1967) Prog. Inorg. Chem. 8, 391-444
- 113 Creutz, C. (1983) Prog. Inorg. Chem. 30, 1-73 114 Marcus, R.A. (1965) J. Chem. Phys. 43, 1261-1274 115 Marcus, R.A. and Sutin, N. (1985) Comments Inorg. Chem.
- 4, in the press 116 Anderson, P.W. (1965) in Magnetism (Rado, G.T. and Suhl, H., eds.), Vol. 1, pp. 25-81, Academic Press, New
- 117 Okamura, M.Y., Fredkin, D.R., Isaacson, R.A. and Feher, G. (1979) in Tunneling in Biological Systems (Chance, B., DeVault, D.C., Frauenfelder, H., Marcus, R.A., Schrieffer,
 - J.B. and Sutin, N., eds.), pp. 729-743, Academic Press,

- 119 Sutton, J.E., Sutton, P.M. and Taube, H. (1979) Inorg. Chem. 18, 1017-1021
- 120 Rieder, K. and Taube, H. (1977) J. Am. Chem. Soc. 99,
 - 7891-7894 121 Creutz, C. and Taube, H. (1973) J. Am. Chem. Soc. 95,
 - 1086-1094 122 Fürholz, U., Burgi, H.-B., Wagner, F.E., Stebler, A., Amm-
 - eter, J.H., Krausz, E., Clark, R.J.H., Stead, M.J. and Ludi, A. (1984) J. Am. Chem. Soc. 106, 121-123
 - 123 Root, L.J. and Ondrechen, M.J. (1982) Chem. Phys. Lett.
 - 93, 421-424 124 Ondrechen, M.J., Ellis, D.L. and Ratner, M.A. (1984)
 - Chem. Phys. Lett. 109, 50-55 125 Ko, J. and Ondrechen, M.J. (1984) Chem. Phys. Lett. 112,
 - 507-512 126 Curtis, J.C. and Meyer, T.J. (1982) Inorg. Chem. 21,
- 1562-1571 127 Stein, C.A. and Taube, H. (1981) J. Am. Chem. Soc. 103,
- 693-695 128 Stein, C.A., Lewis, N.A. and Seitz, G. (1982) J. Am. Chem.
 - Soc. 104, 2596-2599 129 Redi, M. and Hopfield, J.J. (1980) J. Chem. Phys. 72,
 - 6651-6660 130 Delahay, P. and Dziedzic, A. (1984) J. Chem. Phys. 80,
- 5381-5387 131 Delahay, P. and Dziedzic, A. (1984) J. Chem. Phys. 80, 5793-5798
- 132 Saji, T., Maruyama, Y., and Aoyagui, S. (1978) J. Électroanal. Chem. 86, 219-222
- 133 Weaver, M.J. (1980) J. Phys. Chem. 84, 568-576 134 Weaver, M.J. and Hupp, J.T. (1982) Am. Chem. Soc.
- Symp. Series 198, 181-208
- 135 Saveant, J.-M. and Tessier, D. (1982) Faraday Disc. Chem. Soc. 74, 57-72
- 136 Rackovsky, S. and Goldstein, D.A. (1984) Proc. Natl. Acad. Sci. USA 81, 5901-5905
- 137 Churg, A.K., Weiss, R.M., Warshel, A. and Takano, T. (1983) J. Phys. Chem. 87, 1683-1694
- 138 Alexandrov, I.V., Khairutdinov, R.F. and Zamaraev, K.I. (1978) Chem. Phys. 32, 123-141 139 Miller, J.R. (1975) Science 189, 221-222
- 140 Reference deleted 141 Miller, J.R. and Beitz, J.V. (1981) J. Chem. Phys. 74,
 - 6746-6756 142 Calcaterra, L.T., Closs, G.L. and Miller, J.R. (1983) J. Am. Chem. Soc. 105, 670-671
- 143 Mann, B. and Kuhn, H. (1971) J. Appl. Phys. 42, 4398-4405

144 Polymeropoulos, E.E. (1977) J. Appl. Phys. 48, 2404-2407

145 Sugi, M., Fukui, T. and Iizima, S. (1975) Appl. Phys. Lett.

- 27, 559-561 146 Sugi, M., Nembach, K., Möbius, D. and Kuhn, H. (1974)
- Sol. Stat. Comm. 15, 1867-1870
- 147 Kuhn, H. (1979) J. Photochem. 10, 111-132 148 Killesreiter, H. and Baessler, H. (1971) Chem. Phys. Lett.

51, 657-668

- 11, 411-414 149 Killesreiter, H. and Baessler, H. (1972) Phys. Stat. Sol. B
- 118 Creutz, C. (1978) Inorg. Chem. 17, 3723-3725

New York

- 150 Lindsey, J.S., Mauzerall, D.C., and Linschitz, H. (1983) J. Am. Chem. Soc. 105, 6528-6529
- 151 Kostić, N.M., Margalit, R., Che, C.-M. and Gray, H.B.
- (1983) J. Am. Chem. Soc. 105, 7765-7767 152 McCourty, J.L., Blough, N.V. and Hoffman, B.M. (1983)
- J. Am. Chem. Soc. 105, 4470-4472 153 Peterson-Kennedy, S.E., McGourty, J.L. and Hoffman,
- B.M. (1984) J. Am. Chem. Soc. 106, 5010-5012 154 McLendon, G.L., Winkler, J.R., Nocera, D.G., Mauk,
 - M.R., Mauk, A.G. and Gray, H.B. (1985) J. Am. Chem. Soc., 107, 739-740
- 155 Nocera, D.G., Winkler, J.R., Yokom, K.M., Bordignon, E. and Gray, H.B. (1984) J. Am. Chem. Soc. 106, 5145-5150
- 156 Isied, S.S., Kuehn, C. and Worosila, G. (1984) J. Am.
- Chem. Soc. 106, 1722-1726 157 Isied, S.S. (1984) Progr. Inorg. Chem. 32, 443-517
- 158 Brunschwig, B.S., Ehrenson, S. and Sutin, N. (1984) J. Am. Chem. Soc. 106, 6858-6859
- 159 Wherland, S. and Gray, H.B. (1977) in Biological Aspects of Inorganic Chemistry, (Addison, A.W., Cullen, W.R., Dolphin, D. and James, B.R., eds.), pp. 289-368, Wiley, New York
- 160 Margalit, R. and Schejter, A. (1973) Eur. J. Biochem. 32, 492-499
- 161 Koppenol, W.H., Vroonland, C.A.J. and Braams, R. (1978) Biochim, Biophys, Acta 503, 499-508
- 162 Morton, R.A., Overnell, J. and Harbury, H.A. (1970) J. Biol. Chem. 245, 4653-4657
- 163 Goldkorn, T. and Schejter, A. (1979) J. Biol. Chem. 254, 12562-12566
- 164 McArdle, J.V., Coyle, C.L., Gray, H.B., Yoneda, G.S. and Holwerda, R.A. (1977) J. Am. Chem. Soc. 99, 2483-2489
- 165 Rawlings, J., Wherland, S. and Gray, H.B. (1977) J. Am. Chem. Soc. 99, 1968-1971 166 Segal, M.G. and Sykes, A.G. (1978) J. Am. Chem. Soc.
- 100, 4585-4592
- 167 Sailasuta, N., Anson, F.C. and Gray, H.B. (1979) J. Am. Chem. Soc. 101, 455-458
- 168 Yee, E.L. and Weaver, M.J. (1980) Inorg. Chem. 19, 1077-1079
- 169 Feinberg, B.A., Ryan, M.D. and Wei, J.-F. (1977) Biochem. Biophys. Res. Commun. 79, 769-775
- 170 Feinberg, B.A. and Ryan, M.D. (1981) in Topics in Bioelectrochemistry and Bioenergetics (Milazzo, G., ed.), Vol.
- 4, 225-270, Wiley and Sons, New York 171 Rosenberg, R.C., Wherland, S., Holwerda, R.A. and Gray, H.B. (1976) J. Am. Chem. Soc. 98, 6364-6369
- 172 Koppenol, W.H. (1980) Biophys. J. 29, 493-507
- 173 Koppenol, W.H. and Margoliash, E. (1982) J. Biol. Chem.
- 257, 4426-4437 174 Van Leeuwen, J.W., Mofers, F.J.M. and Veerman, E.C.I.
- (1981) Biochim. Biophys. Acta, 635, 431-439
- 175 Van Leeuwen, J.W. (1983) Biochim. Biophys. Acta 743, 408-421
- 176 Bronsted, J.N. (1922) J. Am. Chem. Soc. 44, 877-898
- 177 Meyer, T.E., Watkins, J.A., Przysiecki, C.T., Tollin, G. and Cusanovich, M.A. (1984) Biochem., 23, 4761-4767
- 178 Chapman, S.K., Sinclair-Day, J.D., Sykes, A.G., Tam, S.C.

- and Williams, R.J.P. (1983) J. Chem. Soc., Chem. Commun., 1152-1154
- 179 Beoku-Betts, D. and Sykes, A.G. (1985) Inorg. Chem. 24, 1142-1147
- 180 Matthew, J.B., Weber, P.C., Salemme, F.R. and Richards, F.M. (1983) Nature 301, 169-171
- 181 Dickerson, R.E. and Timkovich, R., (1975) in The Enzymes (Boyer, P.D., ed.) 3rd Edn., Vol. XIA, pp. 397-547, Academic Press, New York
- 182 Takano, T. and Dickerson, R.E. (1981) J. Mol. Biol. 153, 95-115
- 183 Moore, G.R., Huang, Z.-X., Eley, C.G.S., Barker, H.A., Williams, G., Robinson, M.N. and Williams, R.J.P. (1982) Faraday Disc. Chem. Soc. 74, 311-329
- 184 Brown, G.M., Krentzien, H.J., Abe, M. and Taube, H. (1979) Inorg. Chem. 18, 3374-3379
- 185 Sutin, N. (1977) Am. Chem. Soc. Adv. Chem. Ser. 162, 156-172
- 186 Wilkins, R.G. and Yelin, R.E. (1968) Inorg. Chem. 7, 2667-2669 187 Campion, R.J., Deck, C.F., King, P. and Wahl, A.C.
- (1967) Inorg. Chem. 6, 672-681 188 McArdle, J.V., Gray, H.B., Creutz, C. and Sutin, N. (1974)
- J. Am. Chem. Soc. 96, 5737-5741 189 Ewall, R.X. and Bennett, L.E. (1974) J. Am. Chem. Soc.
- 96, 940-942 190 Cummins, D. and Gray, H.B. (1977) J. Am. Chem. Soc. 99, 5158-5167
- 191 Creutz, C. and Sutin, N. (1974) J. Biol. Chem. 249,
- 6788-6795 192 Hodges, H.L., Holwerda, R.A. and Gray, H.B. (1974) J.
- Am. Chem. Soc. 96, 3132-3137 193 Gupta, R.K. (1973) Biochim. Biophys. Acta 292, 291-295
- 194 Holwerda, R.A., Knaff, D.B., Gray, H.B., Clemmer, J.D., Crowley, R., Smith, J.M. and Mauk, A.G. (1980) J. Am. Chem. Soc. 102, 1142-1146
- 195 Butler, J., Davies, D.M., Sykes, A.G., Koppenol, W.H., Osheroff, N. and Margoliash, E. (1981) J. Am. Chem. Soc. 103, 469-471
- 196 Ohno, N. and Cusanovich, M.A. (1981) Biophys. J. 36, 589-605
- 197 Butler, J., Davies, D.M. and Sykes, A.G. (1981) J. Inorg.
- Biochem, 15, 41-53 198 Chapman, S.K., Davies, D.M., Watson, A.D. and Sykes,
- A.G. (1983) Am. Chem. Soc. Symp. Series 211, 177-194 199 Potasek, M.J. and Hopfield, J.J. (1977) Proc. Natl. Acad.
- USA 74, 229-233 200 Dickerson, R.E. (1979) in Tunneling in Biological Systems
- (Chance, B., DeVault, D.C., Frauenfelder, H., Marcus, R.A., Schrieffer, J.B. and Sutin, N. eds.), p. 545, Academic Press, New York
- 201 Austin, R.H. and Hopfield, J.J., cited in: Redi, M. and Hopfield, J.J. (1980) J. Chem. Phys. 72, 6651-6660
- 202 Harbury, H.A. and Marks, R.H.L. (1973) in Inorganic. Biochemistry, (Eichhorn, G.L., ed.), Vol. 2, pp. 902-954, Elsevier, New York
- 203 Reference deleted
- 204 Schomacker, K.T., Bangcharoenpaurpong, O. and Champion, P.M. (1984) J. Chem. Phys. 80, 4701-4717

- 205 Dixon, D.W., Barbush, M. and Shiragi, A. (1984) J. Am. Chem. Soc. 106, 4638-4639 206 Keller, R.M., Wüthrich, K. and Pecht, I. (1976) FEBS"
- Lett. 70, 180-184 207 Yokom, K.M., Ph.D. Thesis, California Institute of Tech-
- nology, Pasadena, CA., 1982 208 Margalit, R., Kostić, N.M., Che, C.-M., Blair, D.F., Chiang,
- H.-J., Pecht, I., Shelton, J.R., Schroeder, W.A. and Gray, H.B. (1984) Proc. Natl. Acad. Sci. USA 81, 6554-6558 209 Winkler, J.R., Nocera, D.G., Yokom, K.M., Bordignon, E.
- and Gray, H.B. (1982) J. Am. Chem. Soc. 104, 5798-5800 210 Cho, K.C., Che, C.M., Cheng, F.C. and Choy, C.L. (1984)
- J. Am. Chem. Soc., 106, 6843-6844 211 Dutton, P.L., Leigh, J.S., Prince, R.C. and Tiede, D.M.
 - (1979) in Tunneling in Biological Systems (Chance, B., DeVault, D.C., Frauenfelder, H., Marcus, R.A., Schrieffer, J.B. and Sutin, N. eds.), pp. 319-352, Academic Press,
- New York 212 Blankenship, D.E. and Parson, W.W. (1979) in Photo-

Biophys. Acta 503, 524-544

pp. 71-114, Elsevier, New York 213 DeVault, D. and Chance, B. (1966) Biophys. J. 6, 825-847 214 Tiede, D.M., Leigh, J.S. and Dutton, P.L. (1978) Biochim.

synthesis in Relation to Model Systems (Barber, J., ed.),

- 215 Buhks, E., Bixon, M. and Jortner, J. (1981) Chem. Phys. 55, 41-48 216 Jortner, J. (1980) Biochim. Biophys. Acta 594, 193-230
- 217 Deisenhofer, J., Epp, O., Miki, K., Huber, R. and Michel, H. (1984) J. Mol. Biol. 180, 385-398 218 Feher, G., Hoff, A.J., Isaacson, R.A. and Ackerson, L.C.
- (1975) Ann. N.Y. Acad. Sci. 244, 239-259 219 Norris, J.R., Scheer, H. and Katz, J.J. (1975) Ann. N.Y.
- Acad. Sci. 244, 261-280 220 Davis, M.S., Forman, A. Hanson, L.K., Thornber, J.P. and Fajer, J. (1979) J. Phys. Chem. 83, 3325-3332
- 221 Haberkorn, H., Michel-Beyerle, M.E. and Marcus, R.A. (1979) Proc. Natl. Acad. Sci. USA 76, 4185-4188
- 222 Thornber, J.P. and Olson, J.M. (1971) Photochem. Photobiol. 14, 329-341
- 223 Prince, R.C., Leigh, J.S. and Dutton, P.L. (1976) Biochim. Biophys. Acta 440, 622-636

224 Holten, D., Windsor, M.W., Parson, W.W. and Thornber, J.P. (1978) Biochim. Biophys. Acta 501, 112-126

- 225 Kihara, T. and McCray, J.A. (1973) Biochim. Biophys. Acta 292, 297-309 226 Netzel, T.L., Bergkamp, M.A. and Chang, C.-K. (1982) J. Am. Chem. Soc. 104, 1952-1957
- 227 Fujita, I., Fajer, J., Chang, C.-K., Wang, C.-B., Bergkamp, M.A. and Netzel, T.L. (1982) J. Phys. Chem. 86, 3754–3759 228 Netzel, T.L. (1982) in Biological Events Probed by Ultra-
- fast Laser Spectroscopy (Alfano, R.R., ed.), pp. 79-117, Academic Press, New York 229 Mataga, N., Karen, A. and Okada, T. (1984) J. Am. Chem.

Michel, H. (1985) in Workshop on Antennas and Reaction

(1980) Biochim. Biophys. Acta 592, 461-472

Soc. 106, 2442-2443

- 230 Holten, D., Hoganson, C., Windsor, M.W., Schenck, C.C., Parson, W.W., Migus, A., Fork, R.L. and Shank, C.V.

- 231 Zinth, W., Franz, M.A., Nuss, M.C., Kaiser, W. and

- 233 Closs, G.L. and Sitzmann, E.V. (1981) J. Am. Chem. Soc. 103, 3217-3219
- 234 Zankel, K.L., Reed, D.W. and Clayton, R.K. (1968) Proc.
 - Natl. Acad. Sci. U.S.A. 61, 1243-1249
- 235 Dutton, P.I. (1971) Biochim. Biophys. Acta 226, 63-80

FEBS Lett. 89, 205-210

Biophys. 172, 329-334

Biophys. J. 24, 361-369

254, 5388-5396

21, 1843-1846

254 Reference deleted

3392-3398

747-751

Biochem, 16, 4975-4978

Academic Press, New York

Chem. 259, 11001-11009

E. (1975) Biochem. J. 147, 145-153

(1976) Biochem. J. 157, 591-598

Chem. Soc. 106, 5043-5045

- Biophys. Acta 767, 345-361

42. 519-530

239 Reference deleted

Verlag, Heidelberg, in the press

245 Chapman, S.K., Davies, D.M., Vuik, C.P.J. and Sykes, A.G. (1984) J. Am. Chem. Soc. 106, 2692-2696

247 Salemme, F.R. (1977) Annu. Rev. Biochem. 46, 299-329 248 Dailey, H.A. and Strittmatter, P. (1979) J. Biol. Chem.

249 Ng, S., Smith, M.B., Smith, H.T. and Millet, F. (1977)

250 Strittmatter, P. (1964) in Rapid Mixing and Sampling Techniques in Biochemistry (Chance, B., Eisenhardt, R.H.,

251 Mauk, M.R., Reid, L.S. and Mauk, A.G. (1982) Biochem.

252 Argos, P. and Mathews, F.S. (1975) J. Biol. Chem. 250,

253 Reid, L.S., Taniguchi, V.T., Gray, H.B. and Mauk, A.G.

255 Simolo, K.P., McLendon, G.L., Mauk, M.R. and Mauk,

256 Malmström, B. (1979) Biochim. Biophys. Acta 549,

257 Brudvig, G.W., Blair, D.F. and Chan, S.I. (1984) J. Biol.

260 Greenwood, C., Brittain, T., Wilson, M. and Brunori, M.

261 Yonetani, T. and Ray, G.S. (1965) J. Biol. Chem. 240,

258 Andreasson, L.E. (1975) Eur. J. Biochem. 53, 591-597 259 Wilson, M.T., Greenwood, C., Brunori, M. and Antonini,

A.G. (1984) J. Am. Chem. Soc. 106, 5012-5013

(1982) J. Am. Chem. Soc. 104, 7516-7519

Gibson, Q.H. and Lonberg-Holm, K.K., eds.), pp. 71-84,

246 Salemme, F.R. (1976) J. Molec. Biol. 102, 563-568

Centers of Photosynthetic Bacteria-Structure, Interactions and Dynamics (Michel-Beyerle, M.E., ed.), Springer-

232 Shuvalov, V.A. and Klevanik, A.V. (1983) FEBS Lett. 160,

- 236 Woodbury, N.W.T. and Parson, W.W. (1984) Biochim.
- 237 Klimov, V.V., Shuvalov, V.A., Krakhmaleva, I.N.,

 - Klevanik, A.V. and Krasnovsky, A.A. (1977) Biokhimia
- 238 Pacham, N.K., Berriman, J.A. and Jackson, J.B. (1978)
- 240 Sutin, N. (1982) Accounts Chem. Res. 15, 275-282
- 241 Prince, R.C. and Dutton, P.L. (1976) Arch. Biochem.
- 242 Gunner, M.R., Liang, Y., Nagus, D.K., Hochstrasser, R.M. and Dutton, P.L. (1982) Biophys. J. 37, 226A
- 243 Pellin, M.J., Wraight, C.A. and Kaufman, K.J. (1978)

289 Kang, C.H., Brautigan, D.L., Osheroff, N. and Margoliash,

290 Purcell, W.L. and Erman, J.E. (1976) J. Am. Chem. Soc.

291 Yandell, J.K. and Yonetani, T. (1983) Biochim. Biophys.

292 Yonetani, T. (1976) in The Enzymes (Boyer, P.D., ed.), Vol. 13, pp. 345-361, Academic Press, New York

293 Ho, P.S., Hoffman, B.M., Kang, C.H. and Margoliash, E.

294 Poulos, T.L. and Kraut, J. (1980) J. Biol. Chem. 255,

295 Fujita, I., Hanson, L.K., Walker, F.A. and Fajer, J. (1983)

296 Poulos, T.L., Freer, S.T., Alden, R.A., Edwards, S.L.,

297 Poulos, T.L. and Kraut, J. (1980) J. Biol. Chem. 255,

299 Finzel, B.C., Poulos, T.L. and Kraut, J. (1984) J. Biol.

298 Kraut, J. (1981) Biochem. Soc. Trans. 9, 197-202

300 Marcus, R.A. (1960) Disc. Faraday Soc. 29, 129-130

301 Noyes, R.M. (1961) Prog. Reaction Kinetics 1, 129-160 302 Debye, P. (1942) Trans. Amer. Electrochem. Soc. 82,

303 Sutin, N. and Brunschwig, B.S. (1982) Am. Chem. Soc.

304 Marcus, R.A. (1981) Int. J. Chem. Kinetics 13, 865-872

306 Jortner, J. (1960) J. Am. Chem. Soc. 102, 6676-6686

305 Dogonadze, R.R. (1971) in Reactions of Molecules at

307 Beratan, D.N. and Hopfield, J.J. (1984) J. Chem. Phys. 81,

308 Marcus, R.A. (1977) in Special Topics in Electrochemistry.

(Rock, P.A., ed.), pp. 161-179, Elsevier, New York

310 Kuznetsov, A.M. and Ulstrup, J. (1981) J. Chem. Phys. 75,

313 Halpern, J. and Orgel, L.E. (1960) Disc. Faraday Soc. 29,

314 Kosower, E.M. and Huppert, D. (1983) Chem. Phys. Letter.

315 Ovchinnikova, M.Ya. (1981) Teor. Eksp. Khim. 17,

309 McConnell, H.M. (1961) J. Chem. Phys. 35, 508-515

311 Davydov, A.S. (1978) Phys. Stat. Sol. B. 90, 457-464

312 Larsson, S. (1984) J. Phys. Chem. 88, 1321-1323

651-654 (Theor. Exper. Chem. 17, 507-510) 316 Calef, D.F. and Wolynes, P.G. (1983) J. Phys. Chem. 87,

Electrodes (Hush, N.S., ed.), Wiley, New York, Ch. 3,

Skoglund, U., Takio, K., Eriksson, B., Xuong, N.-H., Yonetani, T. and Kraut, J. (1980) J. Biol. Chem. 255,

E. (1978) J. Biol. Chem. 253, 6502-6510

(1983) J. Biol. Chem. 258, 4356-4363

J. Am. Chem. Soc. 105, 3296-3300

98, 7033-7037

8199-8205

575-580

265-271

Eqn. 201

5753-5759

2047-2055

96, 433-435

3387-3400

381-397

32 - 41

10322-10330

Chem. 259, 13027-13036

Symp. Ser. 198, 105--125

Acta 748, 263-270

- Dockter, M.E., Steinemann, A. and Schatz, G. (1978) J. Biol. Chem. 253, 311-317 Farver, O. and Pecht, I. (1984) in Copper Proteins and
- Copper Enzymes (Lontie, R., ed.), pp. 183-215, CRC
- Press, Boca Raton, FL Colman, P.M., Freeman, H.C., Guss, J.M., Murata, M.,
- Norris, V.A., Ramshaw, J.A.M. and Venkatappa, M.P. (1980) Nature 272, 319-324 Adman, E.T., Stenkamp, R.E., Sieker, L.C. and Jensen,
- L.H. (1978) Mol. Biol. 123, 35-47 Solomon, E.I., Hare, J.W., Dooley, D.M., Dawson, J.H.,
- Stephens, P.J. and Gray, H.B. (1980) J. Am. Chem. Soc. 102, 168-178 Cookson, D.J., Hayes, M.T. and Wright, P.E. (1980) Bio-
- chem. Biophys. Acta 591, 162-176
- Lappin, A.G., Segal, M.G., Weatherburn, D.C. and Sykes, A.G. (1979) J. Chem. Soc. Chem. Comm. 38-39
- Handford, P.M., Hill, H.A.O., Lee, R.W.-K., Henderson,
- R.A. and Sykes, A.G. (1980) J. Inorg. Biochem. 13, 83-88 Wherland, S., Holwerda, R.A., Rosenberg, R.C. and Gray,
- H.B. (1975) J. Am. Chem. Soc. 97, 5260-5262 Chapman, S.K., Knox, C.V., Kathirgamathan, P. and
- Sykes, A.G. (1984) J. Chem. Soc. Dalton Trans. 2769-2773 Lappin, A.G. (1981) in Metal Ions in Biological Systems, (Sigel, H. ed.), Vol. 13, pp. 15-71, Marcel Dekker, New
- York Augustin, M.A., Chapman, S.K., Davies, D.M., Watson, A.D. and Sykes, A.G. (1984) J. Inorg. Biochem. 20, 281 - 289
- Sisley, M.J., Segal, M.G., Stanley, C.S., Adzamli, I.K. and Sykes, A.G. (1983) J. Am. Chem. Soc. 105, 222-228 Taniguchi, V.T., Sailasuta-Scott, N., Anson, F.C. and Gray, H.B. (1980) Pure and Appl. Chem. 52, 2275-2281
- Canters, G.W., Hill, H.A.O., Kitchen, N.A. and Adman, E.T. (1984) J. Mag. Res. 57, 1-23 Beattie, J.K., Fensom, D.J., Freeman, H.C., Woodcock, E.,
- Hill, H.A.O. and Stokes, A.M. (1975) Biochim. Biophys. Acta 405, 109-114 Dahlin, S., Reinhammar, B. and Wilson, M.T., (1984)
- Biochem. J. 218, 609-614 Farver, O. and Pecht, I. (1981) in Copper Proteins (Spiro, T.G., ed.), pp. 151-192, Wiley, New York
- Goldberg, M. and Pecht, I. (1976) Biochem. 15, 4197-4208 Mauk, A.G., Scott, R.A. and Gray, H.B. (1980) J. Am. Chem. Soc. 102, 4360-4363 Brunswig, B.S., De Laire, P.J., English, A.M., Goldberg,

Wherland, S. and Pecht, I. (1978) Biochem. 17, 2585-2591

Norris, G.E., Anderson, B.F. and Baker, E.N. (1983) J.

Moore, G.R. and Williams, R.J.P. (1976) Coord. Chem.

Yonetani, T. and Ray, G.S. (1966) J. Biol. Chem. 241,

the press

700-706

Mol. Biol. 165, 501-521

J. Biol. Chem. 252, 919-926

Rev. 18, 125-197

- M., Gray, H.B. and Sutin, N. (1985) Inorg. Chem. 24, in
- Wilson, M.T., Silvestrini, M.C., Morpurgo, L. and Brunori, M. (1979) J. Inorg. Biochem. 11, 95-100
- 317 Reference deleted
- 318 Kakitani, T. and Mataga, N. (1985) Chem. Phys. 93,

- 319 Dietz, R. and Peover, M.E. (1968) Discussions Faraday

- Soc. 45, 154–166

- 320 Kojima, H. and Bard, A.J. (1975) J. Am. Chem. Soc. 97,

- 6317-6324

- Kang, C.H., Ferguson-Miller, S. and Margoliash, E. (1978)

- 321 Kojima, H. and Bard, A.J. (1975) J. Electroanal. Chem. 63, 117-129
- 322 Peover, M.E. and Powell, J.S. (1969) J. Electroanal. Chem. 20, 427-433
- 323 Huebert, B.J. and Smith, D.E. (1971) J. Electroanal. Chem. 31, 333-348
- 324 Garreau, D., Saveant, J.E. and Tessier, D. (1979) J. Phys. Chem. 83, 3003-3007
- 325 Fawcett, W.R. and Lasia, A. (1978) J. Phys. Chem. 82, 1114-1121
- 326 Rehm, D. and Weller, A. (1969) Ber. Bunsenges. Phys. Chem. 73, 834-839

- 327 Markham, J.J. (1959) Rev. Mod. Phys. 31, 956-989328 Ballhausen, C. (1979) Molecular Electronic Structures of
 - Transition Metal Complexes, pp. 112–115, McGraw-Hill, Maidenhead
- 329 Sarai, A. (1980) Biochim. Biophys. Acta 589, 71-83
- 330 Kakitani, T. and Kakitani, H. (1981) Biochim. Biophys. Acta 635, 498-514
- 331 Marcus, R.A. and Sutin, N. (1985) in Workshop on Antennas and Reaction Centers of Photosynthetic Bacteria-Structure, Interactions and Dynamics (Michel-Beyerle, M.E., ed.), Springer-Verlag, Heidelberg, in the press