



My first encounters with McGill University came when I was still in a baby carriage. My mother used to wheel me about the campus when we lived in that neighborhood and, as she recounted years later, she would tell me that I would go to McGill. There was some precedent for my going there, since two of my father's brothers received their M.D.'s at McGill.

I have always loved going to school. Since neither of my parents had a higher education, my academic "idols" were these two paternal uncles and one of their uncles, my great-uncle, Henrik Steen (né Markus). My admiration for him, living in faraway Sweden, was not because of a teol.dr. (which he received from the University of Uppsala in 1915) nor because of the many books he wrote—I knew nothing of that—but rather because he was reputed to speak 13 languages. I learned decades later that the number was only 9! Growing up, mostly in Montreal, I was an only child of loving parents. I admired my father's athletic prowess—he excelled in several sports—and my mother's expressive singing and piano playing.

My interest in the sciences started with mathematics in the very beginning, and later with chemistry in early high school and the proverbial home chemistry set. My education at Baron Byng High School was excellent, with dedicated masters (boys and girls were separate). I spent the next years at McGill University, for both undergraduate and, as was the custom of the time, graduate study. Our graduate supervisor, Carl A. Winkler, specialized in rates of chemical reactions. He himself had received his Ph.D. as a student of Cyril Hinshelwood at Oxford. Hinshelwood was later the recipient of the Nobel Prize for his work on chemical kinetics. Winkler brought to his laboratory an enthusiastic joyousness in research and was much loved by his students.

During my McGill years, I took a number of math courses, more than other students in chemistry. Upon receiving a Ph.D. from McGill University in 1946, I joined the new post-doctoral program at the National Research Council of Canada in Ottawa. This program at NRC later became famous, but at the time it was still in its infancy and our titles were Junior Research Officers. The photochemistry group was headed by E.W.R. Steacie, an international figure in the study of free-radical reactions and a major force in the development of the basic research program at NRC. I benefitted from the quality of his research on gas phase reaction rates. Like my research on chemical reaction rates in solution at McGill (kinetics of nitration), it was experimental in nature. There were no theoretical chemists in Canada at the time, and as students I don't think we ever considered how or where theories were conceived.

About 1948 a fellow post-doctoral at NRC, Walter Trost, and I formed a two-man seminar to study theoretical papers related to our experimental work. This adventure led me to explore the possibility of going on a second post-doctoral, but in theoretical work, which seemed like a radical step at the time. I had a tendency to break the glass vacuum apparatus, due to a still present impetuous haste, with time-consuming consequences. Nevertheless, the realization that breaking a pencil point would have far less disastrous consequences played little or no role, I believe, in this decision to explore theory!

I applied in 1948 to six well-known theoreticians in the U.S. for a postdoctoral research fellowship. The possibility that one of them might take on an untested applicant, an applicant hardly qualified for theoretical research, was probably too much to hope for. Oscar K. Rice at the University of North Carolina alone responded favorably, subject to the success of an application he would make to the Office of Naval Research for this purpose. It was, and in February 1949 I took the train south, heading for the University of North Carolina in Chapel Hill. I was impressed on arrival there by the red clay, the sandy walks, and the graciousness of the people.

After that, I never looked back. Being exposed to theory, stimulated by a basic love of concepts and mathematics, was a marvelous experience. During the first three months I read everything I could lay my hands on regarding reaction rate theory, including Marcelin's classic 1915 theory which came within one small step of the Transition State Theory of 1935. I read numerous theoretical papers in German, a primary language for the "chemical dynamics" field in the 1920s and 1930s, attended my first formal course in quantum mechanics, given by Nathan Rosen in the Physics Department, and was guided by Oscar in a two-man weekly seminar in which I described a paper I had read and he pointed out assumptions in it that I had overlooked. My life as a working theorist began three months after this preliminary study and background reading, when Oscar gently nudged me toward working on a particular problem.

Fortunately for me, Oscar's gamble paid off. Some three months later, I had formulated a particular case of what was later entitled by B. Seymour Rabinovitch, RRKM theory ("Rice-Ramsperger-Kassel-Marcus"). In it, I blended statistical ideas from the RRK theory of the 1920s with those of the transition state theory of the mid-1930s. The work was published in 1951. In 1952 I wrote the generalization of it for other reactions. In addition, six months after arrival in Chapel Hill, I was also blessed by marriage to Laura Hearne, an attractive graduate student in sociology at UNC. She is here with me at this ceremony. Our three sons, Alan, Kenneth and Raymond, and two daughters-in-law are also present today.

In 1951, I attempted to secure a faculty position. This effort met with little success (35 letters did not yield 35 no's, since not everyone replied!). Very fortunately, that spring I met Dean Raymond Kirk of the Polytechnic Institute of Brooklyn at an American Chemical Society meeting in Cleveland, which I was attending primarily to seek a faculty position. This

meeting with Dean Kirk, so vital for my subsequent career, was arranged by Seymour Yolles, a graduate student at UNC in a course I taught during Rice's illness. Seymour had been a student at Brooklyn Poly and learned, upon accidentally encountering Dr. Kirk, that Kirk was seeking new faculty. After a subsequent interview at Brooklyn Poly, I was hired, and life as a fully independent researcher began.

I undertook an experimental research program on both gas phase and solution reaction rates, wrote the 1952 RRKM papers, and wondered what to do next in theoretical research. I felt at the time that it was pointless to continue with RRKM since few experimental data were available. Some of our experiments were intended to produce more.

After some minor pieces of theoretical study that I worked on, a student in my statistical mechanics class brought to my attention a problem in polyelectrolytes. Reading everything I could about electrostatics, I wrote two papers on that topic in 1954/55. This electrostatics background made me fully ready in 1955 to treat a problem I had just read about on electron transfers. I comment on this next period on electron transfer research in my Nobel Lecture. About 1960, it became clear that it was best for me to bring the experimental part of my research program to a close—there was too much to do on the theoretical aspects—and I began the process of winding down the experiments. I spent a year and a half during 1960–61 at the Courant Mathematical Institute at New York University, auditing many courses which were, in part, beyond me, but which were, nevertheless, highly instructive.

In 1964, I joined the faculty of the University of Illinois in Urbana-Champaign and I never undertook any further experiments there. At Illinois, my interests in electron transfer continued, together with interests in other aspects of reaction dynamics, including designing “natural collision coordinates”, learning about action-angle variables, introducing the latter into molecular collisions, reaction dynamics, and later into semiclassical theories of collisions and of bound states, and spending much of my free time in the astronomy library learning more about classical mechanics, celestial mechanics, quasiperiodic motion, and chaos. I spent the academic year of 1975–76 in Europe, first as Visiting Professor at the University of Oxford and later as a Humboldt Awardee at the Technical University of Munich, where I was first exposed to the problem of electron transfer in photosynthesis.

In 1978, I accepted an offer from the California Institute of Technology to come there as the Arthur Amos Noyes Professor of Chemistry. My semiclassical interlude of 1970–80 was intellectually a very stimulating one, but it involved for me less interaction with experiments than had my earlier work on unimolecular reaction rates or on electron transfers. Accordingly, prompted by the extensive experimental work of my colleagues at Caltech in these fields of unimolecular reactions, intramolecular dynamics and of electron transfer processes, as well as by the rapidly growing experimental work in both broad areas world-wide, I turned once again to those particu-

lar topics and to the many new types of studies that were being made. Their scope and challenge continues to grow to this day in both fields. Life would be indeed easier if the experimentalists would only pause for a little while!

There was a time when I had wondered about how much time and energy had been lost doing experiments during most of my stay at Brooklyn Poly—experiments on gas phase reactions, flash photolysis, isotopic exchange electron transfer, bipolar electrolytes, nitration, and photoelectrochemistry, among others—and during all of my stay at NRC and at McGill. In retrospect, I realized that this experimental background heavily flavored my attitude and interests in theoretical research. In the latter I drew, in most but not all cases, upon experimental findings or puzzles for theoretical problems to study. The growth of experiments in these fields has served as a continually rejuvenating influence. This interaction of experiment and theory, each stimulating the other, has been and continues to be one of the joys of my experience.

Honors received for the theoretical work include the Irving Langmuir and the Peter Debye Awards of the American Chemical Society (1978, 1988), the Willard Gibbs, Theodore William Richards, and Pauling Medals, and the Remsen and Edgar Fahs Smith Awards, from various sections of the ACS, (1988, 1990, 1991, 1991, 1991), the Robinson and the Centenary Medals of the Faraday Division of the Royal Society of Chemistry (1982, 1988), Columbia University's Chandler Medal (1983) and Ohio State's William Lloyd Evans Award (1990), a Professorial Fellowship at University College, Oxford (1975 to 1976) and a Visiting Professorship in Theoretical Chemistry at Oxford during that period, the Wolf Prize in Chemistry (1985), the National Medal of Science (1989), the Hirschfelder Prize in Chemistry (1993), election to the National Academy of Sciences (1970), the American Academy of Arts and Sciences (1973), the American Philosophical Society (1990), honorary membership in the Royal Society of Chemistry (1991), and foreign membership in the Royal Society (London) (1987) and in the Royal Society of Canada (1993). Honorary degrees were conferred by the University of Chicago and by Goteborg, Polytechnic, McGill, and Queen's Universities and by the University of New Brunswick (1983, 1986, 1987, 1988, 1993, 1993). A commemorative issue of the *Journal of Physical Chemistry* was published in 1986.

ELECTRON TRANSFER REACTIONS IN CHEMISTRY: THEORY AND EXPERIMENT

Nobel Lecture, December 8, 1992

by

RUDOLPH A. MARCUS

Noyes Laboratory of Chemical Physics, California Institute of Technology,
MS 127-72, Pasadena, CA 91125, USA

ELECTRON TRANSFER EXPERIMENTS SINCE THE LATE 1940s

Since the late 1940s, the field of electron transfer processes has grown enormously, both in chemistry and biology. The development of the field, experimentally and theoretically, as well as its relation to the study of other kinds of chemical reactions, represents to us an intriguing history, one in which many threads have been brought together. In this lecture, some history, recent trends, and my own involvement in this research are described.

The early experiments in the electron transfer field were on “isotopic exchange reactions” (self-exchange reactions) and, later, “cross reactions.” These experiments reflected two principal influences. One of these was the availability after the Second World War of many radioactive isotopes, which permitted the study of a large number of isotopic exchange electron transfer reactions, such as



and



in aqueous solution, where the asterisk denotes a radioactive isotope.

There is a two-fold simplicity in typical self-exchange electron transfer reactions (so-called since other methods beside isotopic exchange were later used to study some of them): (1) the reaction products are identical with the reactants, thus eliminating one factor which usually influences the rate of a chemical reaction in a major way, namely the relative thermodynamic stability of the reactants and products; and (2) no chemical bonds are broken or formed in *simple* electron transfer reactions. Indeed, these self-exchange reactions represent, for these combined reasons, the simplest class of reactions in chemistry. Observations stemming directly from this simplicity were to have major consequences, not only for the electron

transfer field but also, to a lesser extent, for the study of other kinds of chemical reactions as well (cf Shaik *et al*, ref. 2).

A second factor in the growth of the electron transfer field was the introduction of new instrumentation, which permitted the study of the rates of rapid chemical reactions. Electron transfers are frequently rather fast, compared with many reactions which undergo, instead, a breaking of chemical bonds and a forming of new ones. Accordingly, the study of a large body of fast electron transfer reactions became accessible with the introduction of this instrumentation. One example of the latter was the stopped-flow apparatus, pioneered for inorganic electron transfer reactions by N. Sutin. It permitted the study of bimolecular reactions in solution in the millisecond time scale (a fast time scale at the time). Such studies led to the investigation of what has been termed electron transfer “cross reactions,” i.e., electron transfer reactions between two different redox systems, as in



which supplemented the earlier studies of the self-exchange electron transfer reactions. A comparative study of these two types of reaction, self-exchange and cross-reactions, stimulated by theory, was also later to have major consequences for the field and, indeed, for other areas.

Again, in the field of electrochemistry, the new post-war instrumentation in chemical laboratories led to methods which permitted the study of fast electron transfer reactions at metal electrodes. Prior to the late 1940s only relatively slow electrochemical reactions, such as the discharge of an H_3O^+ ion at an electrode to form H_2 , had been investigated extensively. They involved the breaking of chemical bonds and the forming of new ones.

Numerous electron transfer studies have now also been made in other areas, some depicted in Figure 1. Some of these investigations were made possible by a newer technology, lasers particularly, and now include studies in the picosecond and subpicosecond time regimes. Just recently, (non-laser) nanometer-sized electrodes have been introduced to study electrochemical processes that are still faster than those hitherto investigated. Still other recent investigations, important for testing aspects of the electron transfer theory at electrodes, involve the new use of an intervening ordered adsorbed monolayer of long chain organic compounds on the electrode to facilitate the study of various effects, such as varying the metal-solution potential difference on the electrochemical electron transfer rate.

In some studies of electron transfer reactions in solution there has also been a skillful blending of these measurements of chemical reaction rates with various organic or inorganic synthetic methods, as well as with site-directed mutagenesis, to obtain still further hitherto unavailable information. The use of chemically modified proteins to study the distance dependence of electron transfer, notably by Gray and coworkers, has opened a whole new field of activity.

The interaction of theory and experiment in these many electron transfer fields has been particularly extensive and exciting, and each has stimulated

Developments in Electron Transfer Reactions




Figure 1. Examples of topics in the electron transfer field (Marcus and Siddarth, ref. 2).

the other. The present lecture addresses the underlying theory and this interaction.

THE EARLY EXPERIENCE

My own involvement in the electron transfer field began in a rather circuitous way. In an accompanying biographical note I have commented on my earlier background, which was in experimental measurements of reaction rates as a chemistry graduate student at McGill University (1943–46) and as a post-doctoral associate at the National Research Council of Canada (1946–49). A subsequent post-doctoral study at the University of North Carolina (1949–51) on the theory of reaction rates resulted in what is now known in the literature as RRKM theory (Rice, Ramsperger, Kassel, Marcus).

This unimolecular reaction field reflects another long and extensive interaction between theory and experiment. RRKM theory enjoys widespread use and is now usually referred to in the literature only by its

acronym (or by the texts written about it, ref. 4), instead of by citation of the original articles.

After the theoretical post-doctoral, I joined the faculty of the Polytechnic Institute of Brooklyn in 1951 and wondered what theoretical research to do next after writing the RRKM papers (1951 – 52). I remember vividly how a friend of mine, a colleague at Brooklyn Poly, Frank Collins, came down to my office every day with a new idea on the liquid state transport theory which he was developing, while I, for theoretical research, had none. Perhaps this gap in not doing anything immediately in the field of theory was, in retrospect, fortunate: In not continuing with the study of the theory of unimolecular reactions, for which there were too few legitimate experimental data at the time to make the subject one of continued interest, I was open for investigating quite different problems in other areas. I did, however, begin a program of experimental studies in gas phase reactions, prompted by my earlier studies at NRC and by the RRKM work.

In the biographical note I have also recalled how a student in my statistical mechanics class in this period (Abe Kotliar) asked me about a particular problem in polyelectrolytes. It led to my writing two papers on the subject (1954 – 55), one of which required a considerable expansion in my background in electrostatics, so as to analyze different methods for calculating the free energy of these systems: In polyelectrolyte molecules, it may be recalled, the ionic charges along the organic or inorganic molecular backbone interact with each other and with the solvent. In the process, I read the relevant parts of the texts that were readily available to me on electrostatics (Caltech's Mason and Weaver's was later to be particularly helpful!). When shortly thereafter I encountered some papers on electron transfer, a field entirely new to me, I was reasonably well prepared for treating the problems which lay ahead.

DEVELOPING AN ELECTRON TRANSFER THEORY

Introduction

My first contact with electron transfers came in 1955 as a result of chancing upon a 1952 symposium issue on the subject in the *Journal of Physical Chemistry*. An article by Bill Libby caught my eye—a use of the Franck-Condon principle to explain some experimental results, namely, why some isotopic exchange reactions which involve electron transfer between pairs of small cations in aqueous solution, such as reaction (1), are relatively slow, whereas electron transfers involving larger ions, such as $\text{Fe}(\text{CN})_6^{3-} \rightarrow \text{Fe}(\text{CN})_6^{4-}$ and $\text{MnO}_4^- \rightarrow \text{MnO}_4^{2-}$, are relatively fast.

Libby explained this observation in terms of the Franck-Condon principle, as discussed below. The principle was used extensively in the field of spectroscopy for interpreting spectra for the excitation of the molecular electronic-vibrational quantum states. An application of that principle to chemical reaction rates was novel and caught my attention. In that paper Libby gave a “back-of-the-envelope” calculation of the resulting solvation

energy barrier which slowed the reaction. However, I felt instinctively that even though the idea—that somehow the Franck—Condon principle was involved—seemed strikingly right, the calculation itself was incorrect. The next month of study of the problem was, for me, an especially busy one. To place the topic in some perspective I first digress and describe the type of theory that was used for other types of chemical reaction rates at the time and continues to be useful today.

Reaction rate theory

Chemical reactions are often described in terms of the motion of the atoms of the reactants on a potential energy surface. This potential energy surface is really the electronic energy of the entire system, plotted versus the positions of all the atoms. A very common example is the transfer of an atom or a group B from AB to form BC



An example of reaction (4) is the transfer of an H, such as in $IH + Br \rightarrow I + HBr$, or the transfer of a CH_3 group from one aromatic sulfonate to another. To aid in visualizing the motion of the atoms in this reaction, this potential energy function is frequently plotted as constant energy contours in a space whose axes are chosen to be two important relative coordinates such as, in reaction (4), a scaled AB bond length and a scaled distance from the center of mass of AB to C, as in Figure 2.

Potential Energy Contours for an Atom or Group Transfer




Figure 2. Potential energy contours for reaction (4), $AB + C \rightarrow A + BC$, in the collinear case.

A point representing this reacting system begins its trajectory in the lower right region of the figure in a valley in this plot of contours, the “valley of the reactants.” When the system has enough energy, appropriately distributed between the various motions, it can cross the “mountain pass” (saddle-point region) separating the initial valley from the products’ valley in the upper left, and so form the reaction products. There is a line in the figure, XY, analogous to the “continental divide” in the Rocky Mountains in the U.S., which separates systems which could spontaneously flow into the reactants’ valley from those which could flow into the products’ one. In chemists’ terminology this line represents the “transition state” of the reaction.

In transition state theory a quasi-equilibrium between the transition state and the reactants is frequently postulated, and the reaction rate is then calculated using equilibrium statistical mechanics. A fundamental dynamical basis, which replaces this apparently *ad hoc* but common assumption of transition state theory and which is perhaps not as well known in the chemical literature as it deserves to be, was given many years ago by the physicist and one-time chemical engineer, Eugene Wigner (1938). He used a classical mechanical description of the reacting system in the many-dimensional space (of coordinates and momenta). Wigner pointed out that the quasi-equilibrium would follow as a dynamical consequence if each trajectory of a moving point representing the reacting system in this many-dimensional space did not recross the transition state (and if the distribution of the reactants in the reactants’ region were a Boltzmann one). In recent times, the examination of this recrossing has been a common one in classical mechanical trajectory studies of chemical reactions. Usually, recrossings are relatively minor, except in nonadiabatic reactions, where they are readily treated (cf discussion, later).

In practice, transition state theory is generalized so as to include as many coordinates as are needed to describe the reacting system. Further, when the system can “tunnel” quantum mechanically through the potential energy barrier (the “pass”) separating the two valleys, as for example frequently happens at low energies in H-transfer reactions, the method of treating the passage across the transition state region needs, and has received, refinement. (The principal problem encountered here has been the lack of “dynamical separability” of the various motions in the transition state region.)

Electron transfer theory. Formulation

In contrast to the above picture, we have already noted that in simple electron transfer reactions no chemical bonds are broken or formed and so a somewhat different picture of the reaction is needed for the electron transfer reaction.

In his 1952 symposium paper, Libby noted that when an electron is transferred from one reacting ion or molecule to another, the two new molecules or ions formed are in the wrong environment of the solvent

molecules, since the nuclei do not have time to move during the rapid electron jump: in reaction (1) a Fe^{2+} ion would be formed in some configuration of the many nearby dipolar solvent molecules that was appropriate to the original Fe^{3+} ion. Analogous remarks apply to the newly formed Fe^{3+} ion in the reaction. On the other hand, in reactions of "complex ions," such as those in the $\text{Fe}(\text{CN})_6^{-3} - \text{Fe}(\text{CN})_6^{-4}$ and $\text{MnO}_4^- - \text{MnO}_4^{2-}$ self-exchange reactions, the two reactants are larger, and so the change of electric field in the vicinity of each ion, upon electron transfer, would be smaller. The original solvent environment would therefore be less foreign to the newly formed charges, and so the energy barrier to reaction would be less. In this way Libby explained the faster self-exchange electron transfer rate for these complex ions. Further confirmation was noted in the ensuing discussion in the symposium: the self-exchange $\text{Co}(\text{NH}_3)_6^{3+} - \text{Co}(\text{NH}_3)_6^{2+}$ reaction is very slow, and it was pointed out that there was a large difference in the equilibrium Co-N bond lengths in the $3+$ and the $2+$ ions, and so each ion would be formed in a very "foreign" configuration of the vibrational coordinates, even though the ions are "complex ions."

After studying Libby's paper and the symposium discussion, I realized that what troubled me in this picture for reactions occurring in the dark was that energy was not conserved: the ions would be formed in the wrong high-energy environment, but the only way such a non-energy conserving event could happen would be by the absorption of light (a "vertical transition"), and not in the dark. Libby had perceptively introduced the Franck-Condon principle to chemical reactions, but something was missing.

In the present discussion, as well as in Libby's treatment, it was supposed that the electronic interaction of the reactants which causes the electron transfer is relatively weak. That view is still the one that seems appropriate today for most of these reactions. In this case of weak-electronic interaction, the question becomes: how does the reacting system behave in the dark so as to satisfy both the Franck-Condon principle and energy conservation? I realized that fluctuations had to occur in the various nuclear coordinates, such as in the orientation coordinates of the individual solvent molecules and indeed in any other coordinates whose most probable distribution for the products differs from that of the reactants. With such fluctuations, values of the coordinates could be reached which satisfy both the Franck-Condon and energy conservation conditions and so permit the electron transfer to occur in the dark.

For a reaction such as reaction (1), an example of an initial and final configuration of the solvent molecules is depicted in Figure 3. Fluctuations from the original equilibrium ensemble of configurations were ultimately needed, prior to the electron transfer, and were followed by a relaxation to the equilibrium ensemble for the products, after electron transfer.

The theory then proceeded as follows. The potential energy U_r of the entire system, reactants plus solvent, is a function of the many hundreds of relevant coordinates of the system, coordinates which include, among others, the position and orientation of the individual solvent molecules (and

Electron Transfer in Solution




Figure 3. Typical nuclear configurations for reactants, products, and surrounding solvent molecules in reaction (1). The longer M-OH₂ bond length in the +2 state is indicated schematically by the larger ionic radius. (Sutin, ref. 2)

hence of their dipole moments, for example), and the vibrational coordinates of the reactants, particularly those in any inner coordination shell of the reacting ions. (E.g., the inner coordination shell of an ion such as Fe²⁺ or Fe³⁺ in water is known from EXAFS experiments to contain six water molecules.) No longer were there just the two or so important coordinates that were dominant in reaction (4).

Similarly, after the electron transfer, the reacting molecules have the ionic charges appropriate to the reaction products, and so the relevant potential energy function U_p is that for the products plus solvent. These two potential energy surfaces will intersect if the electronic coupling which leads to electron transfer is neglected. For a system with N coordinates this intersection occurs on an $(N-1)$ dimensional surface, which then constitutes in our approximation the transition state of the reaction. The neglected electronic coupling causes a well-known splitting of the two surfaces in the vicinity of their intersection. A schematic profile of the two potential energy surfaces in the N -dimensional space is given in Figure 4. (The splitting is not shown.)

Due to the effect of the previously neglected electronic coupling and the coupling between the electronic motion and the nuclear motion near the intersection surface S , an electron transfer can occur at S . In classical terms, the transfer at S occurs at fixed positions and momenta of the atoms, and so the Franck-Condon principle is satisfied. Since U_r equals U_p at S , energy is also conserved. The details of the electron transfer depend on the extent of electronic coupling and how rapidly the point representing the system in this N -dimensional space crosses S . (It has been treated, for example, using as an approximation the well-known one-dimensional Landau-Zener expression for the transition probability at the near-intersection of two potential energy curves.)

Potential Energy Surfaces, Profile




Figure 4. Profile of potential energy surfaces for reactants plus environment, R , and for products plus environment, P . Solid curves: schematic. Dashed curves: schematic but slightly more realistic. The typical splitting at the intersection of U_r and U_p is not shown in the Figure (Marcus and Siddarth, ref. 2).

When the splitting caused by the electronic coupling between the electron donor and acceptor is large enough at the intersection, a system crossing S from the lower surface on the reactants' side of S continues onto the lower surface on the products' side, and so an electron transfer in the dark has then occurred. When the coupling is, instead, very weak, ("non-adiabatic reactions") the probability of successfully reaching the lower surface on the products' side is small and can be calculated using quantum mechanical perturbation theory, for example, using Fermi's "Golden Rule," an improvement over the 1-dimensional Landau-Zener treatment.

Thus, there is some difference and some similarity with a more conventional type of reaction such as reaction (4), whose potential energy contour plots were depicted in Figure 2. In both cases, fluctuations of coordinates are needed to reach the transition state, but since so many coordinates can now play a significant role in the electron transfer reaction, because of the major and relatively abrupt change in charge distribution on passing through the transition state region, a rather different approach from the conventional one was needed to formulate the details of the theory.

In the initial paper (1956) I formulated the above picture of the mechanism of electron transfer and, to make the calculation of the reaction rate tractable, treated the solvent as a dielectric continuum. In the transition state the position-dependent dielectric polarization $\mathbf{P}_u(\mathbf{r})$ of the solvent, due to the orientation and vibrations of the solvent molecules, was not the one in equilibrium with the reactants' or the products' ionic charges. It represented instead, some macroscopic fluctuation from them. The electronic polarization for the solvent molecules, on the other hand, can rapidly respond to any such fluctuations and so is that which is dictated by the reactants' charges and by the instantaneous $\mathbf{P}_u(\mathbf{r})$.

With these ideas as a basis, what was then needed was a method of calculating the electrostatic free energy G of this system with its still unknown polarization function $\mathbf{P}_u(\mathbf{r})$. I obtained this free energy G by finding a reversible path for reaching this state of the system. Upon then minimizing G , subject to the constraint imposed by the Franck-Condon principle (reflected in the electron transfer occurring at the intersection of the two potential energy surfaces), I was able to find the unknown $\mathbf{P}_u(\mathbf{r})$ and, hence, to find the G for the transition state. That G was then introduced into transition state theory and the reaction rate calculated.

In this research I also read and was influenced by a lovely paper by Platzmann and Franck (1952) on the optical absorption spectra of halide ions in water and later by work of physicists such as Pekar and Frohlich (1954) on the closely related topic of polaron theory. As best as I can recall now, my first expressions for G during this month of intense activity seemed rather clumsy, but then with some rearrangement a simple expression emerged that had the right "feel" to it and that I was also able to obtain by a somewhat independent argument. The expression also reduced reassuringly to the usual one, when the constraint of arbitrary $\mathbf{P}_u(\mathbf{r})$ was removed. Obtaining the result for the mechanism and rate of electron transfer was indeed one of the most thrilling moments of my scientific life.

The expression for the rate constant k of the reaction is given by

$$k = A \exp \left(\frac{-\Delta G^*}{k_B T} \right), \quad (5a)$$

where ΔG^* , in turn, is given by

$$\Delta G^* = \frac{\lambda}{4} \left(1 + \frac{\Delta G^0}{\lambda} \right)^2. \quad (5b)$$

The A in Eq. (5a) is a term depending on the nature of the electron transfer reaction (e.g., bimolecular or intramolecular), ΔG^0 is the standard free energy of reaction (and equals zero for a self-exchange reaction), λ is a "reorganization term," composed of solvational (λ_o) and vibrational (λ_i) components.

$$\lambda = \lambda_o + \lambda_i \quad (6)$$

In a two-sphere model of the reactants, λ_o was expressed in terms of the two ionic radii a_1 and a_2 (including in the radius any inner coordination shell), the center-to-center separation distance R of the reactants, the optical (D_{op}) and static (D_s) dielectric constants of the solvent, and the charge transferred Δe from one reactant to the other:

$$\lambda_o = (\Delta e)^2 \left(\frac{1}{2a_1} + \frac{1}{2a_2} - \frac{1}{R} \right) \left(\frac{1}{D_{op}} - \frac{1}{D_s} \right) \quad (7)$$

For a bimolecular reaction, work terms, principally electrostatic, are involved in bringing the reactants together and in separating the reaction products, but are omitted from Eq. (5) for notational brevity. The expression for the vibrational term λ_i is given by

$$\lambda_i = \frac{1}{2} \sum_j k_j (Q_j^r - Q_j^p)^2 \quad (8)$$

where Q_j^r and Q_j^p are equilibrium values for the j th normal mode coordinate Q , and k_j is a reduced force constant $2k_j^r k_j^p / (k_j^r + k_j^p)$ k_j^r being the force constant for the reactants and k_j^p being that for the products. (I introduced a “symmetrization” approximation for the vibrational part of the potential energy surface, to obtain this simple form of Eqs. (5) to (8), and tested it numerically.)

In 1957 I published the results of a calculation of the λ_i arising from a stretching vibration in the innermost coordination shell of each reactant, (the equation used for λ_i was given in the 1960 paper). An early paper on the purely vibrational contribution using chemical bond length coordinates and neglecting bond-bond correlation had already been published for self-exchange reactions by George and Griffiths in 1956.

I also extended the theory to treat electron transfers at electrodes, and distributed it as an Office of Naval Research Report in 1957, the equations being published later in a journal paper in 1959. I had little prior knowledge of the subject, and my work on electrochemical electron transfers was facilitated considerably by reading a beautiful and logically written survey article of Roger Parsons on the equilibrium electrostatic properties of electrified metal-solution interfaces.

In the 1957 and 1965 work I showed that the electrochemical rate constant was again given by Eqs. (5)–(8), but with A now having a value appropriate to the different “geometry” of the encounter of the participants in the reaction. The $1/2a_2$ in Eq. (7) was now absent (there is only one reacting ion) and R now denotes twice the distance from the center of the reactant’s charge to the electrode (it equals the ion-image distance). A term $e\eta$ replaced the ΔG° in Eq.(5b), where e is the charge transferred between the ion and the electrode, and η is the activation overpotential, namely the metal-solution potential difference, relative to the value it would have if the rate constants for the forward and reverse reactions were equal. These rate constants are equal when the minima of the two G curves in Figure 5 have the same height.

Free Energy Curves




Figure 5. Free energy of reactants plus environment vs. the reaction coordinate q (R curve), and free energy of products plus environment vs. reaction coordinate q (P curve). The three vertical lines on the abscissa denote, from left to right, the value for the reactants, for the transition state, and for the products. (Marcus and Siddarth, ref. 2).

When $|e\eta| < \lambda$, most electrons go into or out of quantum states in the metal that are near the Fermi level. However, because of the continuum of states in the metal, the inverted effect (described below) was now predicted to be absent for this process, i.e., the counterpart of Eq. (5b) is applicable only in the region $|e\eta| < \lambda$: In the case of an intrinsically highly exothermic electron transfer reaction at an electrode, the electron can remove the immediate “exothermicity” by (if entering) going into a high unoccupied quantum state of the metal, or (if leaving) departing from a low occupied quantum state, each far removed from the Fermi level. (The inverted region effect should, however, occur for the electron transfer when the electrode is a narrow band semiconductor.)

After these initial electron transfer studies, which were based on a dielectric continuum approximation for the solvent outside the first coordination shell of each reactant, I introduced a purely molecular treatment of the reacting system. Using statistical mechanics, the solvent was treated as a collection of dipoles in the 1960 paper, and later in 1965 a general charge distribution was used for the solvent molecules and for the reactants. At the same time I found a way in this 1960 paper of introducing rigorously a

global reaction coordinate in this many-dimensional (N) coordinate space of the reacting system. The globally defined coordinate so introduced was equivalent to using $U_p - U_r$, the potential energy difference between the products plus solvent (U_p) and the reactants plus solvent (U_r) (cf A. Warshel, 1987). It was, thereby, a coordinate defined everywhere in this N -dimensional space.

The free energy G_r of a system containing the solvent and the reactants, and that of the corresponding system for the products, G_p , could now be defined along this globally defined reaction coordinate. (In contrast, in reactions such as that depicted by Figure 2, it is customary, instead, to define a reaction coordinate locally, namely, in the vicinity of a path leading from the valley of the reactants through the saddle point region and into the valley of the products.)

The potential energies U_r and U_p in the many-dimensional coordinate space are simple functions of the vibrational coordinates but are complicated functions of the hundreds of relevant solvent coordinates: there are many local minima corresponding to locally stable arrangements of the solvent molecules. However, I introduced a "linear response approximation," in which any hypothetical change in charge of the reactants produces a proportional change in the dielectric polarization of the solvent. (Recently, I utilized a central limit theorem to understand this approximation better—beyond simple perturbation theory, and plan to submit the results for publication shortly.) With this linear approximation the free energies G_r and G_p became simple quadratic functions of the reaction coordinate.

Such an approach had major consequences. This picture permitted a depiction of the reaction in terms of parabolic free energy plots in simple and readily visualized terms, as in Figure 5. With them the trends predicted from the equations were readily understood. It was also important to use the free energy curves, instead of oversimplified potential energy profiles, because of the large entropy changes which occur in many electron transfer cross-reactions, due to changes in strong ion-polar solvent interactions. (The free energy plot is legitimately a one-coordinate plot while the potential energy plot is at most a profile of the complicated U_r and U_p in N -dimensional space.)

With the new statistical mechanical treatment of 1960 and 1965 one could also see how certain relations between rate constants initially derivable from the dielectric continuum-based equations in the 1956 paper could also be valid more generally. The relations were based, in part, on Equations (5) and (initially via (7) and (8)) on the approximate relation

$$\lambda_{12} \cong \frac{1}{2} (\lambda_{11} + \lambda_{22}) \quad (9)$$

where λ_{12} is the λ for the cross-reaction and the λ_{11} and λ_{22} are those of the self-exchange reactions.

Predictions

In the 1960 paper I had listed a number of theoretical predictions resulting from these equations, in part to stimulate discussion with experimentalists in the field at a Faraday Society meeting on oxidation-reduction reactions, where this paper was to be presented. At the time I certainly did not anticipate the subsequent involvement of the many experimentalists in testing these predictions. Among the latter was one which became one of the most widely tested aspects of the theory, namely, the “cross-relation.” This expression, which follows from Eqs. (5) and (9), relates the rate constant k_{12} of a cross-reaction to the two self-exchange rate constants, k_{11} and k_{22} , and to the equilibrium constant K_{12} of the reaction.

$$k_{12} \cong (k_{11}k_{22}K_{12}f_{12})^{1/2}, \quad (10)$$

where f_{12} is a known function of k_{11} , k_{22} and K_{12} and is usually close to unity.

Another prediction in the 1960 paper concerned what I termed there the inverted region: In a series of related reactions, similar in λ but differing in ΔG° , a plot of the activation free energy ΔG^* vs. ΔG° is seen from Eq. (5b) to first decrease as ΔG° is varied from 0 to some negative value, vanish at $\Delta G^\circ = -\lambda$, and then increase when ΔG° is made still more negative. This initial decrease of ΔG^* with increasingly negative ΔG° is the expected trend in chemical reactions and is similar to the usual trend in “Bronsted plots” of acid or base catalyzed reactions and in “Tafel plots” of electrochemical reactions. I termed that region of ΔG° the “normal” region. However, the prediction for the region where $-\Delta G^\circ > \lambda$, the “inverted region,” was the unexpected behavior, or at least unexpected until the present theory was introduced.

This inverted region is also easily visualized using Figures 6 and 7: Successively making ΔG° more negative, by lowering the products’ G curve vertically relative to the reactant curve, decreases the free energy barrier ΔG^* (given by the intersection of the reactants’ and products’ curves): that barrier is seen in Figure 6 to vanish at some ΔG° and then to increase again.

Other predictions dealt with the relation between the electrochemical and the corresponding self-exchange electron transfer rates, the numerical estimate of the reaction rate constant k and, in the case of non-specific solvent effects, the dependence of the reaction rate on solvent dielectric properties. The testing of some of the predictions was delayed by an extended sabbatical in 1960–61, which I spent auditing courses and attending seminars at the nearby Courant Mathematical Institute.

Comparisons of Experiment and Theory

Around 1962, during one of my visits to Brookhaven National Laboratory, I showed Norman Sutin the 1960 predictions. Norman had either measured via his stopped-flow apparatus or otherwise knew rate constants and equilibrium constants which permitted the cross-relation Eq. (10) to be tested. There were about six such sets of data which he had available. I remember vividly the growing sense of excitement we both felt as, one by one, the

The Inverted Region Effect




Figure 6. Plot of the free energy G versus the reaction coordinate q , for reactants' (R) and products' (P), for three different values of ΔG^0 , the cases I to III indicated in Figure 7 (Marcus and Siddarth, ref. 2).

The Inverted Region Effect




Figure 7. Plot of $\ln k_r$ vs $-\Delta G^0$. Points I and III are in the normal and inverted regions, respectively, while point II, where $\ln k_r$ is a maximum, occurs at $-\Delta G^0 = \lambda$ (Marcus and Siddarth, ref. 2).

observed k_{12} 's more or less agreed with the predictions of the relation. I later collected the results of this and of various other tests of the 1960 predictions and published them in 1963. Perhaps by showing that the previously published expressions were not mere abstract formulae, but rather had concrete applications, this 1963 paper, and many tests by Sutin and others, appear to have stimulated numerous subsequent tests of the cross-relation and of the other predictions. A few examples of the cross-relation test are given in Table I.

The encouraging success of the experimental tests given in the 1963 paper suggested that the theory itself was more general than the approximations (e.g., solvent dipoles, unchanged force constants) used in 1960 and stimulated me to give a more general formulation (1965). The latter paper also contains a unified treatment of electron transfers in solution and at metal electrodes, and served, thereby, to generalize my earlier (1957) treatment of the electrochemical electron transfers.

The best experimental evidence for the inverted region was provided in 1984 by Miller, Calcaterra and Closs, almost 25 years after it was predicted.

Experimental Confirmation of Inverted Region




Figure 8. Inverted region effect in chemical electron transfer reactions. (Miller, *et al*, ref. 3).

This successful experimental test, which was later obtained for other electron transfer reactions in other laboratories, is reproduced in Figure 8. Possible reasons for not observing it in the earlier tests are several-fold and have been discussed elsewhere.

Previously, indirect evidence for the inverted region had been obtained by observing that electron transfer reactions with a very negative ΔG^* may result in chemiluminescence: when the G_r and G_p curves intersect at a high ΔG^* because of the inverted region effect, there may be an electron transfer to a more easily accessible G_p curve, one in which one of the products is electronically excited and which intersects the G_r curve in the normal region at a low ΔG^* , as in Figure 9. Indeed, experimentally in some reactions 100 % formation of an electronically excited state of a reaction product has been observed by Bard and coworkers, and results in chemiluminescence.

Another consequence of Eq. (5) is the linear dependence of $k_B T \ln k$ on $-\Delta G^o$ with a slope of 1/2, when $|\Delta G^o/\lambda|$ is small, and a similar behavior at electrodes, with ΔG^o replaced by $e\eta$ the product of the charge transferred

Formation of Electronically Excited Products




Figure 9. A favored formation of an electronically excited state of the products (Marcus and Siddarth, ref. 2).

and the activation overpotential. Extensive verification of both these results has been obtained. More recently, the curvature of plots of $\ln k$ vs. $e\eta$, expected from these equations, has been demonstrated in several experiments. The very recent use of ordered organic molecular monolayers on electrodes, either to slow down the electron transfer rate or to bind a redox-active agent to the electrode, but in either case to avoid or minimize diffusion control of the fast electron transfer processes, has considerably facilitated this study of the curvature in the $\ln k$ vs. $e\eta$ plot.

Comparison of experiment and theory has also included that of the absolute reaction rates of the self-exchange reactions, the effect on the rate of varying the solvent, an effect sometimes complicated by ion pairing in the low dielectric constant media involved, and studies of the related problem of charge transfer spectra, such as



Here, the frequency of the spectral absorption maximum $\nu_{\max}$ is given

$$h\nu_{\max} = \lambda + \Delta G^0. \tag{12}$$

Comparisons with Eq. (12), using Eq.(7) for λ , have included those of the effects of separation distance and of the solvent dielectric constant.

Comparisons have also been made of the self-exchange reaction rates in solution with the rates of the corresponding electron transfer reactions at

Electrochemical vs Self-Exchange Rate Constants




Figure 10. Comparison of isotopic exchange electron transfer rates in solution, covering 20 orders of magnitude, with rates of corresponding electron transfers at metal electrodes. (Cannon, ref. 2).

electrodes. An example of the latter is the plot given in Figure 10, where the self-exchange rates are seen to vary by some twenty orders of magnitude. The discrepancy at high k 's is currently the subject of some reinvestigation of the fast electrode reaction rates, using the new nanotechnology. Most recently, a new type of interfacial electron transfer rate has also been measured, electron transfer at liquid-liquid interfaces. In treating the latter, I extended the "cross relation" to this two-phase system. It is clear that much is to be learned from this new area of investigation. (The study of the transfer of ions across such an interface, on the other hand, goes back to the time of Nernst and of Planck, around the turn of the century.)

Other Applications and Extensions

As noted in Figure 1, one aspect of the electron transfer field has been its continued and, indeed, ever-expanding growth in so many directions. One of these is in the biological field, where there are now detailed experimental and theoretical studies in photosynthetic and other protein systems. The three-dimensional structure of a photosynthetic reaction center, the first membrane protein to be so characterized, was obtained by Deisenhofer, Michel and Huber, who received the Nobel Prize in Chemistry in 1988 for this work. A bacterial photosynthetic system is depicted in Figure 11, where the protein framework holding fast the constituents in this reaction center is not shown.

The Reaction Center




Figure 11. Redox-active species involved in the initial charge separation for a photosynthetic bacterium (cf Deisenhofer *et al*, ref. 3, and Yeates *et al*, ref. 3), with labels added, to conform to the present text; they include a missing Q_B .

In the photosynthetic system there is a transfer of electronic excitation from “antenna” chlorophylls (not shown in Figure 11) to a special pair $BChl_2$. The latter then transfers an electron to a pheophytin BPh within a very short time (~ 3 picoseconds) and from it to a quinone Q_A in 200 psec and thence to the other quinone Q_B . (Other chemical reactions then occur with these separated charges at each side of the membrane, bridged by this photosynthetic reaction center.)

To avoid wasting the excitation energy of the $BChl_2^*$ unduly it is necessary that the $-\Delta G^\circ$ of this first electron transfer to BPh be small. (It is only about 0.25 eV out of an overall excitation energy of $BChl_2^*$ of 1.38 eV.) In order that this electron transfer also be successful in competing with two wasteful processes, the fluorescence and the radiationless transition of $BChl_2^*$, it is also necessary that ΔG^* for that first electron transfer step be small and hence, by Eq. (5b), that the λ be small. The size of the reactants is large, and the immediate protein environment is largely nonpolar, so leading to a small λ (cf Eq.(7)). Nature appears, indeed, to have constructed a system with this desirable property.

Furthermore, to avoid another form of wasting the energy, it is also important that an unwanted back electron transfer reaction from the BPh^- to the $BChl_2^+$ not compete successfully with a second forward electron transfer step from BPh^- to Q_A . That is, it is necessary that the back transfer, a “hole-electron recombination” step, be slow, even though it is a very highly exothermic process (~ 1.1 eV). It has been suggested that the small λ (~ 0.25 eV) and the resulting inverted region effect play a significant role in providing this essential condition for the effectiveness of the photosynthetic reaction center.

There is now a widespread interest in synthesizing systems which can mimic the behavior of nature’s photosynthetic systems, and so offer other routes for the harnessing of solar energy. The current understanding of how nature works has served to provide some guidelines. In this context, as well as that of electron transfer in other proteins, there are also relevant experiments in long range electron transfer. Originally the studies were of electron transfer in rigid glasses and were due to Miller and coworkers. More recently the studies have involved a donor and receptor held together by synthetically made rigid molecular bridges. The effect of varying the bridge length has been studied in the various systems. A theoretical estimate of the distance dependence of electron transfers in a photosynthetic system was first made by Hopfield, who used a square barrier model and an approximate molecular estimate of the barrier height.

Recently, in their studies of long range electron transfer in chemically modified proteins, Gray and coworkers have studied systematically the distance or site dependence of the electronic factor, by attaching an appropriate electron donor or acceptor to a desired site. For each such site the reactant chosen should be such that $-\Delta G^\circ \approx \lambda$, i.e., which has a k at the maximum of the $\ln k$ vs. $-\Delta G^\circ$ curve (cf Eq. (5)). The value of k then no longer depends on a ΔG^* . Since ΔG^* is distance-dependent (cf Eq.(7)), it is

particularly desirable to make $\Delta G^* \approx 0$, so that the relative k 's at the various sites now reflect only the electronic factor. Dutton and coworkers have treated data similarly for a number of reactions by using, where possible, the k at the maximum of each $\ln k$ vs. ΔG° curve. Of particular interest in such studies is whether there is a simple exponential decrease of the electronic factor on the separation distance between donor and acceptor, or whether there are deviations from this monotonic behavior, due to local structural factors.

In a different development, the mechanism of various organic reactions has been explored by several investigators, notably by Eberson (ref. 2), in the light of current electron transfer theory. Other organic reactions have been explored by Shaik and Pross, in their analysis of a possible electron transfer mechanism vs. a conventional mechanism, and by Shaik *et al* (ref.2).

Theoretical calculations of the donor-acceptor electronic interactions, initially by McConnell and by Larsson, and later by others, our group among them, have been used to treat long-range electron transfer. The methods have recently been adapted to large protein systems. In our studies with Siddarth we used an "artificial intelligence" searching technique to limit the number of amino acids used in the latter type of study.

Another area of much current activity in electron transfers is that of solvent dynamics, following the pioneering treatment for general reactions by Kramers (1940). Important later developments for electron transfer were made by many contributors. Solvent dynamics affects the electron transfer reaction rate when the solvent is sufficiently sluggish. As we showed recently with Sumi and Nadler, the solvent dynamics effect can also be modified significantly, when there are vibrational (λ_i) contribution to λ .

Computational studies, such as the insightful one of David Chandler and coworkers on the $\text{Fe}^{2+} + \text{Fe}^{3+}$ self-exchange reaction, have also been employed recently. Using computer simulations they obtained a verification of the parabolic G curves, even for surprisingly high values of the fluctuation in G . They also extended their studies to dynamical and quantum mechanical effects of the nuclear motion. Studies of the quantum mechanical effects on the nuclear motion on electron transfer reactions were initiated in 1959 by Levich and Dogonadze, who assumed a harmonic oscillator model for the polar solvent medium and employed perturbation theory. Their method was related to that used for other problems by Huang and Rhys (1951) and Kubo and Toyozawa (1954).

There were important subsequent developments by various authors on these quantum effects, including the first discussion of quantum effects for the vibrations of the reactants by Sutin in 1962 and the important work of Jortner and coworkers in 1974–75, who combined a Levich and Dogonadze type approach to treat the high frequency vibrations of the reactants with the classical expression which I described earlier for the polar medium. These quantum effects have implications for the temperature dependence of k , among other effects. Proceeding in a different (classical) direction Saveant recently showed how to extend Eq. (5b) to reactions which involved

the rupture of a chemical bond by electron transfer and which he had previously studied experimentally: $M(e) + RX \rightarrow M + R + X^-$, where R is an alkyl group, X a halide and M a metal electrode.

A particularly important early development was that by Taube in the 1950s; he received the Nobel Prize for his work in 1983. Taube introduced the idea of different mechanisms for electron transfer—outer sphere and inner sphere electron transfers, which he had investigated experimentally. His experimental work on charge transfer spectra of strongly interacting systems ("Creutz-Taube" ion, 1959, 1973) and of weakly interacting ones has been similarly influential. Also notable has been Hush's theoretical work on charge transfer spectra, both of intensities and absorption maxima (1967), which supplemented his earlier theoretical study of electron transfer rates (1961).

There has been a "spin-off" of the original electron transfer theory to other types of chemical reactions as well. In particular, the ΔG^* vs ΔG^0 relation and the cross-relation have been extended to these other reactions, such as the transfer of atoms, protons, or methyl groups. (Even an analog of Eqs. (5b) and (9), but for binding energies instead of energy barriers has been introduced to relate the stability of isolated protonbound dimers AHB^+ to those of AHA^+ and BHB^+ !)

Since the transfer of these nuclei involves strong electronic interactions, it is not well represented by intersecting parabolic free energy curves, and so a different theoretical approach was needed. For this purpose I adapted (1968) a "bond-energy-bond-order" model of H. Johnston, in order to treat the problem for a reaction of the type given by Eq.(4). The resulting simple expression for ΔG^* is similar to Eq.(5), when $|\Delta G^0/\lambda|$ is not large ($< 1/2$), but differs from it in not having any inverted region. It has the same λ property as that given by Eq.(9), and has resulted in a cross-relation analogous to Eq. (10). The cross-relation has been tested experimentally for the transfer of methyl groups by E. Lewis, and the ΔG^* vs ΔG^0 relation has been used or tested for other transfers by Alberty and by Kreevoy and their coworkers, among others.

It is naturally gratifying to see one's theories used. A recent article, which showed the considerable growth in the use of papers such as the 1956 and 1964 articles (ref. 5), points up the impressive and continued vitality of the field itself. The remarks above on many areas of electron transfer and on the spin-off of such work on the study of other types of reactions represent a necessarily brief picture of these broad-based investigations.

ACKNOWLEDGMENTS

My acknowledgments are to my many fellow researchers in the electron transfer field, notably Norman Sutin, with whom I have discussed so many of these matters for the past thirty or more years. I also thank my students and post-doctorals, whose presence was a constant source of stimulation to me, both in the electron transfer field and in the other fields of research

Table I. Comparison of Calculated and Experimental k_{12} Values^a

Reaction	$k_{12}, M^{-1} \text{ sec}^{-1}$	
	Observed	Calculated
$\text{IrCl}_6^{3-} + \text{W}(\text{CN})_8^{4-}$	6.1×10^7	6.1×10^7
$\text{IrCl}_6^{3-} + \text{Fe}(\text{CN})_6^{4-}$	3.8×10^6	7×10^6
$\text{IrCl}_6^{3-} + \text{Mo}(\text{CN})_8^{4-}$	1.9×10^6	9×10^6
$\text{Mo}(\text{CN})_8^{3-} + \text{W}(\text{CN})_8^{4-}$	5.0×10^6	4.8×10^6
$\text{Mo}(\text{CN})_8^{3-} + \text{Fe}(\text{CN})_6^{4-}$	3.0×10^4	2.9×10^4
$\text{Fe}(\text{CN})_6^{3-} + \text{W}(\text{CN})_8^{4-}$	4.3×10^4	6.3×10^4
$\text{Ce}^{\text{IV}} + \text{W}(\text{CN})_8^{4-}$	$>10^3$	4×10^3
$\text{Ce}^{\text{IV}} + \text{Fe}(\text{CN})_6^{4-}$	1.9×10^6	8×10^6
$\text{Ce}^{\text{IV}} + \text{Mo}(\text{CN})_8^{4-}$	1.4×10^7	1.3×10^7
$\text{L-Co}[(\text{--})\text{PDTA}]^{2-} + \text{Fe}(\text{bipy})_3^{3+}$	8.1×10^4	$\geq 10^5$
$\text{L-Fe}[(\text{--})\text{PDTA}]^{2-} + \text{Co}(\text{EDTA})^-$	1.3×10^1	1.3×10^1
$\text{L-Fe}[(\text{--})\text{PDTA}]^{2-} + \text{Co}(\text{ox})_3^{3-}$	2.2×10^3	1.0×10^3
$\text{Cr}(\text{EDTA})^{2-} + \text{Fe}(\text{EDTA})^-$	$\geq 10^6$	10^9
$\text{Cr}(\text{EDTA})^{2-} + \text{Co}(\text{EDTA})^-$	$\simeq 3 \times 10^6$	4×10^7
$\text{Fe}(\text{EDTA})^{2-} + \text{Mn}(\text{CyDTA})^-$	$\simeq 4 \times 10^6$	6×10^6
$\text{Co}(\text{EDTA})^{2-} + \text{Mn}(\text{CyDTA})$	9×10^{-1}	2.1
$\text{Fe}(\text{PDTA})^{2-} + \text{Co}(\text{CyDTA})^-$	1.2×10^1	1.8×10^1
$\text{Co}(\text{terpy})_2^{2+} + \text{Co}(\text{bipy})_3^{3+}$	6.4×10	3.2×10
$\text{Co}(\text{terpy})_2^{2+} + \text{Co}(\text{phen})_3^{3+}$	2.8×10^3	1.1×10^3
$\text{Co}(\text{terpy})_2^{2+} + \text{Co}(\text{bipy})(\text{H}_2\text{O})_4^{3+}$	6.8×10^3	6.4×10^4
$\text{Co}(\text{terpy})_2^{2+} + \text{Co}(\text{phen})(\text{H}_2\text{O})_4^{3+}$	1.4×10^3	6.4×10^4
$\text{Co}(\text{terpy})_2^{2+} + \text{Co}(\text{H}_2\text{O})_6^{3+}$	7.4×10^4	2×10^{10}
$\text{Fe}(\text{phen})_3^{2+} + \text{MnO}_4^-$	6×10^3	4×10^3
$\text{Fe}(\text{CN})_6^{4-} + \text{MnO}_4^-$	1.3×10^4	5×10^3
$\text{V}(\text{H}_2\text{O})_6^{2+} + \text{Ru}(\text{NH}_3)_6^{3+}$	1.5×10^3 ^a	4.2×10^3
$\text{Ru}(\text{en})_3^{2+} + \text{Fe}(\text{H}_2\text{O})_6^{3+}$	8.4×10^4	4.2×10^6
$\text{Ru}(\text{NH}_3)_6^{2+} + \text{Fe}(\text{H}_2\text{O})_6^{3+}$	3.4×10^6	7.5×10^6
$\text{Fe}(\text{H}_2\text{O})_6^{2+} + \text{Mn}(\text{H}_2\text{O})_6^{3+}$	1.5×10^4	3×10^4

^a Bennett, ref. 3.

which we have explored. In its earliest stage and for much of this period this research was supported by the Office of Naval Research and also later by the National Science Foundation. The support of both agencies continues to this day and I am very pleased to acknowledge its value and timeliness here.

In my Nobel lecture, I concluded on a personal note with a slide of my great uncle, Henrik Steen (né Markus), who came to Sweden in 1892. He received his doctorate in theology from the University of Uppsala in 1915, and was an educator and a prolific writer of pedagogic books. As I noted in the biographical sketch in *Les Prix Nobel*, he was one of my childhood idols. Coming here, visiting with my Swedish relatives—some thirty or so descendants—has been an especially heartwarming experience for me and for my family. In a sense I feel that I owed him a debt, and that it is most fitting to acknowledge that debt here.

REFERENCES

Some of my relevant articles, largely from the 1956–65 period, are listed in ref. 1 below, and some general references which review the overall literature are listed in ref. 2. Several additional references for the Table and for the Figures are given in ref. 3. Classic texts on unimolecular reactions are given in ref. 4.

1. R. A. Marcus, J. Chem. Phys. **24**, 966 (1956); *ibid.*, **24**, 979 (1956); *ibid.*, **26**, 867 (1957); *ibid.*, **26**, 872 (1957); Trans. N. Y. Acad. Sci. **19**, 423 (1957); ONR Technical Report No. 12, Project NR 051-331 (1957), reproduced in *Special Topics in Electrochemistry*, P. A. Rock, ed., Elsevier, New York, 1977, p. 181; Can. J. Chem. **37**, 155 (1959); Discussions Faraday Soc. **29**, 21 (1960); J. Phys. Chem. **67**, 853, 2889 (1963); J. Chem. Phys. **38**, 1858 (1963); *ibid.*, **39**, 1734 (1963); Ann. Rev. Phys. Chem. **15**, 155 (1964); J. Chem. Phys. **43**, 679 (1965); *ibid.*, **43**, 1261 (1965); *ibid.*, **43**, 2654 (1965), (corr.) **52**, 2803 (1970); J. Phys. Chem. **72**, 891 (1968)
2. R. A. Marcus and N. Sutin, Biochim. Biophys. Acta, **811**, 265 (1985); J.R. Bolton, N. Mataga and G. McLendon (eds.) Adv. Chem. Ser., **228**, (1991), assorted articles; M.D. Newton and N. Sutin, Ann. Rev. Phys. Chem., **35**, 437 (1984); N. Sutin, Prog. Inorg. Chem., **30**, 441 (1983); M.D. Newton, Chem. Rev., **91**, 767 (1991); R.D. Cannon, *Electron Transfer Reactions*, Butterworths, London, 1980; L. Ebersson, *Electron Transfer Reactions in Organic Chemistry*, Springer, New York, 1987; M.A. Fox and M. Chanon (eds.), *Photoinduced Electron Transfer*, Elsevier, New York, 1988, 4 vols.; M.V. Twigg (ed.), *Mechanisms of Inorganic and Organometallic Reactions*, vol. 7, 1991, Chaps. 1 and 2, and earlier volumes; R. A. Marcus and P. Siddarth, in *Photoprocesses in Transition Metal Complexes, Biosystems and Other Molecules: Experiment and Theory*, E. Kochanski, ed., Kluwer, Norwall, Massachusetts, 1992, p. 49; S.S. Shaik, H.B. Schlegel and S. Wolfe, *Theoretical Aspects of Physical Organic Chemistry*, J. Wiley, New York, 1992; N. Sutin, Pure & Applied Chem. **60**, 1817 (1988); Assorted articles in Chem. Revs. **92**, No. 3 (1992); R.A. Marcus Commemorative Issue, J. Phys. Chem. **90**, (1986).
3. L. E. Bennett, Prog. Inorg. Chem. **18**, 1 (1973); J.R. Miller, L.T. Calcaterra and G.L. Closs, J. Am. Chem. Soc. **106**, 3047 (1984); J. Deisenhofer, O. Epp, K. Miki, R. Huber and H. Michel, J. Mol. Biol. **180**, 385 (1984); J. Deisenhofer and H. Michel, Angew. Chem. Int. Ed. Engl. **28**, 829 (1989); T.G. Yeates, H. Komiya, D.C. Rees, J.P. Allen and G. Feher, Proc. Nat. Acad. Sci. **84**, 6438 (1987).
4. P. J. Robinson and H.A. Holbrook, *Unimolecular Reactions*, J. Wiley, New York 1972; W. Forst, *Theory of Unimolecular Reactions*, Academic Press, New York, 1973; cf also the very recent text, R.G. Gilbert and S.C. Smith, *Theory of Unimolecular and Recombination Reactions*, Blackwells, Oxford, 1990.
5. Science Watch, **3**, No.9, November, 1992, p.8.